



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 19, 2026 – 06:55 PM JST

PDB ID : 26PJ / pdb_000026pj
Title : X-ray crystal structure of R28D mutant of the human NBS1 FHA-BRCT1-BRCT2 domain
Authors : Ite, K.; Unno, M.
Deposited on : 2026-05-09
Resolution : 2.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1520 (2.86-2.82)
Clashscore	190562	1559 (2.86-2.82)
Ramachandran outliers	187476	1517 (2.86-2.82)
Sidechain outliers	187428	1518 (2.86-2.82)
RSRZ outliers	180081	1521 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	4% 73% 19% 6% .
1	B	334	5% 74% 16% 6% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5085 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nibrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	328	Total	C	N	O	S	0	0	0
			2525	1606	424	480	15			
1	B	322	Total	C	N	O	S	0	0	0
			2485	1585	413	472	15			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	28	ASP	ARG	engineered mutation	UNP O60934
A	327	LEU	-	expression tag	UNP O60934
A	328	GLU	-	expression tag	UNP O60934
A	329	HIS	-	expression tag	UNP O60934
A	330	HIS	-	expression tag	UNP O60934
A	331	HIS	-	expression tag	UNP O60934
A	332	HIS	-	expression tag	UNP O60934
A	333	HIS	-	expression tag	UNP O60934
A	334	HIS	-	expression tag	UNP O60934
B	28	ASP	ARG	engineered mutation	UNP O60934
B	327	LEU	-	expression tag	UNP O60934
B	328	GLU	-	expression tag	UNP O60934
B	329	HIS	-	expression tag	UNP O60934
B	330	HIS	-	expression tag	UNP O60934
B	331	HIS	-	expression tag	UNP O60934
B	332	HIS	-	expression tag	UNP O60934
B	333	HIS	-	expression tag	UNP O60934
B	334	HIS	-	expression tag	UNP O60934

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	32	Total 32	O 32	0	0
4	B	24	Total 24	O 24	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nibrin



• Molecule 1: Nibrin



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	157.72Å 182.38Å 122.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.45 – 2.84 48.45 – 2.84	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.45-2.84) 99.9 (48.45-2.84)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 2.86Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, R_{free}	0.184 , 0.224 0.189 , 0.221	Depositor DCC
R_{free} test set	2083 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	81.3	Xtriage
Anisotropy	0.597	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 59.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5085	wwPDB-VP
Average B, all atoms (Å ²)	92.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/2575	1.34	18/3487 (0.5%)
1	B	0.65	0/2532	1.29	8/3424 (0.2%)
All	All	0.66	0/5107	1.32	26/6911 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	244	PHE	CA-CB-CG	7.99	121.79	113.80
1	A	244	PHE	CA-CB-CG	7.97	121.77	113.80
1	A	260	HIS	CA-CB-CG	7.90	121.70	113.80
1	A	233	LYS	CB-CA-C	7.19	122.16	110.88
1	A	228	ILE	N-CA-CB	6.95	118.29	110.72
1	A	139	THR	CA-CB-OG1	-6.59	99.72	109.60
1	A	253	THR	CA-CB-OG1	-6.54	99.79	109.60
1	B	243	VAL	N-CA-CB	6.46	118.84	110.57
1	B	253	THR	CA-CB-OG1	-6.26	100.21	109.60
1	A	202	GLU	CB-CA-C	5.89	116.86	110.08
1	A	320	LYS	N-CA-CB	5.84	118.78	110.13
1	A	322	TYR	CB-CA-C	5.73	116.85	109.80
1	A	317	MET	CG-SD-CE	-5.71	88.34	100.90
1	A	61	ASP	CA-CB-CG	5.70	118.30	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	139	THR	CA-CB-OG1	-5.66	101.11	109.60
1	B	6	PRO	CA-C-O	-5.41	114.99	121.48
1	A	23	GLU	CB-CA-C	-5.40	101.44	110.19
1	A	34	LEU	N-CA-CB	-5.40	101.62	110.42
1	A	177	PHE	CA-CB-CG	-5.29	108.51	113.80
1	A	84	GLN	N-CA-CB	5.21	118.68	110.23
1	B	23	GLU	CB-CA-C	-5.18	101.80	110.19
1	A	318	THR	CA-CB-OG1	-5.12	101.92	109.60
1	B	257	GLU	CB-CA-C	5.10	120.57	110.42
1	A	6	PRO	CA-C-O	-5.08	115.39	121.48
1	A	273	THR	CA-CB-OG1	-5.07	102.00	109.60
1	B	219	LYS	CB-CG-CD	5.01	122.81	111.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	103	GLY	Peptide
1	B	89	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2525	0	2516	42	0
1	B	2485	0	2486	45	0
2	A	10	0	0	0	0
2	B	5	0	0	0	0
3	A	4	0	6	0	0
4	A	32	0	0	2	0
4	B	24	0	0	0	0
All	All	5085	0	5008	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:VAL:HG22	1:A:147:CYS:HA	1.39	1.01
1:A:275:ILE:HA	1:A:278:SER:OG	1.83	0.79
1:A:217:GLU:HB2	1:A:317:MET:CE	2.16	0.76
1:A:264:LEU:O	1:A:265:ALA:O	2.05	0.74
1:A:114:VAL:CG2	1:A:147:CYS:HA	2.20	0.71
1:B:269:CYS:SG	1:B:305:ILE:CD1	2.82	0.67
1:B:298:GLN:HE21	1:B:298:GLN:CA	2.08	0.67
1:B:103:GLY:HA2	1:B:105:LYS:HD2	1.77	0.65
1:A:202:GLU:OE2	1:A:204:SER:HB3	1.98	0.63
1:A:264:LEU:HD21	1:A:300:GLN:HG3	1.83	0.61
1:B:298:GLN:HE21	1:B:298:GLN:HA	1.66	0.61
1:A:217:GLU:HB2	1:A:317:MET:HE3	1.84	0.60
1:B:269:CYS:HB2	1:B:305:ILE:CD1	2.32	0.59
1:A:63:ILE:HD12	1:A:63:ILE:O	2.03	0.59
1:A:233:LYS:HE2	1:A:234:GLN:N	2.18	0.58
1:B:103:GLY:CA	1:B:105:LYS:HD2	2.34	0.57
1:A:231:ASN:OD1	1:A:231:ASN:C	2.47	0.57
1:B:231:ASN:OD1	1:B:231:ASN:C	2.48	0.57
1:B:269:CYS:HB2	1:B:305:ILE:HD11	1.88	0.54
1:B:45:HIS:CD2	1:B:76:THR:HG22	2.42	0.54
1:B:25:VAL:HG13	1:B:31:CYS:HB2	1.89	0.53
1:B:144:THR:OG1	1:B:146:GLU:HG3	2.08	0.53
1:B:274:GLY:C	1:B:276:THR:H	2.16	0.53
1:A:264:LEU:HD21	1:A:300:GLN:CG	2.39	0.53
1:B:269:CYS:CB	1:B:305:ILE:HD11	2.39	0.53
1:A:17:ARG:HG2	1:A:17:ARG:NH1	2.25	0.52
1:A:264:LEU:HD21	1:A:300:GLN:CB	2.40	0.52
1:A:157:VAL:HG22	1:A:157:VAL:O	2.10	0.52
1:A:17:ARG:HG2	1:A:17:ARG:HH11	1.74	0.51
1:A:29:LYS:HE2	1:A:30:ASN:OD1	2.10	0.51
1:B:225:LYS:NZ	1:B:323:CYS:O	2.44	0.51
1:A:270:VAL:CG1	1:A:297:LEU:HD11	2.40	0.51
1:B:285:CYS:O	1:B:286:GLN:HB2	2.11	0.50
1:A:134:GLN:HG2	4:A:517:HOH:O	2.12	0.49
1:B:319:THR:O	1:B:319:THR:HG23	2.12	0.49
1:A:169:ARG:HH11	1:A:169:ARG:HG3	1.76	0.49
1:B:20:THR:O	1:B:50:ALA:O	2.29	0.49
1:B:269:CYS:CB	1:B:305:ILE:CD1	2.91	0.49
1:B:45:HIS:NE2	1:B:76:THR:HG22	2.28	0.48
1:A:156:LYS:O	1:A:161:THR:HG21	2.14	0.48
1:A:275:ILE:HA	1:A:278:SER:HG	1.75	0.48
1:B:298:GLN:HA	1:B:298:GLN:NE2	2.30	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:256:ASN:O	1:B:257:GLU:C	2.58	0.47
1:A:3:LYS:NZ	1:A:111:GLU:OE1	2.40	0.46
1:B:17:ARG:HG2	1:B:17:ARG:HH11	1.80	0.46
1:B:77:PHE:CE2	1:B:82:LYS:HB2	2.49	0.46
1:A:250:ARG:HH22	1:A:262:PHE:HE1	1.63	0.46
1:A:256:ASN:O	1:A:257:GLU:C	2.59	0.46
1:B:264:LEU:CD2	1:B:300:GLN:HG2	2.46	0.46
1:B:273:THR:C	1:B:275:ILE:H	2.24	0.46
1:A:319:THR:HG23	1:A:319:THR:O	2.16	0.45
1:A:169:ARG:HG3	1:A:169:ARG:NH1	2.32	0.45
1:A:77:PHE:CE2	1:A:82:LYS:HB2	2.52	0.45
1:B:17:ARG:HG2	1:B:17:ARG:NH1	2.32	0.45
1:B:259:GLU:C	1:B:261:ASN:H	2.23	0.45
1:B:45:HIS:HE2	1:B:76:THR:HG22	1.82	0.44
1:B:54:VAL:HG12	1:B:54:VAL:O	2.17	0.44
1:B:269:CYS:HB2	1:B:305:ILE:HD12	1.98	0.44
1:B:296:MET:O	1:B:296:MET:HG3	2.15	0.44
1:B:276:THR:HB	1:B:277:ASN:H	1.65	0.44
1:A:226:THR:HA	1:A:248:GLU:O	2.17	0.44
1:A:254:GLU:HB2	1:A:289:TRP:CD2	2.52	0.44
1:B:263:PHE:CE2	1:B:293:ILE:HD13	2.52	0.44
1:B:285:CYS:O	1:B:286:GLN:CB	2.65	0.44
1:B:25:VAL:HG13	1:B:31:CYS:CB	2.48	0.43
1:A:264:LEU:C	1:A:265:ALA:O	2.60	0.43
1:B:263:PHE:HE2	1:B:293:ILE:CD1	2.31	0.43
1:B:50:ALA:HA	1:B:65:VAL:O	2.17	0.43
1:A:270:VAL:HG11	1:A:297:LEU:HD11	2.00	0.43
1:B:175:GLU:OE1	1:B:175:GLU:N	2.50	0.43
1:B:210:VAL:HG12	1:B:212:LEU:HD13	2.01	0.43
1:A:105:LYS:C	1:A:106:PHE:CD1	2.97	0.43
1:B:240:SER:O	1:B:241:ALA:C	2.61	0.42
1:B:265:ALA:O	1:B:266:PRO:C	2.62	0.42
1:A:158:THR:HG22	1:A:159:ILE:H	1.84	0.42
1:A:270:VAL:HG13	1:A:297:LEU:HD11	2.01	0.42
1:A:50:ALA:HA	1:A:65:VAL:O	2.18	0.42
1:A:240:SER:O	1:A:241:ALA:C	2.61	0.42
1:B:255:GLU:O	1:B:257:GLU:N	2.50	0.42
1:A:255:GLU:O	1:A:257:GLU:N	2.50	0.42
1:A:253:THR:OG1	1:A:255:GLU:HB2	2.21	0.41
1:B:254:GLU:HB2	1:B:289:TRP:CD2	2.56	0.41
1:A:225:LYS:HE3	4:A:508:HOH:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:317:MET:HE3	1:B:317:MET:HB2	1.82	0.41
1:A:5:LEU:HD23	1:A:15:PRO:HB3	2.02	0.40
1:A:143:TRP:CD1	1:A:163:CYS:HB3	2.57	0.40
1:B:282:ILE:HA	1:B:283:PRO:HD3	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/334 (98%)	302 (93%)	20 (6%)	4 (1%)	10	22
1	B	316/334 (95%)	291 (92%)	21 (7%)	4 (1%)	9	20
All	All	642/668 (96%)	593 (92%)	41 (6%)	8 (1%)	10	22

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	10	PRO
1	A	265	ALA
1	B	275	ILE
1	A	255	GLU
1	A	256	ASN
1	B	265	ALA
1	B	54	VAL
1	B	283	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/292 (95%)	248 (89%)	30 (11%)	6	13
1	B	275/292 (94%)	243 (88%)	32 (12%)	5	11
All	All	553/584 (95%)	491 (89%)	62 (11%)	6	12

All (62) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	VAL
1	A	49	THR
1	A	62	GLU
1	A	67	THR
1	A	92	LYS
1	A	107	ARG
1	A	114	VAL
1	A	133	LEU
1	A	157	VAL
1	A	158	THR
1	A	161	THR
1	A	169	ARG
1	A	204	SER
1	A	221	ILE
1	A	226	THR
1	A	228	ILE
1	A	231	ASN
1	A	233	LYS
1	A	236	LYS
1	A	237	LYS
1	A	240	SER
1	A	243	VAL
1	A	260	HIS
1	A	261	ASN
1	A	264	LEU
1	A	268	THR
1	A	270	VAL
1	A	288	LYS
1	A	319	THR
1	A	327	LEU
1	B	49	THR
1	B	62	GLU
1	B	67	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	76	THR
1	B	125	LYS
1	B	126	THR
1	B	133	LEU
1	B	146	GLU
1	B	169	ARG
1	B	212	LEU
1	B	221	ILE
1	B	226	THR
1	B	231	ASN
1	B	243	VAL
1	B	248	GLU
1	B	255	GLU
1	B	260	HIS
1	B	261	ASN
1	B	268	THR
1	B	276	THR
1	B	281	LEU
1	B	282	ILE
1	B	285	CYS
1	B	296	MET
1	B	297	LEU
1	B	298	GLN
1	B	300	GLN
1	B	317	MET
1	B	318	THR
1	B	319	THR
1	B	320	LYS
1	B	327	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	286	GLN
1	B	44	ASN
1	B	84	GLN
1	B	220	GLN
1	B	234	GLN
1	B	261	ASN
1	B	298	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	402	-	4,4,4	0.55	0	6,6,6	0.49	0
2	PO4	B	401	-	4,4,4	0.74	0	6,6,6	0.83	0
2	PO4	A	401	-	4,4,4	1.20	0	6,6,6	0.41	0
3	EDO	A	403	-	3,3,3	0.16	0	2,2,2	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	403	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	328/334 (98%)	0.08	12 (3%) 45 35	59, 83, 151, 203	0
1	B	322/334 (96%)	0.12	16 (4%) 34 27	56, 89, 149, 188	0
All	All	650/668 (97%)	0.10	28 (4%) 40 31	56, 85, 151, 203	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	11	ALA	7.6
1	B	8	ALA	5.9
1	A	10	PRO	5.7
1	A	260	HIS	5.1
1	B	275	ILE	5.0
1	B	12	GLY	4.8
1	B	328	GLU	4.6
1	B	60	THR	4.3
1	B	9	GLY	4.3
1	A	8	ALA	3.8
1	B	244	PHE	3.5
1	B	54	VAL	3.5
1	A	328	GLU	3.3
1	B	13	GLY	3.3
1	B	225	LYS	3.1
1	B	55	THR	3.1
1	A	9	GLY	3.0
1	A	13	GLY	2.9
1	B	262	PHE	2.8
1	A	244	PHE	2.7
1	B	264	LEU	2.7
1	B	277	ASN	2.6
1	A	56	ASN	2.5
1	A	276	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	267	GLY	2.3
1	A	258	GLU	2.1
1	B	265	ALA	2.1
1	A	60	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PO4	A	402	5/5	0.71	0.10	142,156,162,175	0
2	PO4	B	401	5/5	0.93	0.14	81,85,92,98	0
3	EDO	A	403	4/4	0.94	0.23	108,109,126,132	0
2	PO4	A	401	5/5	0.95	0.10	76,80,90,123	0

6.5 Other polymers [i](#)

There are no such residues in this entry.