Full wwPDB X-ray Structure Validation Report (i)

Jun 18, 2024 – 10:38 AM EDT

PDB ID	:	4A42
Title	:	CpGH89CBM32-6 produced by Clostridium perfringens
Authors	:	Ficko-Blean, E.; Stuart, C.P.; Suits, M.D.; Cid, M.; Tessier, M.; Woods, R.J.; Boraston, A.B.
Deposited on	:	2011-10-06
Resolution	:	1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	4%, 74%, 9%, •, 15%
1	B	149	8%, 69%, 15%, •, 15%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2173 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-N-ACETYLGLUCOSAMINIDASE FAMILY PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	127	1016	630	173	211	2	9	8	0
1	B	126	981	611	167	201	2	8	8	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1473	MSE	-	expression tag	UNP Q0TST1
A	1474	GLY	-	expression tag	UNP Q0TST1
A	1475	SER	-	expression tag	UNP Q0TST1
A	1476	SER	-	expression tag	UNP Q0TST1
A	1477	HIS	-	expression tag	UNP Q0TST1
A	1478	HIS	-	expression tag	UNP Q0TST1
A	1479	HIS	-	expression tag	UNP Q0TST1
A	1480	HIS	-	expression tag	UNP Q0TST1
A	1481	HIS	-	expression tag	UNP Q0TST1
A	1482	HIS	-	expression tag	UNP Q0TST1
A	1483	SER	-	expression tag	UNP Q0TST1
A	1484	SER	-	expression tag	UNP Q0TST1
A	1485	GLY	-	expression tag	UNP Q0TST1
A	1486	LEU	-	expression tag	UNP Q0TST1
A	1487	VAL	-	expression tag	UNP Q0TST1
A	1488	PRO	-	expression tag	UNP Q0TST1
A	1489	ARG	-	expression tag	UNP Q0TST1
A	1490	GLY	-	expression tag	UNP Q0TST1
A	1491	SER	-	expression tag	UNP Q0TST1
A	1492	HIS	-	expression tag	UNP Q0TST1
A	1493	MSE	-	expression tag	UNP Q0TST1
A	1494	ALA	-	expression tag	UNP Q0TST1
A	1495	SER	-	expression tag	UNP Q0TST1
B	1473	MSE	-	expression tag	UNP Q0TST1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1474	GLY	-	expression tag	UNP Q0TST1
B	1475	SER	-	expression tag	UNP Q0TST1
B	1476	SER	-	expression tag	UNP Q0TST1
B	1477	HIS	-	expression tag	UNP Q0TST1
B	1478	HIS	-	expression tag	UNP Q0TST1
B	1479	HIS	-	expression tag	UNP Q0TST1
B	1480	HIS	-	expression tag	UNP Q0TST1
B	1481	HIS	-	expression tag	UNP Q0TST1
B	1482	HIS	-	expression tag	UNP Q0TST1
B	1483	SER	-	expression tag	UNP Q0TST1
B	1484	SER	-	expression tag	UNP Q0TST1
B	1485	GLY	-	expression tag	UNP Q0TST1
B	1486	LEU	-	expression tag	UNP Q0TST1
B	1487	VAL	-	expression tag	UNP Q0TST1
B	1488	PRO	-	expression tag	UNP Q0TST1
B	1489	ARG	-	expression tag	UNP Q0TST1
B	1490	GLY	-	expression tag	UNP Q0TST1
B	1491	SER	-	expression tag	UNP Q0TST1
B	1492	HIS	-	expression tag	UNP Q0TST1
B	1493	MSE	-	expression tag	UNP Q0TST1
B	1494	ALA	-	expression tag	UNP Q0TST1
B	1495	SER	-	expression tag	UNP Q0TST1

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
2	A	1	1	1	0	0
			Total	Ca		
2	B	1	1	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	118	118	118	0	0
			Total	O		
3	B	56	56	56	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ALPHA-N-ACETYLGLUCOSAMINIDASE FAMILY PROTEIN

Chain A:



• Molecule 1: ALPHA-N-ACETYLGLUCOSAMINIDASE FAMILY PROTEIN

Chain B:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α, β, γ	48.81Å 48.81Å 98.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.23 – 1.55 28.23 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.2 (28.23-1.55) 99.9 (28.23-1.55)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	6.11 (at 1.55Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.7_650)	Depositor
R, R_{free}	0.197 , 0.239 0.194 , 0.239	Depositor DCC
R_{free} test set	1685 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å^2)	18.1	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 55.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.047 for h,-k,-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	2173	wwPDB-VP
Average B, all atoms (Å^2)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.97% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.54	0/1045	0.62	0/1411
1	B	0.58	0/1013	0.67	0/1370
All	All	0.56	0/2058	0.64	0/2781

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1016	0	973	13	0
1	B	981	0	918	17	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	118	0	0	3	0
3	B	56	0	0	4	0
All	All	2173	0	1891	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1500[A]:MSE:HE1	3:B:2055:HOH:O	1.20	1.27
1:B:1500[A]:MSE:CE	3:B:2055:HOH:O	1.89	0.91
1:B:1584:GLU:OE1	1:B:1586[A]:LYS:HE2	1.81	0.78
1:B:1500[A]:MSE:HE2	1:B:1615:GLU:OE2	1.84	0.78
1:A:1535:LYS:HA	3:A:2023:HOH:O	1.93	0.69
1:A:1544:GLU:HB3	1:A:1598:ILE:CD1	2.26	0.65
1:B:1500[A]:MSE:CE	1:B:1521:GLY:HA2	2.28	0.63
1:B:1594:LYS:O	1:B:1595:SER:CB	2.47	0.63
1:A:1541:LEU:O	1:A:1542[B]:SER:HB2	2.01	0.60
1:B:1597:LYS:HA	3:B:2053:HOH:O	2.01	0.59
1:B:1500[A]:MSE:HE2	1:B:1521:GLY:HA2	1.83	0.59
1:A:1620:LYS:O	1:A:1621:GLY:C	2.42	0.58
1:B:1500[A]:MSE:HE3	3:B:2003:HOH:O	2.03	0.58
1:A:1506[B]:VAL:HG12	1:A:1539:PHE:CD2	2.39	0.57
1:A:1544:GLU:HB3	1:A:1598:ILE:HD12	1.86	0.56
1:B:1530:GLY:O	1:B:1535:LYS:HE2	2.07	0.54
1:B:1586[B]:LYS:HE2	1:B:1588:VAL:O	2.08	0.54
1:A:1620:LYS:O	1:A:1621:GLY:O	2.30	0.50
1:B:1584:GLU:OE1	1:B:1586[A]:LYS:CE	2.57	0.48
1:B:1497:ASN:ND2	1:B:1500[A]:MSE:HE3	2.28	0.48
1:A:1580[B]:ASN:ND2	1:A:1607:ASN:OD1	2.47	0.47
1:B:1518:LEU:HD13	1:B:1537:VAL:HG11	1.97	0.46
1:A:1580[B]:ASN:ND2	3:A:2094:HOH:O	2.50	0.45
1:B:1560:LEU:HD11	1:B:1606:PHE:CD1	2.53	0.43
1:A:1541:LEU:O	1:A:1542[B]:SER:CB	2.67	0.42
1:A:1580[B]:ASN:HD22	1:A:1580[B]:ASN:HA	1.65	0.41
1:B:1557[A]:ASN:OD1	1:B:1610:ASN:ND2	2.53	0.41
1:A:1605:ASN:HB3	3:A:2048:HOH:O	2.21	0.40
1:A:1509:THR:CG2	1:A:1536:SER:HB3	2.52	0.40
1:B:1526:LEU:HD23	1:B:1614:GLY:HA2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	133/149 (89%)	129 (97%)	2 (2%)	2 (2%)	10 1
1	B	132/149 (89%)	130 (98%)	1 (1%)	1 (1%)	19 4
All	All	265/298 (89%)	259 (98%)	3 (1%)	3 (1%)	19 2

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1595	SER
1	A	1542[A]	SER
1	A	1542[B]	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	114/130 (88%)	109 (96%)	5 (4%)	28 4
1	B	104/130 (80%)	100 (96%)	4 (4%)	33 6
All	All	218/260 (84%)	209 (96%)	9 (4%)	36 5

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1519	ARG
1	A	1571	THR
1	A	1580[A]	ASN
1	A	1580[B]	ASN
1	A	1598	ILE
1	B	1502	ASP
1	B	1519	ARG
1	B	1548[A]	SER
1	B	1548[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1511	ASN
1	B	1568	ASN
1	B	1610	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	126/149 (84%)	0.12	6 (4%)	30	35	9, 22, 37, 47	1 (0%)
1	B	125/149 (83%)	0.73	12 (9%)	8	9	17, 29, 47, 78	0
All	All	251/298 (84%)	0.42	18 (7%)	15	18	9, 25, 43, 78	1 (0%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1596	GLY	6.5
1	A	1495	SER	6.4
1	B	1595	SER	5.7
1	B	1621	GLY	4.2
1	B	1590	SER	4.0
1	B	1594	LYS	3.8
1	A	1533	GLU	3.2
1	B	1619	TYR	3.1
1	B	1496	GLU	3.1
1	A	1570	GLY	3.1
1	B	1549	ALA	2.9
1	B	1547[A]	ILE	2.8
1	A	1534	GLU	2.7
1	A	1621	GLY	2.7
1	A	1528	VAL	2.6
1	B	1518	LEU	2.4
1	B	1598	ILE	2.3
1	B	1592	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	CA	B	1690	1/1	0.99	0.10	21,21,21,21	0
2	CA	A	1690	1/1	1.00	0.08	12,12,12,12	0

6.5 Other polymers (i)

There are no such residues in this entry.