



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 9, 2020 – 06:18 PM BST

PDB ID : 2A74
Title : Human Complement Component C3c
Authors : Janssen, B.J.C.; Huizinga, E.G.; Raaijmakers, H.C.A.; Roos, A.; Daha, M.R.;
Nilsson-Ekdahl, K.; Nilsson, B.; Gros, P.
Deposited on : 2005-07-04
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	643	8% (Poor fit) 65% (0 outliers), 26% (1 outlier), 6% (2 outliers), 1% (3+ outliers)
1	D	643	9% (Poor fit) 66% (0 outliers), 24% (1 outlier), 7% (2 outliers), 3% (3+ outliers)
2	B	188	7% (Poor fit) 78% (0 outliers), 14% (1 outlier), 5% (2 outliers), 5% (3+ outliers)
2	E	188	23% (Poor fit) 69% (0 outliers), 21% (1 outlier), 6% (2 outliers), 1% (3+ outliers)
3	C	343	9% (Poor fit) 51% (0 outliers), 26% (1 outlier), 7% (2 outliers), 14% (3+ outliers)
3	F	343	9% (Poor fit) 61% (0 outliers), 19% (1 outlier), 5% (2 outliers), 13% (3+ outliers)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	G	2	 100%
4	H	2	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NDG	G	2	X	-	-	X
4	NDG	H	2	X	-	-	X
5	GOL	A	712	-	-	X	X
5	GOL	B	702	-	-	-	X
5	GOL	D	711	-	-	X	X
5	GOL	E	705	-	-	-	X
5	GOL	E	706	-	-	-	X
5	GOL	F	707	-	-	-	X
6	NO3	F	802	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 18389 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Complement Component C3c.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	630	4907	3127	828	937	15	0	0	0
1	D	633	4933	3144	833	941	15	0	0	0

- Molecule 2 is a protein called Complement Component C3c.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	183	1480	950	249	276	5	0	0	0
2	E	184	1484	954	250	275	5	0	0	0

- Molecule 3 is a protein called Complement Component C3c.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	296	2407	1517	395	475	20	0	0	0
3	F	298	2421	1524	397	480	20	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	2	28	16	2	10	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
			Total	C	N				O
4	H	2	28	16	2	10	0	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	6	3	3	0	0
5	A	1	6	3	3	0	0
5	A	1	6	3	3	0	0
5	B	1	6	3	3	0	0
5	B	1	6	3	3	0	0
5	C	1	6	3	3	0	0
5	C	1	6	3	3	0	0
5	D	1	6	3	3	0	0
5	E	1	6	3	3	0	0
5	E	1	6	3	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	E	1	Total	C	O	0	0
			6	3	3		
5	F	1	Total	C	O	0	0
			6	3	3		
5	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is NITRATE ION (three-letter code: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	N	O	0	0
			4	1	3		
6	F	1	Total	N	O	0	0
			4	1	3		
6	F	1	Total	N	O	0	0
			4	1	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	188	Total	O	0	0
			188	188		
7	B	58	Total	O	0	0
			58	58		
7	C	71	Total	O	0	0
			71	71		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	148	Total 148	O 148	0	0
7	E	53	Total 53	O 53	0	0
7	F	93	Total 93	O 93	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Complement Component C3c




- Molecule 1: Complement Component C3c





- Molecule 3: Complement Component C3c



- Molecule 4: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	126.87Å 246.86Å 87.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.40 28.35 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.4 (40.00-2.40) 95.0 (28.35-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 2.31Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.215 , 0.275 0.213 , 0.272	Depositor DCC
R_{free} test set	3471 reflections (2.99%)	wwPDB-VP
Wilson B-factor (Å ²)	47.0	Xtrriage
Anisotropy	0.511	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	18389	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.48 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7203e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NDG, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	2/5004 (0.0%)	0.72	11/6799 (0.2%)
1	D	0.65	10/5032 (0.2%)	0.71	10/6839 (0.1%)
2	B	0.49	1/1512 (0.1%)	0.70	1/2055 (0.0%)
2	E	1.21	16/1516 (1.1%)	0.76	6/2060 (0.3%)
3	C	0.40	1/2453 (0.0%)	0.76	16/3305 (0.5%)
3	F	0.59	7/2468 (0.3%)	0.78	11/3327 (0.3%)
All	All	0.62	37/17985 (0.2%)	0.73	55/24385 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	D	0	8
3	C	0	1
3	F	0	3
All	All	0	17

The worst 5 of 37 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	564	GLU	CG-CD	19.98	1.81	1.51
2	E	837	GLU	CD-OE1	16.09	1.43	1.25
2	E	761	LYS	CE-NZ	13.09	1.81	1.49
2	E	758	GLU	CD-OE2	12.47	1.39	1.25
3	F	1397	LYS	CE-NZ	12.42	1.80	1.49

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	564	GLU	OE1-CD-OE2	7.91	132.80	123.30
2	E	761	LYS	CD-CE-NZ	-7.19	95.16	111.70
3	C	1413	ASP	CB-CG-OD2	6.27	123.94	118.30
3	F	1397	LYS	CD-CE-NZ	-6.14	97.58	111.70
1	A	30	ASP	CB-CG-OD2	5.94	123.65	118.30

There are no chirality outliers.

5 of 17 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	255	ASP	Peptide
1	A	403	ARG	Peptide
1	A	549	GLU	Peptide
1	A	63	ASN	Peptide
1	A	69	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4907	0	4970	204	0
1	D	4933	0	4996	195	0
2	B	1480	0	1501	30	0
2	E	1484	0	1511	46	0
3	C	2407	0	2316	97	0
3	F	2421	0	2326	62	0
4	G	28	0	23	2	0
4	H	28	0	19	1	0
5	A	18	0	24	6	0
5	B	12	0	16	2	0
5	C	12	0	16	0	0
5	D	6	0	8	5	0
5	E	18	0	24	2	0
5	F	12	0	16	4	0
6	C	4	0	0	0	0
6	F	8	0	0	0	0
7	A	188	0	0	11	0
7	B	58	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	71	0	0	6	0
7	D	148	0	0	3	0
7	E	53	0	0	2	0
7	F	93	0	0	2	0
All	All	18389	0	17766	601	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 601 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:564:GLU:CG	1:D:564:GLU:CD	1.81	1.49
3:F:1397:LYS:CE	3:F:1397:LYS:NZ	1.80	1.44
1:D:544:LYS:NZ	1:D:544:LYS:CE	1.79	1.43
2:E:761:LYS:CE	2:E:761:LYS:NZ	1.81	1.42
1:A:6:ILE:HD11	1:A:20:MET:CG	1.53	1.39

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	624/643 (97%)	582 (93%)	27 (4%)	15 (2%)	6 6
1	D	629/643 (98%)	593 (94%)	21 (3%)	15 (2%)	6 6
2	B	181/188 (96%)	172 (95%)	8 (4%)	1 (1%)	25 36
2	E	182/188 (97%)	167 (92%)	13 (7%)	2 (1%)	14 20
3	C	290/343 (84%)	270 (93%)	9 (3%)	11 (4%)	3 2
3	F	294/343 (86%)	274 (93%)	11 (4%)	9 (3%)	4 3

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	2200/2348 (94%)	2058 (94%)	89 (4%)	53 (2%)	6 6

5 of 53 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	410	SER
1	A	550	ASP
3	C	1361	THR
1	D	41	PRO
1	D	372	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	555/567 (98%)	498 (90%)	57 (10%)	7 10
1	D	558/567 (98%)	487 (87%)	71 (13%)	4 5
2	B	171/175 (98%)	156 (91%)	15 (9%)	10 15
2	E	171/175 (98%)	155 (91%)	16 (9%)	8 13
3	C	270/309 (87%)	232 (86%)	38 (14%)	3 4
3	F	272/309 (88%)	242 (89%)	30 (11%)	6 8
All	All	1997/2102 (95%)	1770 (89%)	227 (11%)	5 7

5 of 227 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	1555	GLN
1	D	98	VAL
3	F	1449	LEU
3	C	1573	LYS
1	D	34	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 49 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	1462	ASN
1	D	10	ASN
2	E	834	GLN
3	C	1579	HIS
1	D	38	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NDG	G	1	4	14,14,15	0.62	0	17,19,21	0.65	0
4	NDG	G	2	4	14,14,15	0.54	0	17,19,21	0.77	0
4	NDG	H	1	4	14,14,15	0.55	0	17,19,21	1.06	1 (5%)
4	NDG	H	2	4	14,14,15	0.87	1 (7%)	17,19,21	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NDG	G	1	4	-	0/6/23/26	0/1/1/1
4	NDG	G	2	4	1/1/5/7	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NDG	H	1	4	-	2/6/23/26	0/1/1/1
4	NDG	H	2	4	1/1/5/7	4/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	2	NDG	C1-C2	-2.64	1.48	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	1	NDG	O4-C4-C5	2.70	116.00	109.30

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	G	2	NDG	C1
4	H	2	NDG	C1

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	H	2	NDG	C8-C7-N2-C2
4	H	2	NDG	O7-C7-N2-C2
4	G	2	NDG	O5-C5-C6-O6
4	H	1	NDG	O5-C5-C6-O6
4	H	2	NDG	O5-C5-C6-O6


There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	1	NDG	2	0
4	G	2	NDG	1	0
4	H	1	NDG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	B	702	-	5,5,5	0.40	0	5,5,5	0.22	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	D	711	-	5,5,5	0.38	0	5,5,5	0.35	0
5	GOL	A	713	-	5,5,5	0.43	0	5,5,5	0.37	0
5	GOL	A	703	-	5,5,5	0.34	0	5,5,5	0.20	0
5	GOL	F	707	-	5,5,5	0.37	0	5,5,5	0.27	0
6	NO3	F	803	-	1,3,3	3.54	1 (100%)	0,3,3	0.00	-
5	GOL	C	709	-	5,5,5	0.31	0	5,5,5	0.40	0
5	GOL	F	710	-	5,5,5	0.35	0	5,5,5	0.22	0
5	GOL	E	706	-	5,5,5	0.36	0	5,5,5	0.47	0
5	GOL	E	704	-	5,5,5	0.42	0	5,5,5	0.46	0
6	NO3	C	801	-	1,3,3	3.61	1 (100%)	0,3,3	0.00	-
5	GOL	C	708	-	5,5,5	0.30	0	5,5,5	0.42	0
5	GOL	A	712	-	5,5,5	0.32	0	5,5,5	0.38	0
5	GOL	B	701	-	5,5,5	0.40	0	5,5,5	0.49	0
6	NO3	F	802	-	1,3,3	3.61	1 (100%)	0,3,3	0.00	-
5	GOL	E	705	-	5,5,5	0.37	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	702	-	-	3/4/4/4	-
5	GOL	D	711	-	-	0/4/4/4	-
5	GOL	A	713	-	-	1/4/4/4	-
5	GOL	A	703	-	-	2/4/4/4	-
5	GOL	F	707	-	-	4/4/4/4	-
5	GOL	C	709	-	-	0/4/4/4	-
5	GOL	F	710	-	-	3/4/4/4	-
5	GOL	E	706	-	-	3/4/4/4	-
5	GOL	E	704	-	-	2/4/4/4	-
5	GOL	C	708	-	-	2/4/4/4	-
5	GOL	A	712	-	-	0/4/4/4	-
5	GOL	B	701	-	-	4/4/4/4	-
5	GOL	E	705	-	-	2/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	801	NO3	O1-N	3.61	1.40	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	F	802	NO3	O1-N	3.61	1.40	1.24
6	F	803	NO3	O1-N	3.54	1.40	1.24

There are no bond angle outliers.

There are no chirality outliers.

5 of 26 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	702	GOL	O1-C1-C2-C3
5	A	703	GOL	C1-C2-C3-O3
5	F	707	GOL	O1-C1-C2-C3
5	F	710	GOL	O1-C1-C2-C3
5	E	705	GOL	O1-C1-C2-C3

There are no ring outliers.

7 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	711	GOL	5	0
5	A	703	GOL	2	0
5	F	707	GOL	3	0
5	F	710	GOL	1	0
5	A	712	GOL	4	0
5	B	701	GOL	2	0
5	E	705	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	630/643 (97%)	0.49	53 (8%) 11 10	30, 44, 55, 73	0
1	D	633/643 (98%)	0.56	58 (9%) 9 8	33, 44, 56, 78	0
2	B	183/188 (97%)	0.48	14 (7%) 13 12	32, 44, 52, 59	0
2	E	184/188 (97%)	1.14	44 (23%) 0 0	37, 44, 51, 71	0
3	C	296/343 (86%)	0.65	31 (10%) 6 5	33, 45, 56, 65	0
3	F	298/343 (86%)	0.73	32 (10%) 6 5	30, 45, 57, 68	0
All	All	2224/2348 (94%)	0.62	232 (10%) 6 6	30, 44, 55, 78	0

The worst 5 of 232 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	1502	ASP	11.1
3	C	1416	PHE	10.8
3	F	1501	SER	10.2
2	E	899	ILE	7.5
1	D	551	ARG	7.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NDG	G	2	14/15	0.64	0.47	64,66,69,69	0
4	NDG	H	2	14/15	0.68	0.56	75,76,78,79	0
4	NDG	H	1	14/15	0.73	0.40	61,66,70,73	0
4	NDG	G	1	14/15	0.86	0.29	54,57,59,62	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	713	6/6	0.45	0.29	61,65,66,68	0
5	GOL	E	706	6/6	0.48	0.57	57,60,63,64	0
5	GOL	F	707	6/6	0.61	0.54	86,88,88,89	0
5	GOL	A	712	6/6	0.64	0.68	78,81,82,85	0
6	NO3	F	802	4/4	0.64	0.58	89,89,89,90	0
5	GOL	E	705	6/6	0.65	0.48	62,66,66,67	0
5	GOL	B	702	6/6	0.68	0.56	87,88,88,89	0
6	NO3	F	803	4/4	0.69	0.37	73,73,73,73	0
5	GOL	D	711	6/6	0.73	0.77	81,83,84,88	0
5	GOL	A	703	6/6	0.73	0.32	60,62,63,65	0
5	GOL	F	710	6/6	0.77	0.20	72,73,73,74	0
6	NO3	C	801	4/4	0.79	0.32	77,77,77,77	0
5	GOL	C	708	6/6	0.79	0.22	66,72,73,75	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	GOL	C	709	6/6	0.80	0.28	90,90,91,91	0
5	GOL	E	704	6/6	0.86	0.24	45,49,51,52	0
5	GOL	B	701	6/6	0.93	0.12	45,46,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.