



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 22, 2023 – 08:06 AM EDT

PDB ID : 3AEQ  
Title : Structure of the light-independent protochlorophyllide reductase catalyzing a key reduction for greening in the dark  
Authors : Muraki, N.; Nomata, J.; Shiba, T.; Fujita, Y.; Kurisu, G.  
Deposited on : 2010-02-10  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	437	
1	C	437	
2	B	525	
2	D	525	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PMR	B	526	X	-	-	-
4	PMR	D	526	X	-	-	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 13035 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Light-independent protochlorophyllide reductase subunit N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	415	3164	2007	558	584	15	0	0	0
1	C	414	3169	2010	558	586	15	0	1	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MET	-	expression tag	UNP P26164
A	-11	ALA	-	expression tag	UNP P26164
A	-10	SER	-	expression tag	UNP P26164
A	-9	TRP	-	expression tag	UNP P26164
A	-8	SER	-	expression tag	UNP P26164
A	-7	HIS	-	expression tag	UNP P26164
A	-6	ALA	-	expression tag	UNP P26164
A	-5	PRO	-	expression tag	UNP P26164
A	-4	LYS	-	expression tag	UNP P26164
A	-3	PHE	-	expression tag	UNP P26164
A	-2	GLU	-	expression tag	UNP P26164
A	-1	LYS	-	expression tag	UNP P26164
A	0	GLY	-	expression tag	UNP P26164
A	1	ALA	-	expression tag	UNP P26164
C	-12	MET	-	expression tag	UNP P26164
C	-11	ALA	-	expression tag	UNP P26164
C	-10	SER	-	expression tag	UNP P26164
C	-9	TRP	-	expression tag	UNP P26164
C	-8	SER	-	expression tag	UNP P26164
C	-7	HIS	-	expression tag	UNP P26164
C	-6	ALA	-	expression tag	UNP P26164
C	-5	PRO	-	expression tag	UNP P26164
C	-4	LYS	-	expression tag	UNP P26164
C	-3	PHE	-	expression tag	UNP P26164
C	-2	GLU	-	expression tag	UNP P26164

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	LYS	-	expression tag	UNP P26164
C	0	GLY	-	expression tag	UNP P26164
C	1	ALA	-	expression tag	UNP P26164

- Molecule 2 is a protein called Light-independent protochlorophyllide reductase subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	419	Total 3297	C 2100	N 569	O 606	S 22	0	16	0
2	D	419	Total 3286	C 2092	N 568	O 604	S 22	0	16	0

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	S		
3	A	1	Total 8	Fe 4	S 4	0	0
3	C	1	Total 8	Fe 4	S 4	0	0

- Molecule 4 is Protochlorophyllide (three-letter code: PMR) (formula: C<sub>35</sub>H<sub>32</sub>MgN<sub>4</sub>O<sub>5</sub>).







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.63Å 81.28Å 177.10Å 90.00° 100.43° 90.00°	Depositor
Resolution (Å)	47.25 – 2.90 47.24 – 2.90	Depositor EDS
% Data completeness (in resolution range)	88.2 (47.25-2.90) 88.2 (47.24-2.90)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.01 (at 2.91Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.235 , 0.298 0.236 , 0.296	Depositor DCC
$R_{free}$ test set	2167 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	58.0	Xtrriage
Anisotropy	0.086	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 46.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.84	EDS
Total number of atoms	13035	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, PMR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/3227	0.62	1/4382 (0.0%)
1	C	0.49	0/3232	0.64	1/4389 (0.0%)
2	B	0.50	0/3369	0.63	1/4581 (0.0%)
2	D	0.50	0/3357	0.64	0/4565
All	All	0.49	0/13185	0.63	3/17917 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	418	LEU	CA-CB-CG	6.58	130.43	115.30
1	C	96	LEU	CA-CB-CG	6.22	129.61	115.30
2	B	264	LEU	CA-CB-CG	5.08	126.98	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	410[A]	LEU	Peptide
2	D	412[A]	GLU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3164	0	3199	56	0
1	C	3169	0	3201	51	0
2	B	3297	0	3324	113	0
2	D	3286	0	3309	90	0
3	A	8	0	0	0	0
3	C	8	0	0	1	0
4	B	45	0	31	9	0
4	D	45	0	31	18	0
5	A	3	0	0	0	0
5	B	6	0	0	0	0
5	C	1	0	0	0	0
5	D	3	0	0	0	0
All	All	13035	0	13095	302	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 302 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:408[A]:MET:HB3	4:B:526:PMR:O2A	1.44	1.17
2:D:410[A]:LEU:O	2:D:413[A]:HIS:HB2	1.47	1.12
2:D:408[A]:MET:HG2	4:D:526:PMR:O1A	1.47	1.11
2:D:408[A]:MET:CG	4:D:526:PMR:CGA	2.29	1.09
2:B:414[B]:LEU:HA	2:B:417[B]:MET:CB	1.86	1.04

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	413/437 (94%)	384 (93%)	27 (6%)	2 (0%)	29	61
1	C	413/437 (94%)	386 (94%)	26 (6%)	1 (0%)	47	78
2	B	428/525 (82%)	402 (94%)	21 (5%)	5 (1%)	13	40
2	D	427/525 (81%)	398 (93%)	23 (5%)	6 (1%)	11	36
All	All	1681/1924 (87%)	1570 (93%)	97 (6%)	14 (1%)	22	51

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	413[A]	HIS
2	B	413[B]	HIS
1	C	343	SER
2	D	408[A]	MET
2	D	408[B]	MET

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/344 (96%)	294 (89%)	35 (11%)	6	20
1	C	330/344 (96%)	291 (88%)	39 (12%)	5	16
2	B	347/417 (83%)	313 (90%)	34 (10%)	8	24
2	D	345/417 (83%)	313 (91%)	32 (9%)	9	27
All	All	1351/1522 (89%)	1211 (90%)	140 (10%)	7	21

5 of 140 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	105	ASN
2	D	142[A]	ARG
2	D	279	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	168	HIS
2	B	164	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	360	HIS
2	D	380	GLN
2	D	31	HIS
2	D	229	GLN
2	D	13	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	PMR	D	526	5	48,53,53	2.19	13 (27%)	54,89,89	1.40	5 (9%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SF4	C	425	1,2	0,12,12	-	-	-	-	-
4	PMR	B	526	5	48,53,53	2.26	13 (27%)	54,89,89	1.50	6 (11%)
3	SF4	A	425	1,2	0,12,12	-	-	-	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PMR	D	526	5	3/3/16/20	4/15/111/111	-
3	SF4	C	425	1,2	-	-	0/6/5/5
4	PMR	B	526	5	2/2/16/20	7/15/111/111	-
3	SF4	A	425	1,2	-	-	0/6/5/5

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	526	PMR	C4D-CHA	-6.03	1.37	1.45
4	D	526	PMR	C4D-CHA	-5.83	1.37	1.45
4	B	526	PMR	MG-NB	-5.70	1.94	2.05
4	D	526	PMR	MG-NB	-5.56	1.94	2.05
4	B	526	PMR	MG-NC	-4.51	1.95	2.06

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	526	PMR	C1A-NA-C4A	-6.11	103.96	106.71
4	D	526	PMR	O2D-CGD-CBD	4.28	118.87	111.27
4	D	526	PMR	C1A-NA-C4A	-4.13	104.85	106.71
4	B	526	PMR	O2D-CGD-CBD	4.11	118.57	111.27
4	B	526	PMR	C3C-C4C-NC	3.44	113.12	109.88

All (5) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	526	PMR	NC
4	B	526	PMR	NB
4	D	526	PMR	NA
4	D	526	PMR	NC
4	D	526	PMR	NB

5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	526	PMR	C4C-C3C-CAC-CBC
4	B	526	PMR	C2C-C3C-CAC-CBC
4	D	526	PMR	C2C-C3C-CAC-CBC
4	D	526	PMR	C4C-C3C-CAC-CBC
4	B	526	PMR	C2B-C3B-CAB-CBB


There are no ring outliers.

3 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	526	PMR	18	0
3	C	425	SF4	1	0
4	B	526	PMR	9	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	415/437 (94%)	0.04	0 <b>100</b> <b>100</b>	41, 52, 64, 69	0
1	C	414/437 (94%)	0.02	2 (0%) <b>91</b> <b>91</b>	40, 52, 64, 69	0
2	B	419/525 (79%)	0.03	12 (2%) 51 47	39, 47, 54, 64	5 (1%)
2	D	419/525 (79%)	0.03	11 (2%) 56 52	39, 47, 55, 65	6 (1%)
All	All	1667/1924 (86%)	0.03	25 (1%) <b>73</b> <b>73</b>	39, 49, 60, 69	11 (0%)

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	413[A]	HIS	5.1
2	B	413[A]	HIS	3.6
2	B	416[B]	THR	3.4
2	D	409[A]	GLY	3.1
2	B	408[A]	MET	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	PMR	D	526	45/45	0.85	0.44	23,26,27,28	45
4	PMR	B	526	45/45	0.93	0.37	15,18,23,26	45
3	SF4	A	425	8/8	0.98	0.13	31,33,33,34	0
3	SF4	C	425	8/8	0.99	0.12	33,34,35,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.