Full wwPDB EM Validation Report ⓘ

Mar 8, 2026 – 12:01 AM UTC

PDB ID	:	9AU0 / pdb_00009au0
EMDB ID	:	EMD-43839
Title	:	Cryo-EM structure of the BW245C-bound prostaglandin D2 receptor (DP1)-Gs complex
Authors	:	Davoudinasab, B.; Cherezov, V.; Han, G.W.; Kim, D.
Deposited on	:	2024-02-27
Resolution	:	2.45 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev132
Mogul	:	2022.3.0, CSD as543be (2022)
MolProbity	:	4-5-2 with Phenix2.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	5931 (1.96 - 2.95)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	394	55% 5% 40%
2	B	345	90% 8%
3	C	71	83% 17%
4	N	139	80% 12% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	R	385	

2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 8576 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Guanine nucleotide-binding protein G(s) subunit alpha isoforms short.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	236	Total	C	N	O	S	0	0
			1945	1225	351	362	7		

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	54	ASN	SER	engineered mutation	UNP P63092
A	226	ALA	GLY	engineered mutation	UNP P63092
A	268	ALA	GLU	engineered mutation	UNP P63092
A	271	LYS	ASN	engineered mutation	UNP P63092
A	274	ASP	LYS	engineered mutation	UNP P63092
A	280	LYS	ARG	engineered mutation	UNP P63092
A	284	ASP	THR	engineered mutation	UNP P63092
A	285	THR	ILE	engineered mutation	UNP P63092

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	339	Total	C	N	O	S	0	0
			2597	1602	465	509	21		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLY	-	expression tag	UNP P62873
B	-3	PRO	-	expression tag	UNP P62873
B	-2	GLY	-	expression tag	UNP P62873
B	-1	SER	-	expression tag	UNP P62873
B	0	SER	-	expression tag	UNP P62873
B	1	GLN	-	expression tag	UNP P62873

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	59	Total	C	N	O	S	0	0
			453	282	80	88	3		

- Molecule 4 is a protein called Nanobody-35.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	N	128	Total	C	N	O	S	0	0
			973	605	170	192	6		

- Molecule 5 is a protein called Prostaglandin D2 receptor.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	R	307	Total	C	N	O	S	0	0
			2394	1574	396	403	21		

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	-25	MET	-	expression tag	UNP Q13258
R	-24	LYS	-	expression tag	UNP Q13258
R	-23	THR	-	expression tag	UNP Q13258
R	-22	ILE	-	expression tag	UNP Q13258
R	-21	ILE	-	expression tag	UNP Q13258
R	-20	ALA	-	expression tag	UNP Q13258
R	-19	LEU	-	expression tag	UNP Q13258
R	-18	SER	-	expression tag	UNP Q13258
R	-17	TYR	-	expression tag	UNP Q13258
R	-16	ILE	-	expression tag	UNP Q13258
R	-15	PHE	-	expression tag	UNP Q13258
R	-14	CYS	-	expression tag	UNP Q13258
R	-13	LEU	-	expression tag	UNP Q13258
R	-12	VAL	-	expression tag	UNP Q13258
R	-11	PHE	-	expression tag	UNP Q13258
R	-10	ALA	-	expression tag	UNP Q13258
R	-9	ASP	-	expression tag	UNP Q13258
R	-8	TYR	-	expression tag	UNP Q13258
R	-7	LYS	-	expression tag	UNP Q13258
R	-6	ASP	-	expression tag	UNP Q13258
R	-5	ASP	-	expression tag	UNP Q13258
R	-4	ASP	-	expression tag	UNP Q13258

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
R	-3	ASP	-	expression tag	UNP Q13258
R	-2	GLY	-	expression tag	UNP Q13258
R	-1	ALA	-	expression tag	UNP Q13258
R	0	PRO	-	expression tag	UNP Q13258

- Molecule 6 is 7-{(4S)-3-[(3R)-3-cyclohexyl-3-hydroxypropyl]-2,5-dioxoimidazolidin-4-yl}heptanoic acid (CCD ID: A1AF8) (formula: $C_{19}H_{32}N_2O_5$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
6	R	1	26	19	2	5	0

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				AltConf
7	R	1	Total	C	N	O	0
			14	8	1	5	

- Molecule 8 is CHOLESTEROL HEMISUCCINATE (CCD ID: Y01) (formula: $C_{31}H_{50}O_4$).



Mol	Chain	Residues	Atoms			AltConf
8	R	1	Total	C	O	0
			35	31	4	

- Molecule 9 is water.

Mol	Chain	Residues	Atoms	AltConf
9	A	32	Total O 32 32	0
9	B	59	Total O 59 59	0
9	C	2	Total O 2 2	0
9	N	22	Total O 22 22	0
9	R	24	Total O 24 24	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Guanine nucleotide-binding protein G(s) subunit alpha isoforms short

Chain A:



• Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1

Chain B:



• Molecule 3: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2

Chain C:



• Molecule 4: Nanobody-35

Chain N:



• Molecule 5: Prostaglandin D2 receptor

Chain R:



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	1072665	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	65	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	61.441	Depositor
Minimum map value	-36.907	Depositor
Average map value	-0.008	Depositor
Map value standard deviation	1.030	Depositor
Recommended contour level	4.3	Depositor
Map size (Å)	302.4, 302.4, 302.4	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.84, 0.84, 0.84	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: Y01, NAG, A1AF8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.11	0/1981	0.27	0/2666
2	B	0.13	0/2644	0.32	0/3585
3	C	0.09	0/459	0.20	0/620
4	N	0.13	0/993	0.31	0/1345
5	R	0.16	0/2452	0.28	0/3324
All	All	0.13	0/8529	0.29	0/11540

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1945	0	1911	13	0
2	B	2597	0	2495	18	0
3	C	453	0	460	0	0
4	N	973	0	938	9	0
5	R	2394	0	2442	15	0
6	R	26	0	0	0	0
7	R	14	0	13	0	0
8	R	35	0	49	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	A	32	0	0	0	0
9	B	59	0	0	0	0
9	C	2	0	0	0	0
9	N	22	0	0	0	0
9	R	24	0	0	0	0
All	All	8576	0	8308	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:51:SER:HA	1:A:249:ALA:HB2	1.75	0.68
1:A:291:LEU:O	1:A:294:GLN:NE2	2.33	0.62
2:B:294:CYS:HB3	2:B:308:LEU:HB2	1.82	0.60
1:A:228:ARG:NH1	2:B:162:GLY:O	2.34	0.60
5:R:9:GLN:O	5:R:94:ARG:NH1	2.33	0.59
1:A:59:GLN:OE1	1:A:371:ASN:ND2	2.36	0.59
1:A:371:ASN:OD1	1:A:374:ARG:NH2	2.37	0.57
4:N:50:ASP:OD1	4:N:59:SER:OG	2.22	0.57
5:R:266:LEU:HD13	5:R:327:ARG:HE	1.70	0.57
1:A:385:ARG:NH1	5:R:260:GLU:OE2	2.38	0.56
1:A:321:PRO:HG2	1:A:324:ALA:HB2	1.87	0.56
1:A:228:ARG:NH1	2:B:186:ASP:OD1	2.38	0.55
4:N:52:SER:OG	4:N:55:GLY:O	2.25	0.54
1:A:254:ASN:ND2	1:A:306:SER:OG	2.41	0.53
2:B:62:HIS:HD2	2:B:105:TYR:H	1.56	0.53
1:A:295:ASP:N	1:A:295:ASP:OD1	2.42	0.53
5:R:284:ARG:NH2	5:R:301:GLU:OE1	2.34	0.53
4:N:32:TYR:O	4:N:72:ARG:NH2	2.38	0.52
2:B:166:CYS:HB2	2:B:180:PHE:HB2	1.92	0.52
2:B:42:ARG:NH1	2:B:43:ILE:O	2.44	0.51
5:R:230:ARG:NH1	5:R:233:GLN:OE1	2.43	0.51
2:B:271:CYS:HB2	2:B:290:ASP:HB2	1.92	0.51
5:R:3:SER:OG	5:R:298:ARG:NH1	2.45	0.50
4:N:51:ILE:HD13	4:N:72:ARG:HB2	1.94	0.49
4:N:22:CYS:HB3	4:N:79:LEU:HB3	1.93	0.49
4:N:37:VAL:HG12	4:N:47:TRP:HA	1.95	0.49
5:R:281:VAL:HG21	5:R:312:LEU:HD22	1.94	0.49
5:R:176:GLN:NE2	5:R:180:GLY:O	2.45	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:25:CYS:SG	2:B:259:GLN:NE2	2.87	0.47
2:B:165:THR:HG22	2:B:181:THR:HG22	1.95	0.47
2:B:51:LEU:HB2	2:B:336:LEU:HB2	1.97	0.46
5:R:89:GLN:OE1	5:R:91:ARG:NH2	2.36	0.46
2:B:320:VAL:HG22	2:B:327:VAL:HG12	1.98	0.46
2:B:192:LEU:HD23	2:B:199:PHE:HB3	1.98	0.45
5:R:43:ARG:HD3	5:R:48:TRP:HB2	1.99	0.45
2:B:303:ASP:OD1	2:B:303:ASP:N	2.47	0.44
5:R:60:VAL:HG13	5:R:149:GLY:HA3	1.99	0.44
4:N:33:LYS:HE3	4:N:107:CYS:SG	2.58	0.44
2:B:62:HIS:CD2	2:B:105:TYR:H	2.35	0.44
5:R:173:LYS:HD3	5:R:196:VAL:HG22	1.99	0.43
2:B:266:HIS:ND1	2:B:268:ASN:OD1	2.51	0.43
5:R:82:VAL:HG12	5:R:93:LEU:HD11	1.99	0.43
4:N:71:SER:HB3	4:N:80:TYR:HB2	2.00	0.43
5:R:280:PRO:O	5:R:284:ARG:HG2	2.19	0.43
1:A:10:GLU:OE1	1:A:13:ARG:NH2	2.52	0.43
2:B:57:LYS:HE2	2:B:59:TYR:HE1	1.83	0.42
2:B:212:ASP:HB2	2:B:219:ARG:HE	1.85	0.42
2:B:180:PHE:CE1	2:B:216:GLY:HA2	2.55	0.42
1:A:229:ASP:OD1	1:A:229:ASP:N	2.53	0.41
4:N:39:GLN:HB2	4:N:45:LEU:HD23	2.02	0.41
1:A:207:ILE:HD11	1:A:222:PHE:HB3	2.02	0.40
5:R:287:TYR:CZ	5:R:291:LYS:HD3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/394 (58%)	227 (99%)	3 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	337/345 (98%)	326 (97%)	11 (3%)	0	100	100
3	C	57/71 (80%)	55 (96%)	2 (4%)	0	100	100
4	N	126/139 (91%)	124 (98%)	2 (2%)	0	100	100
5	R	301/385 (78%)	297 (99%)	4 (1%)	0	100	100
All	All	1051/1334 (79%)	1029 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/349 (60%)	209 (100%)	0	100	100
2	B	280/286 (98%)	279 (100%)	1 (0%)	84	89
3	C	48/58 (83%)	48 (100%)	0	100	100
4	N	106/116 (91%)	106 (100%)	0	100	100
5	R	258/331 (78%)	257 (100%)	1 (0%)	84	89
All	All	901/1140 (79%)	899 (100%)	2 (0%)	85	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	59	TYR
5	R	65	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	220	HIS
1	A	267	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	278	ASN
1	A	279	ASN
2	B	44	GLN
2	B	62	HIS
2	B	156	GLN
2	B	259	GLN
4	N	82	GLN
4	N	120	GLN
4	N	123	GLN
5	R	176	GLN
5	R	224	ASN
5	R	229	HIS
5	R	263	HIS

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
6	A1AF8	R	401	-	27,27,27	0.58	0	30,35,35	0.84	1 (3%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
7	NAG	R	402	5	14,14,15	0.72	0	17,19,21	0.93	1 (5%)
8	Y01	R	403	-	38,38,38	0.71	1 (2%)	57,57,57	1.45	10 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	A1AF8	R	401	-	-	4/18/42/42	0/2/2/2
7	NAG	R	402	5	-	2/6/23/26	0/1/1/1
8	Y01	R	403	-	-	3/19/77/77	0/4/4/4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	R	403	Y01	CBH-CBF	-2.36	1.52	1.56

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
8	R	403	Y01	OAW-CAY-CAM	4.05	120.25	111.48
8	R	403	Y01	CAK-CBD-CBG	-2.64	107.19	110.93
8	R	403	Y01	CBF-CBD-CBG	-2.48	105.85	109.09
8	R	403	Y01	CAV-CAZ-CAI	-2.33	117.42	120.57
8	R	403	Y01	CAP-CAQ-CBG	-2.27	100.69	105.14
7	R	402	NAG	C2-N2-C7	2.23	125.89	122.90
8	R	403	Y01	CAP-CBE-CBB	-2.19	108.86	112.18
8	R	403	Y01	CAV-CAZ-CBH	2.17	119.20	116.42
8	R	403	Y01	CBI-CBG-CBD	-2.14	111.37	114.41
8	R	403	Y01	CAD-CBH-CBF	-2.09	109.31	111.66
8	R	403	Y01	CAQ-CBG-CBI	-2.05	101.43	103.84
6	R	401	A1AF8	C15-N14-C04	2.02	125.54	122.19

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	R	402	NAG	C8-C7-N2-C2
7	R	402	NAG	O7-C7-N2-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	R	401	A1AF8	C05-C06-C07-C08
8	R	403	Y01	CAO-CAJ-CAN-CBA
6	R	401	A1AF8	C07-C08-C09-C10
6	R	401	A1AF8	C08-C09-C10-C11
8	R	403	Y01	CAL-CAM-CAY-OAW
8	R	403	Y01	CAX-CAL-CAM-CAY
6	R	401	A1AF8	C06-C07-C08-C09

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand A1AF8 R 401



Bond lengths



Bond angles



Torsions



Rings

Ligand Y01 R 403



Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation

This section contains visualisations of the EMDB entry EMD-43839. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 180



Y Index: 180



Z Index: 180

6.2.2 Raw map



X Index: 180



Y Index: 180



Z Index: 180

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 182



Y Index: 173



Z Index: 228

6.3.2 Raw map



X Index: 182



Y Index: 173



Z Index: 136

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



X Y Z

6.4.2 Raw map



X Y Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

This section was not generated.

6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 66 nm^3; this corresponds to an approximate mass of 59 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.408 Å^{-1}

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.408 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.45	-	-
Author-provided FSC curve	2.45	2.83	2.48
Unmasked-calculated*	3.07	3.51	3.13

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.07 differs from the reported value 2.45 by more than 10 %

9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-43839 and PDB model 9AU0. Per-residue inclusion information can be found in section 3 on page 9.

9.1 Map-model overlay

This section was not generated.

9.2 Q-score mapped to coordinate model



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model

This section was not generated.

9.4 Atom inclusion (i)



At the recommended contour level, 93% of all backbone atoms, 93% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (4.3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9260	0.6330
A	0.8470	0.6010
B	0.9590	0.6650
C	0.8270	0.5940
N	0.9610	0.6550
R	0.9660	0.6210