wwPDB X-ray Structure Validation Summary Report (i)

Sep 25, 2023 – 11:19 PM EDT

PDB ID	:	6BAP
Title	:	Stigmatella aurantiaca bacterial phytochrome PAS-GAF-PHY, T289H mutant
Authors	:	Schmidt, M.; Stojkovic, E.
Deposited on	:	2017-10-14
Resolution	:	2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	515	3% 63% 30% • • •
1	B	515	3% 57% 36% 5% • •
1	C	515	32% 38% 46% 11% • •

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 12158 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Photoreceptor-histidine kinase BphP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	505	3982	2513	726	730	13	0	0	0
1	B	505	3982	2513	726	730	13	0	0	0
1	C	500	3941	2488	720	720	13	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	289	HIS	THR	engineered mutation	UNP Q097N3
B	289	HIS	THR	engineered mutation	UNP Q097N3
C	289	HIS	THR	engineered mutation	UNP Q097N3

- Molecule 2 is 3-[5-[(Z)-(4-ethenyl-3-methyl-5-oxidanylidene-pyrrol-2-ylidene)methyl]-2-[[5-[(Z)-(3-ethenyl-4-methyl-5-oxidanylidene-pyrrol-2-ylidene)methyl]-3-(3-hydroxy-3-oxopropyl)-4-methyl-1H-pyrrol-2-yl]methyl]-4-methyl-1H-pyrrol-3-yl]propanoic acid (three-letter code: BLR) (formula: $C_{33}H_{36}N_4O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	43	33	4	6	0	0
2	B	1	43	33	4	6	0	0
2	C	1	43	33	4	6	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	46	46	46	0	0
3	B	42	42	42	0	0
3	C	36	36	36	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Photoreceptor-histidine kinase BphP

Chain A: 3% 63% 30%



• Molecule 1: Photoreceptor-histidine kinase BphP

Chain B: 3% 57% 36% 5%





- Molecule 1: Photoreceptor-histidine kinase BphP

Chain C:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α, β, γ	83.31Å 83.31Å 475.05Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.28 – 2.65 40.28 – 2.59	Depositor EDS
% Data completeness (in resolution range)	98.8 (40.28-2.65) 98.6 (40.28-2.59)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$< I/\sigma(I) >$ [1]	1.49 (at 2.58Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.241 , 0.315 0.241 , 0.313	Depositor DCC
R_{free} test set	3051 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å^2)	70.1	Xtriage
Anisotropy	0.079	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.31 , 63.3	EDS
L-test for twinning[2]	$< \|L\| > = 0.44$, $< L^2 > = 0.26$	Xtriage
Estimated twinning fraction	0.068 for -h,-k,l	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	12158	wwPDB-VP
Average B, all atoms (Å^2)	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.66% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BLR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.61	4/4075 (0.1%)	0.77	6/5529 (0.1%)
1	B	0.59	3/4075 (0.1%)	0.75	4/5529 (0.1%)
1	C	0.93	16/4034 (0.4%)	0.91	18/5473 (0.3%)
All	All	0.73	23/12184 (0.2%)	0.81	28/16531 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	4
1	C	0	14
All	All	0	19

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	298	PRO	N-CD	-34.79	0.99	1.47
1	C	74	PRO	N-CD	-17.44	1.23	1.47
1	B	495	GLU	CD-OE1	-9.56	1.15	1.25
1	C	173	ARG	CZ-NH2	-8.98	1.21	1.33
1	C	100	ARG	CZ-NH1	-8.33	1.22	1.33

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	413	TYR	C-N-CD	-10.37	97.78	120.60
1	A	394	TRP	CA-CB-CG	10.16	133.01	113.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	145	LEU	CB-CG-CD1	-9.99	94.02	111.00
1	C	223	ARG	CG-CD-NE	-9.22	92.44	111.80
1	C	173	ARG	NE-CZ-NH2	-7.84	116.38	120.30

There are no chirality outliers.

5 of 19 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	400	GLY	Peptide
1	B	325	TYR	Peptide
1	B	326	ASP	Peptide
1	B	393	GLU	Peptide
1	B	422	VAL	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3982	0	3925	175	0
1	B	3982	0	3925	194	0
1	C	3941	0	3886	413	0
2	A	43	0	32	5	0
2	B	43	0	32	4	0
2	C	43	0	33	15	0
3	A	46	0	0	7	0
3	B	42	0	0	5	0
3	C	36	0	0	19	0
All	All	12158	0	11833	758	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 758 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:ARG:NH1	1:B:307:PHE:CA	1.72	1.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:ARG:NH1	1:B:307:PHE:HA	1.19	1.44
1:C:429:PHE:O	1:C:501:ARG:NH1	1.65	1.29
1:A:142:ARG:HH11	1:B:307:PHE:CB	1.54	1.19
1:C:154:LEU:HD21	1:C:155:GLN:NE2	1.65	1.12

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	503/515 (98%)	446 (89%)	44 (9%)	13 (3%)	5 7
1	B	503/515 (98%)	444 (88%)	39 (8%)	20 (4%)	3 3
1	C	498/515 (97%)	332 (67%)	122 (24%)	44 (9%)	1 0
All	All	1504/1545 (97%)	1222 (81%)	205 (14%)	77 (5%)	2 2

5 of 77 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	THR
1	A	44	VAL
1	A	389	THR
1	A	458	LYS
1	B	14	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	419/427 (98%)	405 (97%)	14 (3%)	38	54
1	B	419/427 (98%)	409 (98%)	10 (2%)	49	67
1	C	414/427 (97%)	386 (93%)	28 (7%)	16	24
All	All	1252/1281 (98%)	1200 (96%)	52 (4%)	30	45

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	129	ARG
1	C	203	HIS
1	C	490	LYS
1	C	136	SER
1	C	154	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	155	GLN
1	C	364	HIS
1	C	510	GLN
1	C	417	GLN
1	C	58	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	BLR	A	601	1	40,46,46	5.14	17 (42%)	44,67,67	1.57	9 (20%)
2	BLR	B	601	1	40,46,46	5.12	17 (42%)	44,67,67	1.45	9 (20%)
2	BLR	C	601	1	40,46,46	5.19	17 (42%)	44,67,67	1.60	8 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BLR	A	601	1	-	9/24/58/58	0/4/4/4
2	BLR	B	601	1	-	11/24/58/58	0/4/4/4
2	BLR	C	601	1	-	14/24/58/58	0/4/4/4

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	BLR	CHA-C4D	-24.25	1.33	1.51
2	A	601	BLR	CHA-C4D	-24.11	1.33	1.51
2	C	601	BLR	CHA-C4D	-23.42	1.33	1.51
2	C	601	BLR	C3D-C4D	10.77	1.55	1.39
2	C	601	BLR	C1B-NB	9.76	1.54	1.37

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	C	601	BLR	CAD-C3D-C4D	4.18	130.23	127.30
2	C	601	BLR	C4C-NC-C1C	-4.07	105.48	110.67
2	A	601	BLR	CAA-C2A-C1A	4.00	130.10	127.30
2	A	601	BLR	CAD-C3D-C4D	-3.43	124.89	127.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	601	BLR	C4C-NC-C1C	-3.25	106.53	110.67

There are no chirality outliers.

5 of 34 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	BLR	C2A-C1A-CHA-C4D
2	A	601	BLR	NA-C4A-CHB-C1B
2	A	601	BLR	C3A-C4A-CHB-C1B
2	B	601	BLR	C2A-C1A-CHA-C4D
2	B	601	BLR	C3D-C4D-CHA-C1A

There are no ring outliers.

3 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	BLR	5	0
2	B	601	BLR	4	0
2	C	601	BLR	15	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand BLR A 601



Bond lengths



Bond angles



Torsions



Rings

Ligand BLR B 601



Bond lengths



Bond angles



Torsions



Rings

Ligand BLR C 601



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	505/515 (98%)	0.03	18 (3%)	42	39	35, 64, 99, 162	0
1	B	505/515 (98%)	0.05	18 (3%)	42	39	35, 69, 109, 135	1 (0%)
1	C	500/515 (97%)	2.02	167 (33%)	0	0	54, 107, 209, 230	3 (0%)
All	All	1510/1545 (97%)	0.70	203 (13%)	3	2	35, 74, 189, 230	4 (0%)

The worst 5 of 203 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	337	ALA	32.8
1	C	380	GLY	17.2
1	C	368	ALA	15.6
1	C	378	VAL	14.9
1	C	491	ALA	14.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	BLR	C	601	43/43	0.78	0.24	86,97,113,118	0
2	BLR	A	601	43/43	0.94	0.20	47,60,78,94	1
2	BLR	B	601	43/43	0.95	0.18	43,51,63,67	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around BLR C 601:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)







Electron density around BLR A 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BLR B 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.