



Full wwPDB X-ray Structure Validation Report i

Jun 22, 2024 – 06:52 PM EDT

PDB ID : 6BFI
Title : Vinculin homolog in a sponge (phylum Porifera) reveals vertebrate-like cell adhesions involved in early multicellular evolution
Authors : Weis, W.I.; Chodaparambil, J.V.
Deposited on : 2017-10-26
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 2 unique types of molecules in this entry. The entry contains 12363 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VIN1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	805	Total	C 6124	N 3804	O 1098	S 1186	36	0	2	0
1	B	806	Total	C 6106	N 3794	O 1091	S 1185	36	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	62	Total O 62 62	0	0
2	B	71	Total O 71 71	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: VIN1



- Molecule 1: VIN1



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	50.34Å 93.68Å 101.63Å 114.81° 93.29° 90.23°	Depositor
Resolution (Å)	36.50 – 2.30 92.06 – 2.30	Depositor EDS
% Data completeness (in resolution range)	92.6 (36.50-2.30) 92.6 (92.06-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.37 (at 2.29Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ????)	Depositor
R , R_{free}	0.218 , 0.266 0.223 , 0.270	Depositor DCC
R_{free} test set	7546 reflections (10.10%)	wwPDB-VP
Wilson B-factor (Å ²)	49.0	Xtriage
Anisotropy	0.609	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 44.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for -h,k,-k-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12363	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.25	0/6190	0.37	0/8348
1	B	0.25	0/6169	0.38	0/8322
All	All	0.25	0/12359	0.38	0/16670

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6124	0	6295	49	0
1	B	6106	0	6272	54	0
2	A	62	0	0	2	0
2	B	71	0	0	0	0
All	All	12363	0	12567	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:695:VAL:HG11	1:A:811:ILE:HD11	1.63	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:695:VAL:HG11	1:B:811:ILE:HD11	1.67	0.75
1:A:362:GLN:OE1	1:A:365:ARG:NH2	2.27	0.68
1:B:176:GLU:OE2	1:B:199:ARG:NH2	2.31	0.64
1:A:362:GLN:NE2	1:A:498:GLU:OE2	2.31	0.61
1:B:304:ARG:NH2	1:B:427:SER:OG	2.34	0.60
1:B:372:VAL:HG21	1:B:494:ILE:HG13	1.83	0.60
1:A:517:ILE:HD13	1:A:526:LYS:HB3	1.84	0.59
1:B:303:ARG:NH2	1:B:326:ASP:OD2	2.35	0.59
1:B:152:ASP:N	1:B:152:ASP:OD1	2.36	0.59
1:B:683:GLN:HB2	1:B:834:LYS:HE2	1.84	0.59
1:A:698:MET:HE2	1:A:732:ASN:HB3	1.85	0.58
1:B:574:PRO:HA	1:B:577:ILE:HG12	1.85	0.58
1:B:21:VAL:HG13	1:B:112:ALA:HB1	1.85	0.58
1:A:203:PRO:HA	1:A:206:ILE:HG12	1.85	0.57
1:A:751:ARG:NH2	2:A:902:HOH:O	2.36	0.57
1:B:750:LYS:HA	1:B:753:LYS:HG2	1.86	0.57
1:B:214:LEU:HB3	1:B:465:ASN:HD21	1.69	0.57
1:B:44:VAL:HG11	1:B:89:SER:HA	1.88	0.56
1:A:142:GLN:HE22	1:A:243:GLU:HG2	1.70	0.56
1:B:744:ALA:HB2	1:B:756:LEU:HD23	1.87	0.56
1:A:744:ALA:HB2	1:A:756:LEU:HD23	1.88	0.56
1:B:324:GLU:OE2	1:B:356:LYS:NZ	2.39	0.55
1:B:10:LEU:HD11	1:B:124:MET:HE3	1.87	0.55
1:B:580:LYS:HA	1:B:583:VAL:HG13	1.89	0.55
1:B:276:HIS:CG	1:B:621:ARG:HD3	2.42	0.55
1:A:29:GLU:OE2	1:A:717:ARG:N	2.39	0.54
1:B:513:GLU:OE2	1:B:516:ARG:NH1	2.41	0.54
1:A:68:GLU:HG3	1:B:570:LYS:HA	1.90	0.53
1:A:372:VAL:HG21	1:A:494:ILE:HG13	1.90	0.53
1:B:44:VAL:HG21	1:B:92:ALA:HB2	1.91	0.53
1:A:521:GLU:OE2	1:A:592:GLN:NE2	2.41	0.51
1:B:824:ASP:OD1	1:B:825:SER:N	2.43	0.51
1:A:184:ASN:ND2	1:A:255:PHE:H	2.09	0.51
1:B:279:GLN:HG3	1:B:357:LEU:HD23	1.92	0.51
1:A:818:CYS:SG	1:A:832:TRP:NE1	2.84	0.51
1:A:610:ARG:O	1:A:614:GLU:HB2	2.11	0.50
1:B:62:VAL:O	1:B:71:LYS:NZ	2.44	0.50
1:B:157:LEU:HD13	1:B:209:ILE:HG21	1.94	0.50
1:A:113:ARG:HD3	1:A:780:SER:HB3	1.94	0.49
1:A:142:GLN:NE2	1:A:243:GLU:HG2	2.28	0.49
1:A:342:ASP:O	1:A:347:ARG:NH1	2.47	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179:HIS:CG	1:B:180:PRO:HD3	2.49	0.48
1:A:27:PHE:CE1	1:A:38:GLU:HG3	2.49	0.48
1:A:218:LYS:HD3	1:A:218:LYS:HA	1.74	0.48
1:A:303:ARG:HE	1:A:322:ALA:HB1	1.79	0.48
1:A:157:LEU:O	1:A:161:LEU:HB2	2.13	0.47
1:B:513:GLU:HG2	1:B:530:LYS:HE2	1.95	0.47
1:A:7:THR:HG22	1:A:10:LEU:H	1.79	0.46
1:B:697:GLN:HG3	1:B:701:LYS:HE2	1.96	0.46
1:B:153:VAL:HG23	1:B:155:ALA:H	1.80	0.46
1:A:184:ASN:HD21	1:A:255:PHE:H	1.62	0.46
1:A:130:GLU:O	1:A:134:MET:HG3	2.15	0.46
1:B:126:ALA:O	1:B:130:GLU:HG2	2.16	0.46
1:B:553:GLU:HG2	1:B:750:LYS:HE3	1.99	0.45
1:A:375:ASP:OD2	1:A:420:THR:OG1	2.34	0.45
1:B:202:ILE:HB	1:B:203:PRO:HD3	1.97	0.45
1:B:27:PHE:CE1	1:B:38:GLU:HG3	2.52	0.45
1:A:303:ARG:NH2	1:A:326:ASP:OD2	2.50	0.45
1:B:680:GLU:HG2	1:B:808:LYS:HG3	1.99	0.44
1:A:256:GLU:OE1	1:A:256:GLU:N	2.50	0.44
1:A:574:PRO:HA	1:A:577:ILE:HG12	1.99	0.44
1:A:172:MET:O	1:A:176:GLU:HB2	2.17	0.43
1:A:719:LYS:NZ	1:A:781:MET:O	2.51	0.43
1:A:74:LEU:HB3	1:A:75:PRO:HD3	2.01	0.43
1:B:121:ASP:HA	1:B:124:MET:HB2	2.00	0.43
1:B:142:GLN:HB2	1:B:242:ILE:HG21	2.00	0.43
1:B:54:LEU:HD23	1:B:122:LEU:HD23	1.99	0.43
1:B:696:LYS:O	1:B:700:ARG:HG3	2.19	0.43
1:A:108:LEU:O	1:A:112:ALA:HB2	2.19	0.43
1:A:580:LYS:O	1:A:583:VAL:HG22	2.19	0.43
1:B:580:LYS:O	1:B:583:VAL:HG22	2.19	0.43
1:B:207:SER:O	1:B:211:VAL:HG13	2.19	0.43
1:B:530:LYS:HD3	1:B:530:LYS:HA	1.86	0.43
1:A:179:HIS:CG	1:A:180:PRO:HD3	2.54	0.42
1:B:148:ALA:HA	1:B:151:ILE:HD13	2.00	0.42
1:B:551:ASN:OD1	1:B:751:ARG:HG3	2.18	0.42
1:A:199[A]:ARG:NH1	2:A:908:HOH:O	2.52	0.42
1:A:190:MET:O	1:A:194:GLU:HG2	2.20	0.42
1:A:619:PRO:HA	1:A:620:PRO:HD3	1.95	0.42
1:B:67:ASP:HB3	1:B:70:PHE:HB3	2.02	0.42
1:B:88:LEU:HD23	1:B:88:LEU:HA	1.84	0.42
1:B:208:SER:O	1:B:211:VAL:HG22	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:705:MET:O	1:A:709:THR:HG23	2.20	0.42
1:A:122:LEU:HD23	1:A:122:LEU:HA	1.89	0.42
1:B:500:MET:SD	1:B:541:VAL:HG13	2.60	0.41
1:A:763:ILE:N	1:A:764:PRO:HD2	2.35	0.41
1:B:708:TYR:CE2	1:B:721:ASP:HB3	2.55	0.41
1:B:83:GLN:O	1:B:86:VAL:HG12	2.20	0.41
1:B:403:PHE:CG	1:B:460:LEU:HD22	2.55	0.41
1:A:682:ASP:HA	1:A:696:LYS:HE3	2.02	0.41
1:A:157:LEU:HD13	1:A:209:ILE:HG21	2.01	0.41
1:B:430:CYS:HB3	1:B:493:VAL:HG13	2.03	0.41
1:A:226:ASP:OD2	1:A:457:ARG:HD2	2.21	0.41
1:A:494:ILE:HD13	1:A:494:ILE:HA	1.87	0.41
1:A:549:ALA:HB2	1:A:561:LEU:HD23	2.03	0.41
1:A:668:ASN:HB3	1:A:671:HIS:HB3	2.03	0.41
1:B:763:ILE:N	1:B:764:PRO:HD2	2.35	0.41
1:A:186:ALA:O	1:A:190:MET:HG3	2.21	0.40
1:B:382:PRO:HG2	1:B:413:HIS:CG	2.56	0.40
1:B:511:VAL:HG13	1:B:599:ALA:HB1	2.02	0.40
1:B:103:GLU:O	1:B:107:LYS:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	797/846 (94%)	780 (98%)	15 (2%)	2 (0%)	41 50
1	B	796/846 (94%)	775 (97%)	20 (2%)	1 (0%)	51 64
All	All	1593/1692 (94%)	1555 (98%)	35 (2%)	3 (0%)	47 58

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	284	SER
1	A	620	PRO
1	B	102	PRO

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	654/686 (95%)	647 (99%)	7 (1%)	73 86
1	B	652/686 (95%)	646 (99%)	6 (1%)	78 89
All	All	1306/1372 (95%)	1293 (99%)	13 (1%)	76 87

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	PHE
1	A	146	ASP
1	A	290	SER
1	A	363	GLU
1	A	524	SER
1	A	583	VAL
1	A	621	ARG
1	B	5	PHE
1	B	19	GLN
1	B	35	LEU
1	B	83	GLN
1	B	556	GLU
1	B	583	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	805/846 (95%)	0.21	14 (1%) 70 76	45, 71, 118, 177	0
1	B	806/846 (95%)	0.23	19 (2%) 59 66	44, 72, 123, 165	0
All	All	1611/1692 (95%)	0.22	33 (2%) 65 71	44, 71, 119, 177	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	784	GLY	5.9
1	A	591	ILE	5.0
1	B	525	MET	4.5
1	B	522	GLU	4.0
1	B	35	LEU	3.9
1	B	784	GLY	3.5
1	B	62	VAL	3.4
1	A	822	ARG	3.3
1	B	646	LEU	3.1
1	A	145	LEU	2.9
1	B	836	GLY	2.9
1	A	824	ASP	2.8
1	B	61	MET	2.8
1	A	583	VAL	2.8
1	B	220	GLY	2.8
1	B	1	MET	2.8
1	A	258	GLU	2.6
1	A	157	LEU	2.5
1	A	821	LEU	2.4
1	B	572	ILE	2.4
1	A	520	ALA	2.4
1	B	826	PRO	2.3
1	B	787	ARG	2.3
1	A	525	MET	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	152	ASP	2.3
1	A	37	LYS	2.2
1	B	255	PHE	2.1
1	B	579	ALA	2.1
1	B	141	VAL	2.1
1	A	739	LEU	2.1
1	B	344	ARG	2.1
1	A	687	LYS	2.1
1	B	679	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.