wwPDB NMR Structure Validation Summary Report (i)

Feb 8, 2022 – 12:43 PM EST

PDB ID	:	1BVM
Title	:	SOLUTION NMR STRUCTURE OF BOVINE PANCREATIC PHOSPHOLIPASE A2, 20 STRUCTURES
Authors	:	Yuan, C.-H.; Byeon, I.-J.L.; Li, Y.; Tsai, M.-D.
Deposited on	:	1998-09-14

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.26
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	123	

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:2-A:30, A:35-A:63, A:69-A:123 (113)	0.39	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 5, 6, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 20
2	4, 7, 19
Single-model clusters	18

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1842 atoms, of which 885 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PROTEIN (PHOSPHOLIPASE A2).

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	123	1842	586	885	164	192	15	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	122	ASN	LYS	variant	UNP P00593

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: PROTEIN (PHOSPHOLIPASE A2)

Chain A: 41% 46% • • 8%

A1 L2 W3 Q4 C11 K12 I13 S16 E17 D21 F22 N23 N24 Y25 G26 C27 Y28 L31 G32 G33 S34 G35 T36 P37 V38 D39 D40 L41 D42 R43 Q46 D49 K53 Q54 A55 K56 K57 L58 D59 S60 C61 K62 V63 L64 V65 D66 N67 P68 Y69 T70 N71 N72 Y73

S74 Y75 S76 C77 S78 N79 N80 E81 I82 T83 C84 S85 N88 N89 E92 R100 I104 C105 F106 S107 K108 V109 K113 K116 N117 L118 D119 K120 K121 N122 C123

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 2. Colouring as in section 4.1 above.

- Molecule 1: PROTEIN (PHOSPHOLIPASE A2)

Chain A: 36% 50% 6% • 8%

A1 L2 W3 Q4 C11 K12 I13 S16 L19 L20 N23 N24 Y25 G26 C27 Y28 C29 G30 L31 G32 G33 S34 G35 T36 P37 V38 D39 D40 R43 Q46 T47 H48 D49 Y52 K53 Q54 A55 K56 K57 L58 D59 S60 C61 K62 V63 L64 V65 D66 N67 P68 Y69 T70 N71

N72 Y73 S74 Y75 S76 C77 S78 N79 N80 E81 I82 T83 C84 S85 N89 E92 A93 R100 I104 C105 F106 S107 K108 V109 N112 K113 E114 H115 K116 N117 L118 D119 K120 K121 N122 C123

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 40 calculated structures, 20 were deposited, based on the following criterion: *CLOSEST TO THE MEAN STRUCTURE WHICH SHOWS LITTLE RESTRAINT VIOLATION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	
X-PLOR	structure solution	

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.9±0.4
All	All	0	37

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	100	ARG	Sidechain	19
1	A	43	ARG	Sidechain	18

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	892	819	819	45±6
All	All	17840	16380	16380	895

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

5 of 332 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:38:VAL:CG1	1:A:118:LEU:HD22	0.95	1.92	18	5
1:A:38:VAL:HG11	1:A:118:LEU:HD22	0.89	1.43	10	10
1:A:105:CYS:O	1:A:109:VAL:HG22	0.89	1.67	8	16
1:A:17:GLU:CB	1:A:20:LEU:HD12	0.76	2.10	19	3
1:A:55:ALA:HB2	1:A:94:PHE:CD1	0.72	2.19	17	3

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	112/123 (91%)	98±3 (87±2%)	10±3 (9±3%)	4±2 (4±2%)	5	32
All	All	2240/2460 (91%)	1953 (87%)	200 (9%)	87 (4%)	5	32

5 of 18 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	78	SER	20
1	A	26	GLY	14
1	A	70	THR	10
1	A	73	TYR	7
1	A	29	CYS	5

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	104/111 (94%)	86±3 (83±3%)	18±3 (17±3%)	4	39
All	All	2080/2220 (94%)	1717 (83%)	363 (17%)	4	39

5 of 56 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	76	SER	19
1	A	43	ARG	18
1	A	58	LEU	18
1	A	2	LEU	15
1	A	16	SER	15

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided