



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 16, 2024 – 03:51 AM EDT

PDB ID : 2C4F  
Title : crystal structure of factor VII.stf complexed with pd0297121  
Authors : Kohrt, J.T.; Zhang, E.  
Deposited on : 2005-10-18  
Resolution : 1.72 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6152 (1.74-1.70)
Ramachandran outliers	138981	6051 (1.74-1.70)
Sidechain outliers	138945	6051 (1.74-1.70)
RSRZ outliers	127900	5629 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	254	<div> <div>10%</div> <div>76%</div> <div>22%</div> <div>.</div> </div>
2	L	142	<div> <div>17%</div> <div>70%</div> <div>25%</div> <div>..</div> </div>
3	T	75	<div> <div>%</div> <div>87%</div> <div>13%</div> </div>
4	U	116	<div> <div>17%</div> <div>75%</div> <div>25%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CGU	L	35	-	-	-	X

## 2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 5058 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called COAGULATION FACTOR VII PRECURSOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	254	Total	C	N	O	S	0	0	0
			1974	1253	351	357	13			

- Molecule 2 is a protein called COAGULATION FACTOR VII PRECURSOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	139	Total	C	N	O	S	0	0	0
			1117	673	185	244	15			

- Molecule 3 is a protein called TISSUE FACTOR PRECURSOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	T	75	Total	C	N	O	S	0	0	0
			615	397	95	121	2			


- Molecule 4 is a protein called TISSUE FACTOR PRECURSOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	U	116	Total	C	N	O	S	0	0	0
			940	593	155	189	3			

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	1	Total	Ca	0	0
			1	1		
5	L	4	Total	Ca	0	0
			4	4		


- Molecule 6 is 2-{{6-{3-[AMINO(IMINO)METHYL]PHENOXY}-4-(DIISOPROPYLAMINO)-3,5-DIFLUOROPYRIDIN-2-YL]OXY}-5-[(ISOBUTYLAMINO)CARBONYL]BEN ZOIC ACID (three-letter code: GIL) (formula: C<sub>30</sub>H<sub>35</sub>F<sub>2</sub>N<sub>5</sub>O<sub>5</sub>).



- GLC
- 
- Chemical structure of  $\alpha$ -D-glucopyranose (GLC) in a chair conformation. The structure shows the anomeric carbon (C1) at the top right, with its hydroxyl group (O1) in the  $\alpha$  position (pointing down). The other carbons are labeled C2(R), C3(S), C4(S), C5(R), and C6. The hydroxyl groups are labeled O1, O2, O3, O4, O5, and O6. The structure is shown with stereochemistry: C1(R), C2(R), C3(S), C4(S), C5(R), and C6(S).


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	L	1	Total	C	O	0	0
			12	6	6		

- 



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	L	1	Total	C	O	0	0
			11	6	5		

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	U	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	H	154	Total 154	O 154	0	0
10	L	63	Total 63	O 63	0	0
10	T	58	Total 58	O 58	0	0
10	U	53	Total 53	O 53	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: COAGULATION FACTOR VII PRECURSOR




#### • Molecule 2: COAGULATION FACTOR VII PRECURSOR



#### • Molecule 3: TISSUE FACTOR PRECURSOR



#### • Molecule 4: TISSUE FACTOR PRECURSOR





  
  
P210

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.99Å 81.30Å 125.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.72 19.54 – 1.73	Depositor EDS
% Data completeness (in resolution range)	88.8 (50.00-1.72) 92.0 (19.54-1.73)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	-0.06 (at 1.72Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.229 , (Not available) 0.231 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.6	Xtriage
Anisotropy	0.391	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 51.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5058	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GIL, FUC, CA, CGU, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	H	0.27	0/2024	0.65	1/2755 (0.0%)
2	L	0.26	0/1011	0.53	0/1349
3	T	0.29	0/631	0.68	1/860 (0.1%)
4	U	0.27	0/958	0.63	0/1299
All	All	0.27	0/4624	0.63	2/6263 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	199	HIS	N-CA-C	-6.55	93.30	111.00
3	T	20	LYS	N-CA-C	-5.07	97.31	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1974	0	1950	48	0
2	L	1117	0	970	34	0
3	T	615	0	598	9	0
4	U	940	0	911	25	0
5	H	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	L	4	0	0	0	0
6	H	42	0	33	3	0
7	L	12	0	12	2	0
8	L	11	0	12	4	0
9	U	14	0	13	1	0
10	H	154	0	0	3	0
10	L	63	0	0	3	0
10	T	58	0	0	2	0
10	U	53	0	0	4	0
All	All	5058	0	4499	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:68:LEU:HD12	1:H:112:VAL:HG11	1.37	1.04
2:L:60:SER:HB3	8:L:203:FUC:O1	1.59	1.01
2:L:74:PRO:HA	4:U:135:ARG:HH12	1.38	0.87
2:L:31:PHE:HB2	2:L:37:THR:HG22	1.61	0.81
2:L:60:SER:HB3	8:L:203:FUC:C1	2.10	0.81

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	H	252/254 (99%)	238 (94%)	12 (5%)	2 (1%)	19 6
2	L	127/142 (89%)	122 (96%)	5 (4%)	0	100 100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	T	73/75 (97%)	72 (99%)	1 (1%)	0	100	100
4	U	112/116 (97%)	107 (96%)	5 (4%)	0	100	100
All	All	564/587 (96%)	539 (96%)	23 (4%)	2 (0%)	34	18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	219	GLY
1	H	170(F)	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	216/216 (100%)	208 (96%)	8 (4%)	34	14
2	L	113/114 (99%)	111 (98%)	2 (2%)	59	41
3	T	70/70 (100%)	69 (99%)	1 (1%)	67	52
4	U	109/109 (100%)	108 (99%)	1 (1%)	78	69
All	All	508/509 (100%)	496 (98%)	12 (2%)	49	29

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	247	ARG
2	L	36	ARG
4	U	199	ASN
2	L	110	ARG
1	H	84	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	233	GLN
1	H	239	GLN
4	U	199	ASN
3	T	31	ASN
4	U	171	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	CGU	L	14	5,2	9,11,12	1.19	1 (11%)	10,14,16	1.45	1 (10%)
2	CGU	L	29	5,2	9,11,12	1.25	1 (11%)	10,14,16	1.41	1 (10%)
2	CGU	L	20	2	9,11,12	1.21	1 (11%)	10,14,16	1.41	1 (10%)
2	CGU	L	26	5,2	9,11,12	1.25	1 (11%)	10,14,16	1.70	2 (20%)
2	CGU	L	35	2	9,11,12	1.17	1 (11%)	10,14,16	1.45	1 (10%)
2	CGU	L	6	2	9,11,12	1.20	1 (11%)	10,14,16	1.34	1 (10%)
2	CGU	L	7	2	9,11,12	1.17	1 (11%)	10,14,16	1.49	1 (10%)
2	CGU	L	16	5,2	9,11,12	1.25	1 (11%)	10,14,16	1.48	1 (10%)
2	CGU	L	19	5,2	9,11,12	1.20	1 (11%)	10,14,16	1.39	1 (10%)
2	CGU	L	25	5,2	9,11,12	1.24	1 (11%)	10,14,16	1.45	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CGU	L	14	5,2	-	2/13/14/16	-
2	CGU	L	29	5,2	-	0/13/14/16	-
2	CGU	L	20	2	-	1/13/14/16	-
2	CGU	L	26	5,2	-	7/13/14/16	-
2	CGU	L	35	2	-	3/13/14/16	-
2	CGU	L	6	2	-	4/13/14/16	-
2	CGU	L	7	2	-	1/13/14/16	-
2	CGU	L	16	5,2	-	6/13/14/16	-
2	CGU	L	19	5,2	-	2/13/14/16	-
2	CGU	L	25	5,2	-	0/13/14/16	-

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	25	CGU	OE22-CD2	-2.48	1.22	1.30
2	L	20	CGU	OE22-CD2	-2.44	1.22	1.30
2	L	6	CGU	OE22-CD2	-2.44	1.22	1.30
2	L	19	CGU	OE22-CD2	-2.43	1.22	1.30
2	L	14	CGU	OE22-CD2	-2.42	1.22	1.30

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	26	CGU	CB-CG-CD2	-3.96	105.07	113.11
2	L	16	CGU	CB-CG-CD2	-3.62	105.76	113.11
2	L	7	CGU	CB-CG-CD2	-3.47	106.06	113.11
2	L	35	CGU	CB-CG-CD2	-3.35	106.32	113.11
2	L	14	CGU	CB-CG-CD2	-3.31	106.39	113.11

There are no chirality outliers.

5 of 26 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	L	7	CGU	O-C-CA-CB
2	L	20	CGU	N-CA-CB-CG
2	L	26	CGU	CA-CB-CG-CD1
2	L	35	CGU	N-CA-CB-CG
2	L	35	CGU	OE12-CD1-CG-CD2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	L	35	CGU	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 5 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	GLC	L	202	-	12,12,12	0.28	0	17,17,17	0.33	0
9	NAG	U	301	4	14,14,15	0.55	0	17,19,21	0.66	0
6	GIL	H	302	-	44,44,44	1.67	11 (25%)	58,63,63	1.40	7 (12%)
8	FUC	L	203	-	11,11,11	0.32	0	16,16,16	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GLC	L	202	-	-	0/2/22/22	0/1/1/1
9	NAG	U	301	4	-	3/6/23/26	0/1/1/1
6	GIL	H	302	-	-	16/37/37/37	0/3/3/3
8	FUC	L	203	-	-	-	0/1/1/1

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	H	302	GIL	C12-C11	3.85	1.43	1.38
6	H	302	GIL	C15-N3	3.20	1.36	1.32
6	H	302	GIL	C23-C24	2.89	1.43	1.39

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	H	302	GIL	C13-C12	2.88	1.44	1.39
6	H	302	GIL	C11-N3	2.62	1.36	1.32

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	H	302	GIL	C16-N4-C10	5.21	127.64	116.87
6	H	302	GIL	O1-C11-C12	3.87	122.52	117.67
6	H	302	GIL	C17-C10-N4	2.73	117.10	112.09
6	H	302	GIL	O2-C15-C14	-2.67	114.32	117.67
6	H	302	GIL	O1-C11-N3	-2.50	115.42	119.61

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	H	302	GIL	C12-C11-O1-C5
6	H	302	GIL	O5-C8-N5-C9
6	H	302	GIL	C30-C27-C9-N5
9	U	301	NAG	C8-C7-N2-C2
9	U	301	NAG	O7-C7-N2-C2


There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	L	202	GLC	2	0
9	U	301	NAG	1	0
6	H	302	GIL	3	0
8	L	203	FUC	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

The following chains have linkage breaks:

Mol	Chain	Number of breaks
4	U	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	U	158:TRP	C	163:SER	N	9.35

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	H	254/254 (100%)	0.64	26 (10%) <b>6</b> <b>7</b>	10, 19, 39, 61	1 (0%)
2	L	129/142 (90%)	1.04	24 (18%) <b>1</b> <b>1</b>	16, 27, 44, 51	0
3	T	75/75 (100%)	0.37	1 (1%) <b>77</b> <b>81</b>	12, 18, 25, 30	0
4	U	116/116 (100%)	1.16	20 (17%) <b>1</b> <b>1</b>	15, 23, 50, 55	1 (0%)
All	All	574/587 (97%)	0.80	71 (12%) <b>4</b> <b>4</b>	10, 22, 41, 61	2 (0%)

The worst 5 of 71 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	U	137	ASN	8.5
4	U	181	LYS	7.8
1	H	248	PRO	7.6
4	U	182	GLY	7.2
2	L	9	ARG	6.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CGU	L	35	12/13	<b>0.42</b>	<b>0.47</b>	39,51,53,54	0
2	CGU	L	14	12/13	0.71	0.29	37,46,48,49	0
2	CGU	L	20	12/13	0.72	0.34	44,50,52,53	0
2	CGU	L	19	12/13	0.75	0.33	46,51,54,54	0
2	CGU	L	6	12/13	0.81	0.23	26,34,37,38	0
2	CGU	L	7	12/13	0.81	0.20	30,33,36,38	0
2	CGU	L	26	12/13	0.85	0.20	23,29,33,35	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CGU	L	16	12/13	0.86	0.17	34,35,37,37	0
2	CGU	L	25	12/13	0.87	0.20	25,26,28,29	0
2	CGU	L	29	12/13	0.90	0.13	25,27,30,30	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
8	FUC	L	203	11/11	0.67	0.19	35,36,39,42	0
7	GLC	L	202	12/12	0.70	0.19	50,52,52,52	0
9	NAG	U	301	14/15	0.81	0.21	24,31,41,41	0
5	CA	L	205	1/1	0.86	0.14	85,85,85,85	0
6	GIL	H	302	42/42	0.88	0.14	14,25,34,36	0
5	CA	L	206	1/1	0.95	0.14	45,45,45,45	0
5	CA	L	204	1/1	0.96	0.15	46,46,46,46	0
5	CA	L	201	1/1	0.97	0.05	28,28,28,28	0
5	CA	H	301	1/1	0.98	0.13	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around GIL H 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.