Full wwPDB X-ray Structure Validation Report (i)

Jun 16, 2024 – 01:07 PM EDT

PDB ID	:	5C7Z
Title	:	AP2 Mu2 adaptin C-terminal domain complexed with integrin alpha-4 peptide
Authors	:	Owen, D.J.; Evans, P.R.; Wilson, T.A.
Deposited on	:	2015-06-25
Resolution	:	2.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	3% 74% 12% • 10%
2	B	8	62% 12% 25%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2175 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AP-2 complex subunit mu.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	258	2078	1335	366	363	14	0	0	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	236A	MET	-	insertion	UNP P84092
A	236B	GLU	-	insertion	UNP P84092
A	236C	GLN	-	insertion	UNP P84092
A	236D	LYS	-	insertion	UNP P84092
A	236E	LEU	-	insertion	UNP P84092
A	236F	ILE	-	insertion	UNP P84092
A	236G	SER	-	insertion	UNP P84092
A	236H	GLU	-	insertion	UNP P84092
A	236I	GLU	-	insertion	UNP P84092
A	236J	ASP	-	insertion	UNP P84092
A	236K	LEU	-	insertion	UNP P84092

- Molecule 2 is a protein called Integrin alpha-4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	6	52	35	8	9	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	39	39	39	0	0
3	B	6	6	6	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: AP-2 complex subunit mu

Chain A:



• Molecule 2: Integrin alpha-4

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α, β, γ	125.65Å 125.65Å 74.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	74.00 – 2.77 61.42 – 2.77	Depositor EDS
% Data completeness (in resolution range)	99.3 (74.00-2.77) 99.3 (61.42-2.77)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$< I/\sigma(I) >$ [1]	2.09 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, R_{free}	0.174 , 0.224 0.182 , 0.227	Depositor DCC
R_{free} test set	893 reflections (5.25%)	wwPDB-VP
Wilson B-factor (Å^2)	58.0	Xtriage
Anisotropy	0.117	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 46.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.035 for h,-h-k,-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	2175	wwPDB-VP
Average B, all atoms (Å^2)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.96	0/2119	1.07	9/2848 (0.3%)
2	B	1.16	0/52	1.17	1/68 (1.5%)
All	All	0.96	0/2171	1.08	10/2916 (0.3%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	253	ARG	NE-CZ-NH1	6.69	123.65	120.30
1	A	202	MET	CG-SD-CE	5.98	109.77	100.20
1	A	294	ARG	NE-CZ-NH2	-5.65	117.47	120.30
1	A	189	GLY	N-CA-C	5.62	127.16	113.10
2	B	3	LYS	CD-CE-NZ	5.58	124.55	111.70
1	A	170	ARG	NE-CZ-NH1	5.51	123.06	120.30
1	A	434	ARG	NE-CZ-NH2	-5.42	117.59	120.30
1	A	162	ARG	NE-CZ-NH1	5.35	122.97	120.30
1	A	269	ASP	CB-CG-OD2	-5.29	113.54	118.30
1	A	199	ARG	NE-CZ-NH1	5.16	122.88	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2078	0	2170	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	52	0	59	0	0
3	A	39	0	0	2	0
3	B	6	0	0	0	0
All	All	2175	0	2229	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:LYS:O	1:A:260:GLU:N	2.17	0.78
1:A:218:ASP:OD2	1:A:261:ARG:NH1	2.32	0.62
1:A:399:LEU:HD23	1:A:400:LYS:N	2.16	0.60
1:A:323:ARG:NH2	1:A:348:GLU:OE1	2.38	0.53
1:A:380:LYS:HG2	1:A:381:TRP:N	2.25	0.52
1:A:256:LYS:C	1:A:260:GLU:N	2.68	0.47
1:A:163:ARG:HG2	3:A:536:HOH:O	2.15	0.47
1:A:399:LEU:HD23	1:A:399:LEU:C	2.35	0.47
1:A:294:ARG:HD3	3:A:521:HOH:O	2.17	0.45
1:A:212:CYS:HA	1:A:405:LYS:O	2.17	0.45
1:A:278:ARG:HG3	1:A:278:ARG:HH11	1.84	0.43
1:A:249:HIS:HD1	1:A:251:CYS:HG	1.68	0.42
1:A:209:MET:N	1:A:408:GLU:OE1	2.47	0.42
1:A:311:PHE:O	1:A:312:LYS:C	2.58	0.42
1:A:265:PHE:CD1	1:A:265:PHE:N	2.88	0.42

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/288 (88%)	234 (93%)	16 (6%)	2 (1%)	19	47
2	B	4/8 (50%)	4 (100%)	0	0	100	100
All	All	256/296 (86%)	238 (93%)	16 (6%)	2 (1%)	19	47

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	189	GLY
1	A	376	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/260 (90%)	210 (90%)	23 (10%)	8	21
2	B	6/8 (75%)	6 (100%)	0	100	100
All	All	239/268 (89%)	216 (90%)	23 (10%)	8	22

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	ARG
1	A	180	SER
1	A	188	GLN
1	A	197	SER
1	A	213	LYS
1	A	221	VAL
1	A	240	SER
1	A	241	ILE
1	A	253	ARG
1	A	256	LYS
1	A	261	ARG
1	A	276	ARG
1	A	292	LEU
1	A	300	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	304	LYS
1	A	309	SER
1	A	323	ARG
1	A	331	SER
1	A	341	LYS
1	A	376	ASN
1	A	377	ASP
1	A	380	LYS
1	A	434	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	188	GLN
1	A	376	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	258/288 (89%)	0.36	9 (3%) 44 38	36, 57, 101, 125	0
2	B	6/8 (75%)	0.13	0 100 100	43, 62, 82, 86	0
All	All	264/296 (89%)	0.35	9 (3%) 45 39	36, 57, 100, 125	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	238	LYS	4.9
1	A	407	PHE	3.3
1	A	221	VAL	3.1
1	A	220	ILE	2.5
1	A	260	GLU	2.4
1	A	266	ILE	2.3
1	A	211	GLU	2.3
1	A	239	GLN	2.3
1	A	263	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.