Full wwPDB X-ray Structure Validation Report (i)

Dec 14, 2023 – 07:34 am GMT

PDB ID	:	4CED
Title	:	Interrogating HIV integrase for compounds that bind- a SAMPL challenge
Authors	:	Peat, T.S.
Deposited on	:	2013-11-11
Resolution	:	1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	183	2% 66% 15% • 19%
1	B	183	% 70% 11% 19%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 2820 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INTEGRASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	149	1270	816	215	235	4	0	16	0
1	B	149	1269	816	215	234	4	0	16	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	30	MET	-	expression tag	UNP Q76353
A	31	GLY	-	expression tag	UNP Q76353
A	32	SER	-	expression tag	UNP Q76353
A	33	SER	-	expression tag	UNP Q76353
A	34	HIS	-	expression tag	UNP Q76353
A	35	HIS	-	expression tag	UNP Q76353
A	36	HIS	-	expression tag	UNP Q76353
A	37	HIS	-	expression tag	UNP Q76353
A	38	HIS	-	expression tag	UNP Q76353
A	39	HIS	-	expression tag	UNP Q76353
A	40	SER	-	expression tag	UNP Q76353
A	41	SER	-	expression tag	UNP Q76353
A	42	GLY	-	expression tag	UNP Q76353
A	43	LEU	-	expression tag	UNP Q76353
A	44	VAL	-	expression tag	UNP Q76353
A	45	PRO	-	expression tag	UNP Q76353
A	46	ARG	-	expression tag	UNP Q76353
A	47	GLY	-	expression tag	UNP Q76353
A	48	SER	-	expression tag	UNP Q76353
A	49	HIS	-	expression tag	UNP Q76353
A	56	SER	CYS	engineered mutation	UNP Q76353
A	139	ASP	PHE	engineered mutation	UNP Q76353
A	185	HIS	PHE	engineered mutation	UNP Q76353
B	30	MET	-	expression tag	UNP Q76353
B	31	GLY	-	expression tag	UNP Q76353

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	32	SER	-	expression tag	UNP Q76353
B	33	SER	-	expression tag	UNP Q76353
B	34	HIS	-	expression tag	UNP Q76353
B	35	HIS	-	expression tag	UNP Q76353
B	36	HIS	-	expression tag	UNP Q76353
B	37	HIS	-	expression tag	UNP Q76353
B	38	HIS	-	expression tag	UNP Q76353
B	39	HIS	-	expression tag	UNP Q76353
B	40	SER	-	expression tag	UNP Q76353
B	41	SER	-	expression tag	UNP Q76353
B	42	GLY	-	expression tag	UNP Q76353
B	43	LEU	-	expression tag	UNP Q76353
B	44	VAL	-	expression tag	UNP Q76353
B	45	PRO	-	expression tag	UNP Q76353
B	46	ARG	-	expression tag	UNP Q76353
B	47	GLY	-	expression tag	UNP Q76353
B	48	SER	-	expression tag	UNP Q76353
B	49	HIS	-	expression tag	UNP Q76353
B	56	SER	CYS	engineered mutation	UNP Q76353
B	139	ASP	PHE	engineered mutation	UNP Q76353
B	185	HIS	PHE	engineered mutation	UNP Q76353

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	B	1	Total Cl 1 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0

- Molecule 5 is (2S)-6-[[[2-(furan-2-ylmethylcarbamoyl)phenyl]methyl-prop-2-enyl-amino]methyl]-2-(3-hydroxy-3-oxopropyl)-2,3-dihydro-1,4-benzodioxine-5-carboxylic acid (three-letter code: 9NS) (formula: $C_{29}H_{30}N_2O_8$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	B	1	39	29	2	8	0	0
			Total	C	N	O		
5	B	1	39	29	2	8	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	90	90	90	0	0
			Total	O		
6	B	69	69	69	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: INTEGRASE

Chain A:



- Molecule 1: INTEGRASE

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α, β, γ	70.67Å 70.67Å 66.65Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.08 – 1.75 45.08 – 1.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.08-1.75) 100.0 (45.08-1.75)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.11 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.168 , 0.194 0.167 , 0.193	Depositor DCC
R_{free} test set	1904 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å^2)	19.1	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 37.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.024 for -h,-k,l 0.479 for h,-h-k,-l 0.024 for -k,-h,-l	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	2820	wwPDB-VP
Average B, all atoms (Å^2)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, 9NS, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.61	0/1312	0.61	0/1780
1	B	0.62	0/1314	0.60	0/1781
All	All	0.61	0/2626	0.60	0/3561

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1270	0	1298	32	0
1	B	1269	0	1303	27	0
2	A	15	0	0	2	0
2	B	15	0	0	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	6	0	8	1	0
4	B	6	0	8	1	0
5	B	78	0	56	5	0
6	A	90	0	0	1	0
6	B	69	0	0	0	0
All	All	2820	0	2673	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:139:ASP:OD1	1:B:141[B]:ILE:HG22	1.40	1.16
1:A:141[B]:ILE:HG23	1:A:143:TYR:CE2	1.84	1.11
1:A:141[B]:ILE:CG2	1:A:143:TYR:CE2	2.38	1.06
1:B:141[B]:ILE:CG2	1:B:143:TYR:CE2	2.53	0.90
1:A:141[B]:ILE:HD12	1:A:142:PRO:HD2	1.54	0.88
1:B:141[B]:ILE:HG23	1:B:143:TYR:CE2	2.09	0.87
1:A:139:ASP:OD1	1:A:141[B]:ILE:HG22	1.80	0.82
1:B:123:SER:OG	4:B:1214:GOL:H12	1.78	0.81
1:A:141[B]:ILE:HG21	1:A:143:TYR:CE2	2.15	0.81
1:B:141[B]:ILE:HG21	1:B:143:TYR:CE2	2.19	0.78
1:A:141[B]:ILE:HG21	1:A:143:TYR:CZ	2.19	0.77
1:B:139:ASP:OD1	1:B:141[B]:ILE:CG2	2.28	0.76
1:A:123:SER:OG	4:A:1214:GOL:H12	1.87	0.75
1:A:87[B]:GLU:OE2	1:B:103[B]:LYS:NZ	2.23	0.68
1:A:202[B]:ASP:OD1	1:B:209:GLN:NE2	2.29	0.65
1:A:102[B]:LEU:HD11	5:B:1216:9NS:H9	1.80	0.63
1:A:141[B]:ILE:HG23	1:A:143:TYR:CD2	2.33	0.63
1:A:57:SER:N	1:A:58:PRO:HD2	2.15	0.60
1:B:141[B]:ILE:CG2	1:B:143:TYR:HE2	2.13	0.60
1:B:141[B]:ILE:HD12	1:B:142:PRO:HD2	1.83	0.59
1:B:102[B]:LEU:HD11	5:B:1215:9NS:H9	1.84	0.58
1:B:141[B]:ILE:HG12	1:B:143:TYR:CZ	2.39	0.57
1:A:115[B]:THR:HG21	1:A:120:ASN:HB3	1.87	0.56
1:A:115[B]:THR:CG2	1:A:120:ASN:HB3	2.37	0.54
1:B:141[B]:ILE:HG21	1:B:143:TYR:CZ	2.45	0.51
1:A:57:SER:HA	1:A:60[B]:ILE:HG12	1.93	0.51
1:A:71:LYS:NZ	6:A:2011:HOH:O	2.43	0.51
1:B:103[A]:LYS:HE3	2:B:1212:SO4:O4	2.12	0.50
1:B:115[B]:THR:CG2	1:B:120:ASN:HB3	2.42	0.49
1:B:95:GLN:HG3	5:B:1215:9NS:C7	2.43	0.49
1:A:127:LYS:NZ	1:A:137:GLN:HE22	2.11	0.48
1:A:62[B]:GLN:NE2	1:A:151[B]:ILE:HG13	2.29	0.47
1:B:141[B]:ILE:HG23	1:B:143:TYR:HE2	1.69	0.46
1:A:65:CYS:SG	1:A:72:VAL:HG13	2.56	0.45
1:B:65:CYS:SG	1:B:72:VAL:HG13	2.56	0.45
1:A:92[A]:GLU:OE2	1:A:116:ASP:OD2	2.35	0.44
1:A:141[B]:ILE:CG2	1:A:143:TYR:HE2	2.15	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202[B]:ASP:CG	1:B:209:GLN:HE22	2.21	0.44
1:B:115[B]:THR:HG21	1:B:120:ASN:HB3	2.00	0.44
1:A:57:SER:N	1:A:58:PRO:CD	2.80	0.43
1:B:181:PHE:O	1:B:185:HIS:HD2	2.01	0.43
1:A:115[B]:THR:HG21	1:A:120:ASN:CB	2.49	0.43
1:A:103[B]:LYS:NZ	1:B:87[B]:GLU:OE2	2.50	0.43
1:B:205:ALA:O	1:B:209:GLN:HG2	2.19	0.42
1:A:103[A]:LYS:HE3	2:A:1212:SO4:O1	2.20	0.42
1:A:156:LYS:HB3	1:A:156:LYS:HE2	1.70	0.42
1:A:60[B]:ILE:O	1:A:79[B]:VAL:HG22	2.19	0.42
5:B:1216:9NS:C6	5:B:1216:9NS:H271	2.50	0.42
1:B:92[A]:GLU:OE2	1:B:116:ASP:OD2	2.38	0.42
1:B:171:HIS:HD2	2:B:1211:SO4:O1	2.03	0.42
1:A:141[B]:ILE:HG23	1:A:143:TYR:HE2	1.68	0.41
1:A:169:ALA:HB1	1:A:174:THR:HB	2.02	0.41
1:A:115[B]:THR:CG2	1:A:120:ASN:CB	2.99	0.41
1:B:115[B]:THR:CG2	1:B:120:ASN:CB	2.99	0.41
1:B:139:ASP:CG	1:B:141[B]:ILE:HG22	2.30	0.40
1:A:171:HIS:HD2	2:A:1211:SO4:O3	2.04	0.40
5:B:1215:9NS:C6	5:B:1215:9NS:H271	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	161/183 (88%)	161 (100%)	0	0	100	100
1	B	161/183 (88%)	161 (100%)	0	0	100	100
All	All	322/366 (88%)	322 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/150 (93%)	137 (99%)	2 (1%)	67	52
1	B	139/150 (93%)	138 (99%)	1 (1%)	84	75
All	All	278/300 (93%)	275 (99%)	3 (1%)	71	60

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	148	GLN
1	A	156	LYS
1	B	152	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	GLN
1	A	155	ASN
1	A	171	HIS
1	B	95	GLN
1	B	171	HIS
1	B	185	HIS
1	B	209	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GOL	B	1214	-	5,5,5	0.46	0	5,5,5	0.39	0
2	SO4	A	1212	-	4,4,4	0.13	0	6,6,6	0.26	0
2	SO4	B	1211	-	4,4,4	0.19	0	6,6,6	0.29	0
2	SO4	A	1210	-	4,4,4	0.09	0	6,6,6	0.35	0
5	9NS	B	1216	-	39,42,42	1.06	4 (10%)	49,57,57	1.47	9 (18%)
4	GOL	A	1214	-	5,5,5	0.48	0	5,5,5	0.35	0
5	9NS	B	1215	-	39,42,42	1.09	4 (10%)	49,57,57	1.36	9 (18%)
2	SO4	A	1211	-	4,4,4	0.13	0	6,6,6	0.36	0
2	SO4	B	1210	-	4,4,4	0.14	0	6,6,6	0.27	0
2	SO4	B	1212	-	4,4,4	0.32	0	6,6,6	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	1214	-	-	2/4/4/4	-
5	9NS	B	1216	-	-	4/27/38/38	0/4/4/4
4	GOL	A	1214	-	-	2/4/4/4	-
5	9NS	B	1215	-	-	5/27/38/38	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1216	9NS	C26-C16	-2.87	1.48	1.51
5	B	1215	9NS	C26-C16	-2.70	1.48	1.51
5	B	1215	9NS	C25-N31	2.57	1.52	1.47
5	B	1215	9NS	C27-N31	2.53	1.51	1.47
5	B	1216	9NS	C27-N31	2.30	1.51	1.47
5	B	1216	9NS	C19-N30	2.25	1.38	1.33
5	B	1215	9NS	C19-N30	2.24	1.38	1.33
5	B	1216	9NS	C25-N31	2.18	1.51	1.47

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	B	1215	9NS	C29-C23-C22	-3.69	108.05	113.57
5	B	1216	9NS	C13-C25-N31	-3.47	106.67	112.75
5	B	1216	9NS	C29-C28-C21	-3.35	103.61	112.51
5	B	1216	9NS	C18-C27-N31	3.31	119.45	112.89
5	B	1216	9NS	C29-C23-C22	-3.20	108.78	113.57
5	B	1216	9NS	C25-N31-C24	3.12	116.18	110.72
5	B	1215	9NS	C29-C28-C21	-3.01	104.52	112.51
5	B	1215	9NS	C13-C25-N31	-2.99	107.51	112.75
5	B	1216	9NS	O39-C23-C29	2.96	111.31	105.76
5	B	1215	9NS	O39-C23-C29	2.48	110.41	105.76
5	B	1215	9NS	C27-N31-C25	2.36	116.81	111.52
5	B	1216	9NS	O33-C21-O36	2.28	128.98	123.30
5	B	1215	9NS	C25-N31-C24	2.27	114.70	110.72
5	B	1216	9NS	C12-C24-N31	2.25	116.68	112.75
5	B	1215	9NS	C18-C27-N31	2.21	117.26	112.89
5	B	1215	9NS	O33-C21-O36	2.18	128.74	123.30
5	B	1215	9NS	O36-C21-C28	-2.18	116.06	123.08
5	B	1216	9NS	O36-C21-C28	-2.14	116.22	123.08

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1214	GOL	O1-C1-C2-C3
4	B	1214	GOL	O1-C1-C2-O2
4	B	1214	GOL	O1-C1-C2-C3
5	B	1215	9NS	C18-C27-N31-C25
5	B	1215	9NS	C13-C25-N31-C27
4	A	1214	GOL	O1-C1-C2-O2
5	B	1215	9NS	C18-C27-N31-C24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	B	1216	9NS	C18-C27-N31-C24
5	B	1216	9NS	C13-C25-N31-C27
5	B	1216	9NS	C15-C11-C20-O32
5	B	1215	9NS	C15-C11-C20-O35
5	B	1216	9NS	C15-C11-C20-O35
5	B	1215	9NS	C15-C11-C20-O32

There are no ring outliers.

8 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1214	GOL	1	0
2	A	1212	SO4	1	0
2	B	1211	SO4	1	0
5	B	1216	9NS	2	0
4	A	1214	GOL	1	0
5	B	1215	9NS	3	0
2	A	1211	SO4	1	0
2	B	1212	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 9NS B 1216



Bond lengths



Bond angles



Torsions



Rings

Ligand 9NS B 1215



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	149/183 (81%)	-0.30	3 (2%)	65	72	11, 17, 32, 43	21 (14%)
1	B	149/183 (81%)	-0.33	1 (0%)	87	92	11, 17, 32, 41	20 (13%)
All	All	298/366 (81%)	-0.32	4 (1%)	77	83	11, 17, 32, 43	41 (13%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	194	TYR	3.0
1	A	193	GLY	2.7
1	A	188	LYS	2.3
1	B	194	TYR	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	CL	A	1213	1/1	0.79	0.10	62,62,62,62	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	GOL	A	1214	6/6	0.86	0.12	35,37,38,42	0
4	GOL	B	1214	6/6	0.86	0.12	35,37,38,41	0
2	SO4	B	1212	5/5	0.89	0.15	28,30,34,34	5
2	SO4	A	1212	5/5	0.89	0.14	30,33,36,36	5
3	CL	B	1213	1/1	0.90	0.10	63,63,63,63	0
5	9NS	B	1215	39/39	0.93	0.11	16,27,33,34	0
5	9NS	B	1216	39/39	0.93	0.12	16,28,36,37	0
2	SO4	A	1210	5/5	0.95	0.11	37,38,40,41	5
2	SO4	B	1210	5/5	0.96	0.11	39,39,41,41	5
2	SO4	B	1211	5/5	0.97	0.10	27,29,30,31	5
2	SO4	A	1211	5/5	0.98	0.07	25,27,29,29	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 9NS B 1215:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 9NS B 1216:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.