Full wwPDB X-ray Structure Validation Report (i)

Sep 27, 2023 – 01:23 AM EDT

PDB ID	:	6CNZ
Title	:	Crystal Structure of Chorismate Mutase from Burkholderia thailandensis
Authors	:	Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on	:	2018-03-09
Resolution	:	2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	.% 84% 5% 11%
1	B	180	82% 8% 10%
1	C	180	2% 81% 10% 9%
1	D	180	.% 82% 7% • 11%
1	E	180	.% 85% • • 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	180	11% 81% 7% 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	B	305	-	-	X	-

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7800 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chorismate mutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	161	1219	763	227	227	2	0	0	0
			Total	C	N	O	S			
1	B	162	1220	762	230	226	2	0	2	0
			Total	C	N	O	S			
1	C	163	1223	762	229	230	2	0	0	0
			Total	C	N	O	S			
1	D	160	1210	758	226	224	2	0	0	0
			Total	C	N	O	S			
1	E	160	1200	750	220	228	2	0	0	0
			Total	C	N	O	S			
1	F	158	1154	728	207	217	2	0	0	0

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	MET	-	initiating methionine	UNP A0A1B4JXW9
A	24	ALA	-	expression tag	UNP A0A1B4JXW9
A	25	HIS	-	expression tag	UNP A0A1B4JXW9
A	26	HIS	-	expression tag	UNP A0A1B4JXW9
A	27	HIS	-	expression tag	UNP A0A1B4JXW9
A	28	HIS	-	expression tag	UNP A0A1B4JXW9
A	29	HIS	-	expression tag	UNP A0A1B4JXW9
A	30	HIS	-	expression tag	UNP A0A1B4JXW9
A	31	MET	-	expression tag	UNP A0A1B4JXW9
B	23	MET	-	initiating methionine	UNP A0A1B4JXW9
B	24	ALA	-	expression tag	UNP A0A1B4JXW9
B	25	HIS	-	expression tag	UNP A0A1B4JXW9
B	26	HIS	-	expression tag	UNP A0A1B4JXW9
B	27	HIS	-	expression tag	UNP A0A1B4JXW9
B	28	HIS	-	expression tag	UNP A0A1B4JXW9
B	29	HIS	-	expression tag	UNP A0A1B4JXW9
B	30	HIS	-	expression tag	UNP A0A1B4JXW9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	31	MET	-	expression tag	UNP A0A1B4JXW9
C	23	MET	-	initiating methionine	UNP A0A1B4JXW9
C	24	ALA	-	expression tag	UNP A0A1B4JXW9
C	25	HIS	-	expression tag	UNP A0A1B4JXW9
C	26	HIS	-	expression tag	UNP A0A1B4JXW9
C	27	HIS	-	expression tag	UNP A0A1B4JXW9
C	28	HIS	-	expression tag	UNP A0A1B4JXW9
C	29	HIS	-	expression tag	UNP A0A1B4JXW9
C	30	HIS	-	expression tag	UNP A0A1B4JXW9
C	31	MET	-	expression tag	UNP A0A1B4JXW9
D	23	MET	-	initiating methionine	UNP A0A1B4JXW9
D	24	ALA	-	expression tag	UNP A0A1B4JXW9
D	25	HIS	-	expression tag	UNP A0A1B4JXW9
D	26	HIS	-	expression tag	UNP A0A1B4JXW9
D	27	HIS	-	expression tag	UNP A0A1B4JXW9
D	28	HIS	-	expression tag	UNP A0A1B4JXW9
D	29	HIS	-	expression tag	UNP A0A1B4JXW9
D	30	HIS	-	expression tag	UNP A0A1B4JXW9
D	31	MET	-	expression tag	UNP A0A1B4JXW9
E	23	MET	-	initiating methionine	UNP A0A1B4JXW9
E	24	ALA	-	expression tag	UNP A0A1B4JXW9
E	25	HIS	-	expression tag	UNP A0A1B4JXW9
E	26	HIS	-	expression tag	UNP A0A1B4JXW9
E	27	HIS	-	expression tag	UNP A0A1B4JXW9
E	28	HIS	-	expression tag	UNP A0A1B4JXW9
E	29	HIS	-	expression tag	UNP A0A1B4JXW9
E	30	HIS	-	expression tag	UNP A0A1B4JXW9
E	31	MET	-	expression tag	UNP A0A1B4JXW9
F	23	MET	-	initiating methionine	UNP A0A1B4JXW9
F	24	ALA	-	expression tag	UNP A0A1B4JXW9
F	25	HIS	-	expression tag	UNP A0A1B4JXW9
F	26	HIS	-	expression tag	UNP A0A1B4JXW9
F	27	HIS	-	expression tag	UNP A0A1B4JXW9
F	28	HIS	-	expression tag	UNP A0A1B4JXW9
F	29	HIS	-	expression tag	UNP A0A1B4JXW9
F	30	HIS	-	expression tag	UNP A0A1B4JXW9
F	31	MET	-	expression tag	UNP A0A1B4JXW9

- Molecule 2 is NITRATE ION (three-letter code: NO3) (formula: NO_3).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	N	O			
2	A	1	4	1	3		0	0
2	A	1	4	1	3		0	0
2	A	1	4	1	3		0	0
2	B	1	4	1	3		0	0
2	B	1	4	1	3		0	0
2	B	1	4	1	3		0	0
2	C	1	4	1	3		0	0
2	C	1	4	1	3		0	0
2	C	1	4	1	3		0	0
2	C	1	4	1	3		0	0
2	C	1	4	1	3		0	0
2	D	1	4	1	3		0	0
2	D	1	4	1	3		0	0
2	D	1	4	1	3		0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	N	O		
2	E	1	4	1	3	0	0
			Total	N	O		
2	E	1	4	1	3	0	0
			Total	N	O		
2	F	1	4	1	3	0	0
			Total	N	O		
2	F	1	4	1	3	0	0
			Total	N	O		
2	F	1	4	1	3	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	B	1	4	2	2	0	0
			Total	C	O		
3	B	1	4	2	2	0	0
			Total	C	O		
3	B	1	4	2	2	0	0
			Total	C	O		
3	C	1	4	2	2	0	0
			Total	C	O		
3	C	1	4	2	2	0	0
			Total	C	O		
3	D	1	4	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	E	1	4	2	2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	108	108	108	0	0
			Total	O		
4	B	111	111	111	0	0
			Total	O		
4	C	102	102	102	0	0
			Total	O		
4	D	60	60	60	0	0
			Total	O		
4	E	55	55	55	0	0
			Total	O		
4	F	34	34	34	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Chorismate mutase

Chain A:



• Molecule 1: Chorismate mutase

Chain B:



• Molecule 1: Chorismate mutase

Chain C:



• Molecule 1: Chorismate mutase

Chain D:



• Molecule 1: Chorismate mutase

Chain E:



• Molecule 1: Chorismate mutase

Chain F:

11%
81%
7%
12%


MET ALA HIS HIS HIS HIS HIS MET ALA ASP GLY ASP D36 L42 P71 R72 L76 L77 T78 D79 V80 V89 D90 P91 A92 T96 F97 F98 Q101 T141 I145 L148 A149 R150 V151 A152 P153 D156 A157 P158 D159 C160 P161 T173 H191 V192 C193
ALA ALA GLY GLY ALA SER ALA VAL GLY

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	51.71Å 121.38Å 88.67Å 90.00° 99.99° 90.00°	Depositor
Resolution (Å)	31.23 – 2.15 46.96 – 2.15	Depositor EDS
% Data completeness (in resolution range)	98.7 (31.23-2.15) 98.7 (46.96-2.15)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.62 (at 2.16Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.164 , 0.216 0.165 , 0.216	Depositor DCC
R_{free} test set	1968 reflections (3.41%)	wwPDB-VP
Wilson B-factor (Å^2)	31.9	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 45.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	7800	wwPDB-VP
Average B, all atoms (Å^2)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.16% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NO3, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.40	0/1246	0.53	0/1707
1	B	0.43	0/1256	0.56	0/1719
1	C	0.40	0/1249	0.58	1/1710 (0.1%)
1	D	0.38	0/1237	0.52	0/1695
1	E	0.34	0/1227	0.49	0/1683
1	F	0.31	0/1181	0.45	0/1625
All	All	0.38	0/7396	0.52	1/10139 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	174	LEU	CA-CB-CG	5.94	128.96	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	35	ASP	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1219	0	1212	6	0
1	B	1220	0	1210	12	0
1	C	1223	0	1213	8	0
1	D	1210	0	1203	8	0
1	E	1200	0	1174	6	0
1	F	1154	0	1113	6	0
2	A	12	0	0	0	0
2	B	12	0	0	2	0
2	C	20	0	0	0	0
2	D	12	0	0	1	0
2	E	8	0	0	0	0
2	F	12	0	0	0	0
3	B	12	0	18	4	0
3	C	8	0	12	0	0
3	D	4	0	6	0	0
3	E	4	0	6	0	0
4	A	108	0	0	0	0
4	B	111	0	0	0	0
4	C	102	0	0	0	0
4	D	60	0	0	1	0
4	E	55	0	0	1	0
4	F	34	0	0	0	0
All	All	7800	0	7167	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:69:ASP:HB3	3:B:305:EDO:H12	1.73	0.69
1:F:98:PHE:HA	1:F:101:GLN:HG2	1.78	0.64
1:E:93:TYR:CE1	1:E:155:ARG:HD3	2.33	0.63
1:D:150:ARG:NH2	1:F:173:THR:OG1	2.31	0.62
1:B:49:ARG:NH2	2:B:302:NO3:O1	2.25	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:76:LEU:HD23	1:D:102:ILE:HD11	1.87	0.57
1:B:73:GLU:OE2	3:B:305:EDO:O2	2.24	0.55
1:B:134:ARG:NH2	2:B:303:NO3:O2	2.34	0.55
1:D:65:LYS:NZ	4:D:401:HOH:O	2.24	0.55
1:D:134:ARG:NE	2:D:303:NO3:O1	2.37	0.53
1:F:141:THR:O	1:F:145:ILE:HG12	2.10	0.52
1:D:116:TRP:CZ2	1:D:122:PRO:HG3	2.45	0.51
1:B:177:TYR:N	1:B:182:LYS:HE3	2.27	0.50
1:E:142:GLN:OE1	4:E:401:HOH:O	2.20	0.49
1:D:83:ARG:HB2	1:D:145:ILE:HD12	1.95	0.49
1:A:83:ARG:HH12	1:A:142:GLN:HB2	1.77	0.48
1:C:58:HIS:CD2	1:C:122:PRO:HG2	2.48	0.48
1:A:81:GLU:OE1	1:A:95:ARG:NH1	2.45	0.47
1:B:174:LEU:HG	1:C:150:ARG:NH1	2.28	0.47
1:F:152:ALA:HB3	1:F:153:PRO:HD3	1.97	0.47
1:B:88:GLY:HA2	1:E:150:ARG:NH2	2.30	0.45
1:B:176:ARG:HA	1:B:182:LYS:NZ	2.31	0.45
1:C:154:LEU:HD12	1:C:157:ALA:HB3	1.99	0.45
1:A:173:THR:OG1	1:E:150:ARG:NH2	2.48	0.44
1:D:100:ASP:OD2	1:D:191:HIS:ND1	2.43	0.44
1:B:72:ARG:HD2	3:B:305:EDO:O1	2.17	0.44
1:C:73:GLU:OE2	1:C:106:LYS:HE3	2.17	0.44
1:C:116:TRP:CD2	1:C:122:PRO:HD3	2.53	0.44
1:A:169:ALA:O	1:A:173:THR:HG23	2.18	0.43
1:D:160:CYS:HB3	1:D:161:PRO:HD3	2.01	0.43
1:B:73:GLU:HG3	3:B:305:EDO:H11	1.99	0.43
1:E:100:ASP:OD2	1:E:191:HIS:ND1	2.36	0.43
1:C:176:ARG:HA	1:C:182:LYS:NZ	2.33	0.43
1:A:160:CYS:HB3	1:A:161:PRO:HD3	2.01	0.42
1:B:89:VAL:HG22	1:B:149:ALA:HB2	2.00	0.42
1:C:177:TYR:N	1:C:182:LYS:HE3	2.35	0.42
1:C:160:CYS:HB3	1:C:161:PRO:HD3	2.01	0.41
1:E:159:ASP:OD1	1:E:159:ASP:N	2.43	0.41
1:F:42:LEU:HD22	1:F:151:VAL:HG21	2.01	0.41
1:A:144:LEU:HD23	1:A:144:LEU:HA	1.94	0.41
1:B:76:LEU:HD23	1:B:102:ILE:HD11	2.02	0.41
1:F:89:VAL:HG22	1:F:149:ALA:HB2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/180 (88%)	157 (99%)	2 (1%)	0	100	100
1	B	162/180 (90%)	159 (98%)	3 (2%)	0	100	100
1	C	161/180 (89%)	157 (98%)	4 (2%)	0	100	100
1	D	158/180 (88%)	155 (98%)	3 (2%)	0	100	100
1	E	158/180 (88%)	156 (99%)	2 (1%)	0	100	100
1	F	156/180 (87%)	155 (99%)	1 (1%)	0	100	100
All	All	954/1080 (88%)	939 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/137 (90%)	123 (100%)	0	100	100
1	B	123/137 (90%)	123 (100%)	0	100	100
1	C	123/137 (90%)	120 (98%)	3 (2%)	49	51
1	D	122/137 (89%)	121 (99%)	1 (1%)	81	86
1	E	121/137 (88%)	120 (99%)	1 (1%)	81	86
1	F	112/137 (82%)	111 (99%)	1 (1%)	78	83
All	All	724/822 (88%)	718 (99%)	6 (1%)	81	86

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	36	ASP
1	C	155	ARG
1	C	170	ASN
1	D	145	ILE
1	E	155	ARG
1	F	72	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	170	ASN
1	F	107	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NO3	E	303	-	1,3,3	0.75	0	0,3,3	-	-
2	NO3	E	301	-	1,3,3	0.52	0	0,3,3	-	-
2	NO3	F	303	-	1,3,3	0.72	0	0,3,3	-	-
2	NO3	C	307	-	1,3,3	0.70	0	0,3,3	-	-
3	EDO	D	304	-	3,3,3	0.51	0	2,2,2	0.32	0
2	NO3	A	302	-	1,3,3	0.69	0	0,3,3	-	-
2	NO3	B	303	-	1,3,3	0.67	0	0,3,3	-	-
2	NO3	B	301	-	1,3,3	0.73	0	0,3,3	-	-
2	NO3	D	302	-	1,3,3	0.58	0	0,3,3	-	-
3	EDO	B	304	-	3,3,3	0.45	0	2,2,2	0.30	0
2	NO3	B	302	-	1,3,3	0.64	0	0,3,3	-	-
2	NO3	D	303	-	1,3,3	0.71	0	0,3,3	-	-
2	NO3	C	303	-	1,3,3	0.92	0	0,3,3	-	-
2	NO3	C	301	-	1,3,3	0.91	0	0,3,3	-	-
3	EDO	B	306	-	3,3,3	0.56	0	2,2,2	0.11	0
3	EDO	E	302	-	3,3,3	0.48	0	2,2,2	0.31	0
2	NO3	F	302	-	1,3,3	0.74	0	0,3,3	-	-
2	NO3	A	301	-	1,3,3	0.74	0	0,3,3	-	-
3	EDO	B	305	-	3,3,3	0.45	0	2,2,2	0.17	0
2	NO3	C	306	-	1,3,3	0.80	0	0,3,3	-	-
3	EDO	C	305	-	3,3,3	0.43	0	2,2,2	0.47	0
2	NO3	A	303	-	1,3,3	0.75	0	0,3,3	-	-
2	NO3	F	301	-	1,3,3	0.65	0	0,3,3	-	-
3	EDO	C	304	-	3,3,3	0.46	0	2,2,2	0.40	0
2	NO3	D	301	-	1,3,3	0.84	0	0,3,3	-	-
2	NO3	C	302	-	1,3,3	0.64	0	0,3,3	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	305	-	-	1/1/1/1	-
3	EDO	B	304	-	-	1/1/1/1	-
3	EDO	C	305	-	-	0/1/1/1	-
3	EDO	C	304	-	-	0/1/1/1	-
3	EDO	B	306	-	-	0/1/1/1	-
3	EDO	E	302	-	-	0/1/1/1	-
3	EDO	D	304	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	304	EDO	O1-C1-C2-O2
3	B	305	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	303	NO3	1	0
2	B	302	NO3	1	0
2	D	303	NO3	1	0
3	B	305	EDO	4	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	161/180 (89%)	-0.21	1 (0%) 89 91	21, 35, 57, 79	0
1	B	162/180 (90%)	-0.16	0 100 100	18, 32, 56, 93	0
1	C	163/180 (90%)	-0.02	4 (2%) 57 65	21, 32, 63, 88	0
1	D	160/180 (88%)	-0.09	1 (0%) 89 91	28, 42, 71, 83	0
1	E	160/180 (88%)	0.02	2 (1%) 77 82	27, 46, 74, 89	0
1	F	158/180 (87%)	0.62	19 (12%) 4 6	31, 54, 89, 111	0
All	All	964/1080 (89%)	0.02	27 (2%) 53 62	18, 40, 75, 111	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	89	VAL	5.2
1	F	156	ASP	4.5
1	F	92	ALA	4.4
1	F	80	VAL	4.3
1	F	158	PRO	4.3
1	F	76	LEU	4.0
1	C	180	ALA	3.5
1	F	153	PRO	3.4
1	F	90	ASP	3.3
1	F	191	HIS	3.1
1	C	177	TYR	3.1
1	F	161	PRO	3.0
1	F	160	CYS	2.9
1	F	159	ASP	2.7
1	C	118	ALA	2.7
1	F	96	THR	2.6
1	C	120	HIS	2.6
1	F	78	THR	2.5
1	F	192	VAL	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	F	148	LEU	2.5
1	F	91	PRO	2.4
1	F	71	PRO	2.4
1	F	77	LEU	2.3
1	E	120	HIS	2.2
1	E	118	ALA	2.1
1	A	195	ALA	2.0
1	D	85	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	NO3	E	303	4/4	0.74	0.20	72,76,77,78	0
2	NO3	A	303	4/4	0.77	0.20	68,82,84,84	0
2	NO3	F	303	4/4	0.78	0.19	86,93,94,95	0
3	EDO	C	304	4/4	0.80	0.14	64,64,67,68	0
3	EDO	B	306	4/4	0.83	0.25	61,65,70,70	0
2	NO3	A	302	4/4	0.83	0.24	66,68,69,73	0
3	EDO	B	305	4/4	0.84	0.22	41,49,51,63	0
2	NO3	F	302	4/4	0.84	0.13	62,68,69,70	0
2	NO3	B	303	4/4	0.84	0.19	69,71,71,73	0
3	EDO	D	304	4/4	0.85	0.19	43,51,60,67	0
3	EDO	B	304	4/4	0.90	0.21	45,46,48,50	4
2	NO3	D	303	4/4	0.91	0.17	55,60,63,63	0
2	NO3	C	307	4/4	0.91	0.18	99,100,100,100	0
2	NO3	D	302	4/4	0.91	0.30	57,62,62,63	4
2	NO3	C	306	4/4	0.93	0.13	67,68,69,69	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	NO3	C	302	4/4	0.93	0.13	46,53,54,55	0
2	NO3	C	303	4/4	0.93	0.12	44,44,47,49	0
2	NO3	B	302	4/4	0.95	0.13	35,52,52,62	0
3	EDO	E	302	4/4	0.95	0.24	54,58,63,70	0
2	NO3	E	301	4/4	0.96	0.12	47,50,50,53	0
3	EDO	C	305	4/4	0.96	0.15	40,40,44,44	0
2	NO3	C	301	4/4	0.97	0.13	33,42,44,45	0
2	NO3	B	301	4/4	0.98	0.15	40,45,46,46	0
2	NO3	D	301	4/4	0.98	0.10	44,44,45,46	0
2	NO3	F	301	4/4	0.98	0.11	49,53,55,58	0
2	NO3	A	301	4/4	0.99	0.11	33,35,38,38	0

6.5 Other polymers (i)

There are no such residues in this entry.