



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 10, 2026 – 11:15 AM UTC

PDB ID : 9CPD / pdb_00009cpd
BMRB ID : 31188
Title : Structures of small molecules bound to RNA repeat expansions that cause Huntington's disease-like 2 and myotonic dystrophy type 1
Authors : Chen, J.L.; Taghavi, A.; Disney, M.D.; Fountain, M.A.; Childs-Disney, J.L.
Deposited on : 2024-07-18

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR



The overall completeness of chemical shifts assignment is 17%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
RNA backbone	8273	777

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	13	
1	B	13	

2 Ensemble composition and analysis ⓘ

This entry contains 20 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.


3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 873 atoms, of which 304 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called RNA (5'-R(*GP*AP*CP*AP*GP*CP*UP*GP*CP*UP*GP*UP*C)-3').

Mol	Chain	Residues	Atoms						Trace
1	A	13	Total	C	H	N	O	P	0
			415	123	142	48	90	12	
1	B	13	Total	C	H	N	O	P	0
			415	123	142	48	90	12	

- Molecule 2 is 4-[(3-methoxyphenyl)amino]-2-methylquinoline-6-carboximidamide (CCD ID: MQC) (formula: C₁₈H₁₈N₄O) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				
2	B	1	Total	C	H	N	O
			43	18	20	4	1

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble


These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: RNA (5'-R(*GP*AP*CP*AP*GP*CP*UP*GP*CP*UP*GP*UP*C)-3')

Chain A: 



- Molecule 1: RNA (5'-R(*GP*AP*CP*AP*GP*CP*UP*GP*CP*UP*GP*UP*C)-3')

Chain B: 



4.2 Residue scores for the representative (author defined) model from the NMR ensemble


The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: RNA (5'-R(*GP*AP*CP*AP*GP*CP*UP*GP*CP*UP*GP*UP*C)-3')

Chain A: 



- Molecule 1: RNA (5'-R(*GP*AP*CP*AP*GP*CP*UP*GP*CP*UP*GP*UP*C)-3')

Chain B: 



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	structure calculation	20

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	96
Number of shifts mapped to atoms	96
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	17%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MQC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.92±0.03	0±0/304 (0.0± 0.0%)	1.15±0.03	1±1/472 (0.2± 0.2%)
1	B	0.92±0.02	0±0/304 (0.0± 0.0%)	1.15±0.03	0±0/472 (0.0± 0.1%)
All	All	0.92	0/12160 (0.0%)	1.15	25/18880 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.9±0.7
1	B	0.0±0.0	2.8±1.1
All	All	0	94

There are no bond-length outliers.

5 of 7 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	7	U	O4'-C1'-C2'	-6.53	101.07	107.60	16	11
1	A	7	U	N1-C1'-C2'	6.26	121.39	112.00	17	8
1	A	6	C	O4'-C4'-C3'	5.82	109.82	104.00	4	2
1	A	3	C	C4'-C3'-C2'	-5.17	97.42	102.60	14	1
1	B	18	G	O4'-C4'-C3'	5.14	109.14	104.00	16	1

There are no chirality outliers.

5 of 13 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	B	21	G	Sidechain	19
1	A	5	G	Sidechain	17
1	A	8	G	Sidechain	16
1	B	18	G	Sidechain	16
1	B	20	U	Sidechain	11

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	273	142	142	0±0
2	B	23	20	0	0±0
All	All	11380	6080	5678	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:20:U:C6	2:B:101:MQC:C12	0.40	3.05	14	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

6.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	12/13 (92%)	1±0 (8±0%)	0±0 (1±3%)	0.64±0.00

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	B	12/13 (92%)	0±0 (0±0%)	0±0 (0±0%)	0.60±0.00
All	All	480/520 (92%)	20 (4%)	2 (0%)	0.62

The overall RNA backbone suiteness is 0.61.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	7	U	20

All unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	6	C	2

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	MQC	B	101	-	25,25,25	1.46±0.07	4±1 (15±3%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles

that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	MQC	B	101	-	31,35,35	1.29±0.12	4±1 (11±2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MQC	B	101	-	-	0±0,10,10,10	0±0,3,3,3

5 of 8 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	B	101	MQC	C6-N2	3.77	1.40	1.33	1	20
2	B	101	MQC	C1-N1	3.21	1.41	1.27	9	20
2	B	101	MQC	C18-C2	2.77	1.42	1.37	11	11
2	B	101	MQC	C1-N4	2.74	1.41	1.33	14	16
2	B	101	MQC	C8-C9	2.47	1.42	1.38	11	5

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.


Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	101	MQC	C10-N3-C9	4.88	137.01	126.36	8	20
2	B	101	MQC	C9-C17-C5	4.56	120.93	118.01	10	18
2	B	101	MQC	C18-C17-C9	3.82	120.89	123.67	10	17
2	B	101	MQC	C17-C5-N2	2.78	119.85	122.80	10	6
2	B	101	MQC	C8-C6-N2	2.59	119.65	122.60	5	12

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 17% for the well-defined parts and 17% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	96
Number of shifts mapped to atoms	96
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 17%, i.e. 81 atoms were assigned a chemical shift out of a possible 490. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Sugar	48/286 (17%)	48/156 (31%)	0/130 (0%)	0/0 (—%)
Base	33/204 (16%)	33/126 (26%)	0/44 (0%)	0/34 (0%)
Overall	81/490 (17%)	81/282 (29%)	0/174 (0%)	0/34 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots

No *random coil index*(RCI) plot could be generated from the current chemical shift list. RCI is only applicable to proteins

8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	349
Intra-residue ($ i-j =0$)	161
Sequential ($ i-j =1$)	133
Medium range ($ i-j >1$ and $ i-j <5$)	0
Long range ($ i-j \geq 5$)	0
Inter-chain	23
Hydrogen bond restraints	32
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	12.9
Number of long range restraints per residue ¹	0.0

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	None	None
0.2-0.5 (Medium)	None	None
>0.5 (Large)	17.0	17.05

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis ⓘ


9.1 Summary of distance violations ⓘ

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue ($i-j =0$)	161	46.1	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	52	14.9	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	102	29.2	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	7	2.0	0	0.0	0.0	0	0.0	0.0
Sequential ($i-j =1$)	133	38.1	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	14	4.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	93	26.6	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	26	7.4	0	0.0	0.0	0	0.0	0.0
Medium range ($i-j >1$ & $i-j <5$)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Long range ($i-j \geq 5$)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Inter-chain	23	6.6	17	73.9	4.9	17	73.9	4.9
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	18	5.2	13	72.2	3.7	13	72.2	3.7
Sidechain-Sidechain	5	1.4	4	80.0	1.1	4	80.0	1.1
Hydrogen bond	32	9.2	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	349	100.0	17	4.9	4.9	17	4.9	4.9
Backbone-Backbone	66	18.9	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	213	61.0	13	6.1	3.7	13	6.1	3.7
Sidechain-Sidechain	70	20.1	4	5.7	1.1	4	5.7	1.1

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	0	0	0	0	17	17	14.03	16.92	1.16	13.55
2	0	0	0	0	17	17	14.04	17.0	1.16	13.48
3	0	0	0	0	17	17	14.02	17.02	1.16	13.62
4	0	0	0	0	17	17	14.0	16.98	1.15	13.37
5	0	0	0	0	17	17	14.01	16.91	1.16	13.66
6	0	0	0	0	17	17	14.09	16.98	1.14	13.62
7	0	0	0	0	17	17	14.02	17.04	1.15	13.48
8	0	0	0	0	17	17	14.07	16.86	1.18	13.59
9	0	0	0	0	17	17	14.05	16.9	1.13	13.52
10	0	0	0	0	17	17	14.06	17.01	1.15	13.6

Continued on next page...


Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
11	0	0	0	0	17	17	13.98	16.99	1.16	13.46
12	0	0	0	0	17	17	14.04	16.93	1.15	13.61
13	0	0	0	0	17	17	14.01	17.01	1.13	13.57
14	0	0	0	0	17	17	14.03	16.74	1.15	13.65
15	0	0	0	0	17	17	14.05	17.0	1.17	13.55
16	0	0	0	0	17	17	14.01	17.0	1.15	13.49
17	0	0	0	0	17	17	14.06	17.02	1.2	13.44
18	0	0	0	0	17	17	14.09	17.05	1.17	13.61
19	0	0	0	0	17	17	14.01	17.0	1.17	13.56
20	0	0	0	0	17	17	14.12	17.05	1.17	13.85

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model ⓘ



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble ⓘ


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 300(IR:161, SQ:133, MR:0, LR:0, IC:6) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
0	0	0	0	0	0	1	5.0
0	0	0	0	0	0	2	10.0
0	0	0	0	0	0	3	15.0
0	0	0	0	0	0	4	20.0
0	0	0	0	0	0	5	25.0
0	0	0	0	0	0	6	30.0
0	0	0	0	0	0	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	17	17	20	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(3,10)	2:101:B:MQC:H7	1:20:B:U:H6	20	16.97	0.07	17.0
(3,12)	2:101:B:MQC:H7	1:20:B:U:H2'	20	15.44	0.22	15.34
(3,13)	2:101:B:MQC:H7	1:21:B:G:H2'	20	15.29	0.14	15.34
(3,8)	2:101:B:MQC:H6	1:20:B:U:H2'	20	15.27	0.17	15.26
(3,8)	2:101:B:MQC:H6	1:20:B:U:H1'	20	15.27	0.17	15.26
(3,1)	2:101:B:MQC:C15	1:19:B:C:H6	20	15.07	0.32	15.04
(3,16)	2:101:B:MQC:H8	1:19:B:C:H2'	20	14.39	0.12	14.38
(3,6)	2:101:B:MQC:H2	1:6:A:C:H1'	20	14.16	0.13	14.18
(3,4)	2:101:B:MQC:C7	1:6:A:C:H2'	20	13.62	0.21	13.57
(3,4)	2:101:B:MQC:C7	1:6:A:C:H1'	20	13.62	0.21	13.57
(3,9)	2:101:B:MQC:H6	1:21:B:G:H2'	20	13.58	0.13	13.56
(3,11)	2:101:B:MQC:H7	1:19:B:C:H1'	20	13.48	0.12	13.48

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(3,10)	2:101:B:MQC:H7	1:20:B:U:H6	18	17.05
(3,10)	2:101:B:MQC:H7	1:20:B:U:H6	20	17.05
(3,10)	2:101:B:MQC:H7	1:20:B:U:H6	7	17.04
(3,10)	2:101:B:MQC:H7	1:20:B:U:H6	3	17.02
(3,10)	2:101:B:MQC:H7	1:20:B:U:H6	17	17.02
(3,10)	2:101:B:MQC:H7	1:20:B:U:H6	10	17.01
(3,10)	2:101:B:MQC:H7	1:20:B:U:H6	13	17.01
(3,10)	2:101:B:MQC:H7	1:20:B:U:H6	2	17.0
(3,10)	2:101:B:MQC:H7	1:20:B:U:H6	15	17.0
(3,10)	2:101:B:MQC:H7	1:20:B:U:H6	16	17.0

10 Dihedral-angle violation analysis ⓘ

No dihedral-angle restraints found