



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 26, 2026 – 04:40 AM EDT

PDB ID : 9CQH / pdb_00009cqh
Title : CRYSTAL STRUCTURE OF APO CLEAVED N-TERMINAL HIS-TAG
GAGA-DOG HSP47(36-418) IN A C 2 CRYSTAL FORM
Authors : Sheriff, S.
Deposited on : 2024-07-19
Resolution : 2.01 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	<div> <div>16%</div> <div>93% 6% .</div> </div>
1	B	387	<div> <div>14%</div> <div>91% 8% .</div> </div>
1	C	387	<div> <div>19%</div> <div>93% 6% .</div> </div>
1	D	387	<div> <div>31%</div> <div>92% 7% .</div> </div>
1	E	387	<div> <div>17%</div> <div>91% 7% ..</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	387	<div><div></div><div>35%</div><div></div><div>92%</div><div></div><div>6% ..</div></div>
1	G	387	<div><div></div><div>32%</div><div></div><div>93%</div><div></div><div>6% .</div></div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 41762 atoms, of which 20444 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serpin H1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	383	Total	C	H	N	O	S	2879	0	0
			5802	1864	2879	506	540	13			
1	B	382	Total	C	H	N	O	S	2950	1	0
			5911	1885	2950	517	546	13			
1	C	383	Total	C	H	N	O	S	2943	0	0
			5902	1885	2943	515	546	13			
1	D	383	Total	C	H	N	O	S	2935	0	0
			5897	1882	2935	516	551	13			
1	E	383	Total	C	H	N	O	S	2945	1	0
			5910	1887	2945	519	546	13			
1	F	383	Total	C	H	N	O	S	2879	0	0
			5806	1864	2879	506	544	13			
1	G	383	Total	C	H	N	O	S	2897	0	0
			5843	1876	2897	510	547	13			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	32	GLY	-	expression tag	UNP C7C419
A	33	ALA	-	expression tag	UNP C7C419
A	34	GLY	-	expression tag	UNP C7C419
A	35	ALA	-	expression tag	UNP C7C419
B	32	GLY	-	expression tag	UNP C7C419
B	33	ALA	-	expression tag	UNP C7C419
B	34	GLY	-	expression tag	UNP C7C419
B	35	ALA	-	expression tag	UNP C7C419
C	32	GLY	-	expression tag	UNP C7C419
C	33	ALA	-	expression tag	UNP C7C419
C	34	GLY	-	expression tag	UNP C7C419
C	35	ALA	-	expression tag	UNP C7C419
D	32	GLY	-	expression tag	UNP C7C419
D	33	ALA	-	expression tag	UNP C7C419
D	34	GLY	-	expression tag	UNP C7C419

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	35	ALA	-	expression tag	UNP C7C419
E	32	GLY	-	expression tag	UNP C7C419
E	33	ALA	-	expression tag	UNP C7C419
E	34	GLY	-	expression tag	UNP C7C419
E	35	ALA	-	expression tag	UNP C7C419
F	32	GLY	-	expression tag	UNP C7C419
F	33	ALA	-	expression tag	UNP C7C419
F	34	GLY	-	expression tag	UNP C7C419
F	35	ALA	-	expression tag	UNP C7C419
G	32	GLY	-	expression tag	UNP C7C419
G	33	ALA	-	expression tag	UNP C7C419
G	34	GLY	-	expression tag	UNP C7C419
G	35	ALA	-	expression tag	UNP C7C419

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	8	0
			14	3	8	3		
3	D	1	Total	C	H	O	8	0
			14	3	8	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	95	Total	O	0	0
			95	95		
4	B	88	Total	O	0	0
			88	88		
4	C	91	Total	O	0	0
			91	91		
4	D	86	Total	O	0	0
			86	86		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	97	Total 97	O 97	0	0
4	F	85	Total 85	O 85	0	0
4	G	81	Total 81	O 81	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Serpin H1




• Molecule 1: Serpin H1





• Molecule 1: Serpin H1









• Molecule 1: Serpin H1




• Molecule 1: Serpin H1




• Molecule 1: Serpin H1





● Molecule 1: Serpin H1

Chain G: 32% 93% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	180.04Å 115.49Å 187.53Å 90.00° 106.80° 90.00°	Depositor
Resolution (Å)	28.87 – 2.01 28.87 – 2.01	Depositor EDS
% Data completeness (in resolution range)	61.5 (28.87-2.01) 61.5 (28.87-2.01)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 2.01Å)	Xtriage
Refinement program	BUSTER 2.11.8	Depositor
R, R_{free}	0.234 , 0.254 0.226 , 0.246	Depositor DCC
R_{free} test set	7538 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	39.5	Xtriage
Anisotropy	0.043	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	41762	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	2/2984 (0.1%)	0.97	0/4037
1	B	0.75	2/3025 (0.1%)	0.98	1/4087 (0.0%)
1	C	0.76	1/3020 (0.0%)	0.98	1/4080 (0.0%)
1	D	0.76	2/3023 (0.1%)	0.99	1/4087 (0.0%)
1	E	0.77	2/3029 (0.1%)	0.97	0/4094
1	F	0.77	2/2988 (0.1%)	0.99	0/4043
1	G	0.74	2/3007 (0.1%)	0.97	0/4068
All	All	0.76	13/21076 (0.1%)	0.98	3/28496 (0.0%)

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	236	MET	SD-CE	-7.22	1.61	1.79
1	D	236	MET	SD-CE	-6.57	1.63	1.79
1	D	87	ALA	CA-C	6.16	1.60	1.53
1	G	87	ALA	CA-C	6.06	1.60	1.53
1	B	236	MET	SD-CE	-5.93	1.64	1.79

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	288	GLU	CB-CG-CD	7.15	124.76	112.60
1	C	288	GLU	CB-CG-CD	6.42	123.52	112.60
1	B	416	ASP	CA-CB-CG	5.53	118.13	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2923	2879	2879	11	0
1	B	2961	2950	2950	11	0
1	C	2959	2943	2943	7	0
1	D	2962	2935	2935	7	0
1	E	2965	2945	2945	10	0
1	F	2927	2879	2879	11	0
1	G	2946	2897	2897	7	0
2	A	20	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	E	10	0	0	1	0
3	B	6	8	8	0	0
3	D	6	8	8	0	0
4	A	95	0	0	3	0
4	B	88	0	0	1	0
4	C	91	0	0	1	0
4	D	86	0	0	0	0
4	E	97	0	0	1	0
4	F	85	0	0	2	0
4	G	81	0	0	1	0
All	All	21318	20444	20444	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:230:TYR:OH	1:E:413:LYS:HB3	1.75	0.86
1:D:263:LYS:NZ	2:E:501:SO4:O4	2.22	0.72
1:C:230:TYR:OH	1:C:413:LYS:HB3	1.95	0.66
1:A:291:LYS:NZ	4:A:601:HOH:O	2.30	0.64
1:B:209:HIS:HD2	4:B:684:HOH:O	1.83	0.61

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	381/387 (98%)	368 (97%)	13 (3%)	0	100	100
1	B	381/387 (98%)	366 (96%)	12 (3%)	3 (1%)	16	11
1	C	381/387 (98%)	364 (96%)	13 (3%)	4 (1%)	12	8
1	D	381/387 (98%)	370 (97%)	11 (3%)	0	100	100
1	E	382/387 (99%)	366 (96%)	14 (4%)	2 (0%)	24	21
1	F	381/387 (98%)	362 (95%)	19 (5%)	0	100	100
1	G	381/387 (98%)	364 (96%)	16 (4%)	1 (0%)	36	35
All	All	2668/2709 (98%)	2560 (96%)	98 (4%)	10 (0%)	30	27

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	375	GLU
1	B	123	ALA
1	B	374	GLU
1	E	36	LEU
1	G	372	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/328 (92%)	294 (97%)	8 (3%)	40	44
1	B	312/328 (95%)	301 (96%)	11 (4%)	32	32

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	310/328 (94%)	300 (97%)	10 (3%)	34	35
1	D	312/328 (95%)	302 (97%)	10 (3%)	34	35
1	E	310/328 (94%)	298 (96%)	12 (4%)	28	28
1	F	304/328 (93%)	295 (97%)	9 (3%)	36	38
1	G	305/328 (93%)	294 (96%)	11 (4%)	31	31
All	All	2155/2296 (94%)	2084 (97%)	71 (3%)	33	34

5 of 71 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	253	LEU
1	F	391	LEU
1	G	172	SER
1	C	330	ILE
1	C	253	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	274	HIS
1	G	274	HIS
1	E	274	HIS
1	F	298	GLN
1	E	254	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	502	-	5,5,5	0.09	0	5,5,5	0.17	0
2	SO4	A	501	-	4,4,4	0.33	0	6,6,6	0.30	0
2	SO4	B	501	-	4,4,4	0.28	0	6,6,6	0.26	0
2	SO4	A	504	-	4,4,4	0.30	0	6,6,6	0.14	0
3	GOL	D	501	-	5,5,5	0.12	0	5,5,5	0.22	0
2	SO4	A	502	-	4,4,4	0.29	0	6,6,6	0.43	0
2	SO4	E	502	-	4,4,4	0.30	0	6,6,6	0.23	0
2	SO4	E	501	-	4,4,4	0.29	0	6,6,6	1.41	1 (16%)
2	SO4	C	501	-	4,4,4	0.28	0	6,6,6	0.82	0
2	SO4	A	503	-	4,4,4	0.30	0	6,6,6	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	D	501	-	-	0/4/4/4	-
3	GOL	B	502	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	501	SO4	O4-S-O2	-3.14	93.16	109.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	501	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	383/387 (98%)	0.83	61 (15%) 5 4	10, 21, 47, 66	0
1	B	382/387 (98%)	0.76	56 (14%) 6 5	11, 22, 46, 63	1 (0%)
1	C	383/387 (98%)	0.90	73 (19%) 3 3	11, 22, 46, 60	0
1	D	383/387 (98%)	1.27	121 (31%) 1 1	11, 26, 51, 66	0
1	E	383/387 (98%)	0.84	67 (17%) 4 3	11, 22, 45, 57	1 (0%)
1	F	383/387 (98%)	1.56	135 (35%) 1 1	11, 26, 57, 74	0
1	G	383/387 (98%)	1.33	122 (31%) 1 1	11, 25, 55, 70	0
All	All	2680/2709 (98%)	1.07	635 (23%) 2 2	10, 23, 51, 74	2 (0%)

The worst 5 of 635 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	123	ALA	9.6
1	F	122	THR	8.3
1	F	123	ALA	8.2
1	E	120	ASN	7.0
1	A	122	THR	6.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	504	5/5	0.73	0.21	116,116,117,117	0
2	SO4	B	501	5/5	0.76	0.14	100,100,100,100	0
2	SO4	A	502	5/5	0.84	0.12	84,84,84,84	0
2	SO4	A	501	5/5	0.85	0.14	89,89,90,90	0
2	SO4	E	502	5/5	0.87	0.12	111,111,111,111	0
3	GOL	D	501	6/6	0.87	0.14	40,41,41,41	8
2	SO4	A	503	5/5	0.88	0.13	100,100,100,100	0
2	SO4	E	501	5/5	0.93	0.09	61,62,62,62	0
3	GOL	B	502	6/6	0.94	0.09	41,41,42,42	8
2	SO4	C	501	5/5	0.96	0.09	43,43,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.