



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 5, 2026 – 03:54 AM UTC

PDB ID : 9CYN / pdb_00009cyn
Title : Flagellar filament sheath protein FlaA1
Authors : San Martin, F.; Larrieux, N.; Trajtenberg, F.; Buschiazzi, A.
Deposited on : 2024-08-02
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	
1	B	257	
1	C	257	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5528 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endoflagellar filament sheath protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	215	Total	C	N	O	S	0	5	0
			1728	1108	289	329	2			
1	B	210	Total	C	N	O	S	0	0	0
			1641	1053	278	309	1			
1	C	218	Total	C	N	O	S	0	1	0
			1704	1091	288	323	2			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	46	GLY	-	expression tag	UNP Q04UP1
A	47	SER	-	expression tag	UNP Q04UP1
A	48	GLY	-	expression tag	UNP Q04UP1
A	49	SER	-	expression tag	UNP Q04UP1
B	46	GLY	-	expression tag	UNP Q04UP1
B	47	SER	-	expression tag	UNP Q04UP1
B	48	GLY	-	expression tag	UNP Q04UP1
B	49	SER	-	expression tag	UNP Q04UP1
C	46	GLY	-	expression tag	UNP Q04UP1
C	47	SER	-	expression tag	UNP Q04UP1
C	48	GLY	-	expression tag	UNP Q04UP1
C	49	SER	-	expression tag	UNP Q04UP1

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		
2	B	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total	Ca	0	0
			1	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	158	Total	O	0	2
			160	160		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	109	Total 113	O 113	0	4
4	C	143	Total 144	O 144	0	1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Endoflagellar filament sheath protein



• Molecule 1: Endoflagellar filament sheath protein



• Molecule 1: Endoflagellar filament sheath protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.42Å 105.78Å 108.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.82 – 1.82 75.82 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.9 (75.82-1.82) 100.0 (75.82-1.82)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.04 (at 1.82Å)	Xtriage
Refinement program	PHENIX 1.21.1_5286	Depositor
R, R_{free}	0.184 , 0.215 0.184 , 0.215	Depositor DCC
R_{free} test set	4232 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	36.3	Xtriage
Anisotropy	0.376	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5528	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, YCM, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/1777	0.81	1/2416 (0.0%)
1	B	0.67	1/1669 (0.1%)	0.74	0/2264
1	C	0.69	0/1736	0.75	0/2353
All	All	0.71	1/5182 (0.0%)	0.77	1/7033 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	120	THR	CB-CG2	5.14	1.69	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	257	GLU	CB-CG-CD	5.91	122.64	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1728	0	1663	12	0
1	B	1641	0	1557	11	0
1	C	1704	0	1617	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	10	0	0	1	0
3	B	15	0	0	1	0
3	C	10	0	0	0	0
4	A	160	0	0	2	0
4	B	113	0	0	1	0
4	C	144	0	0	0	0
All	All	5528	0	4837	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:49:SER:HB2	1:A:52:LEU:HD12	1.43	1.01
1:A:234:SER:HA	1:A:237[B]:GLN:HE21	1.41	0.85
1:A:233[B]:ASN:ND2	4:A:501:HOH:O	2.20	0.75
1:A:237[A]:GLN:NE2	4:A:502:HOH:O	2.28	0.66
1:B:113:ILE:HD12	1:B:265:GLU:HG3	1.83	0.61
1:C:200:TRP:NE1	1:C:239:LYS:HE3	2.15	0.61
1:B:187:ARG:NH2	3:B:402:SO4:O3	2.21	0.60
1:B:93:VAL:HG22	1:B:102:GLU:HG3	1.90	0.53
1:B:260:TYR:CE1	1:C:274:GLU:OE1	2.62	0.52
1:B:110:PRO:HD2	1:B:113:ILE:HG13	1.92	0.51
1:B:120:THR:O	1:B:122:LYS:HG3	2.13	0.48
1:B:95:GLN:OE1	1:C:234:SER:HA	2.13	0.48
1:A:234:SER:O	1:A:237[B]:GLN:HG2	2.13	0.48
1:A:257:GLU:HG3	1:A:258:THR:N	2.28	0.48
1:C:231:ASP:HB2	1:C:239:LYS:HE2	1.94	0.47
1:B:76:ASP:OD2	1:B:77:LYS:NZ	2.41	0.47
1:B:276:HIS:CD2	4:B:502:HOH:O	2.68	0.47
1:C:174:PRO:HD2	1:C:268:VAL:HG21	1.96	0.46
1:C:174:PRO:O	1:C:175:GLY:O	2.32	0.46
1:C:251[A]:ARG:HB2	1:C:253:ASP:OD1	2.17	0.45
1:C:185:YCM:HB3	1:C:217:TRP:CD2	2.52	0.45
1:B:152:ARG:HA	1:B:152:ARG:HD2	1.82	0.43
1:A:57:VAL:HB	1:A:171:VAL:CG1	2.48	0.43
1:A:254:THR:HA	3:A:402:SO4:O2	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:ASP:CG	1:B:111:GLY:HA3	2.44	0.43
1:C:276:HIS:CD2	1:C:276:HIS:H	2.35	0.43
1:C:284:PHE:O	1:C:285:ASP:HB2	2.18	0.43
1:A:234:SER:HA	1:A:237[B]:GLN:NE2	2.23	0.41
1:A:185:YCM:HD2	1:A:217:TRP:CD1	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/257 (83%)	207 (97%)	6 (3%)	0	100	100
1	B	198/257 (77%)	192 (97%)	6 (3%)	0	100	100
1	C	206/257 (80%)	199 (97%)	7 (3%)	0	100	100
All	All	617/771 (80%)	598 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/220 (84%)	181 (98%)	4 (2%)	45	30
1	B	170/220 (77%)	169 (99%)	1 (1%)	78	74

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	177/220 (80%)	176 (99%)	1 (1%)	78	74
All	All	532/660 (81%)	526 (99%)	6 (1%)	65	56

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	GLN
1	A	100	LEU
1	A	226	GLN
1	A	257	GLU
1	B	144	ARG
1	C	167	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	98	GLN
1	B	80	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	YCM	A	185	1	7,9,10	1.48	2 (28%)	5,10,12	0.48	0
1	YCM	B	185	1	7,9,10	1.11	1 (14%)	5,10,12	1.31	1 (20%)
1	YCM	C	185	1	7,9,10	0.96	0	5,10,12	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	YCM	A	185	1	-	2/6/8/10	-
1	YCM	B	185	1	-	1/6/8/10	-
1	YCM	C	185	1	-	2/6/8/10	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	185	YCM	CD-SG	-2.43	1.75	1.81
1	B	185	YCM	CE-NZ2	2.22	1.40	1.32
1	A	185	YCM	CE-NZ2	2.09	1.39	1.32

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	185	YCM	CE-CD-SG	2.24	120.89	113.81

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	185	YCM	SG-CD-CE-OZ1
1	B	185	YCM	SG-CD-CE-NZ2
1	C	185	YCM	SG-CD-CE-NZ2
1	A	185	YCM	SG-CD-CE-OZ1
1	A	185	YCM	CA-CB-SG-CD

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	185	YCM	1	0
1	C	185	YCM	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	402	-	4,4,4	0.70	0	6,6,6	0.31	0
3	SO4	B	404	-	4,4,4	0.60	0	6,6,6	0.48	0
3	SO4	C	403	-	4,4,4	1.13	0	6,6,6	0.78	0
3	SO4	C	402	-	4,4,4	0.77	0	6,6,6	0.32	0
3	SO4	A	403	-	4,4,4	0.72	0	6,6,6	0.29	0
3	SO4	A	402	-	4,4,4	0.50	0	6,6,6	0.51	0
3	SO4	B	403	-	4,4,4	0.72	0	6,6,6	0.60	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	402	SO4	1	0
3	A	402	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	214/257 (83%)	0.39	28 (13%)	7 7	21, 41, 82, 102	5 (2%)
1	B	209/257 (81%)	0.66	27 (12%)	7 8	31, 47, 89, 121	0
1	C	217/257 (84%)	0.59	29 (13%)	7 7	29, 47, 85, 105	1 (0%)
All	All	640/771 (83%)	0.55	84 (13%)	7 7	21, 45, 86, 121	6 (0%)

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	235	TYR	8.5
1	B	285	ASP	6.4
1	A	277	PHE	5.7
1	A	113	ILE	5.6
1	B	275	VAL	4.9
1	A	84	GLY	4.8
1	B	232	VAL	4.6
1	B	175	GLY	4.5
1	C	118	ALA	4.4
1	B	282	ILE	4.4
1	B	116	VAL	4.4
1	C	232	VAL	4.4
1	A	238	VAL	4.3
1	A	82	ALA	4.0
1	C	81	GLN	4.0
1	B	117	ASP	3.9
1	B	237	GLN	3.9
1	B	272	VAL	3.9
1	C	175	GLY	3.8
1	C	233	ASN	3.8
1	B	238	VAL	3.6
1	A	49	SER	3.6
1	C	281	SER	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	256	GLY	3.5
1	A	114	LYS	3.5
1	C	114	LYS	3.5
1	C	113	ILE	3.5
1	A	47	SER	3.4
1	B	120	THR	3.4
1	C	238	VAL	3.4
1	B	119	GLY	3.4
1	B	51	GLU	3.4
1	C	278	ASP	3.3
1	C	279	GLY	3.3
1	A	235	TYR	3.2
1	B	274	GLU	3.2
1	C	228	VAL	3.2
1	A	237[A]	GLN	3.2
1	C	237	GLN	3.1
1	B	227	GLY	3.1
1	C	46	GLY	3.1
1	C	151	LEU	3.1
1	B	277	PHE	3.1
1	C	153	THR	3.1
1	A	112	ASP	3.0
1	A	166	SER	3.0
1	C	234	SER	3.0
1	C	227	GLY	3.0
1	B	151	LEU	2.9
1	B	233	ASN	2.9
1	C	231	ASP	2.9
1	A	236	PRO	2.8
1	A	234	SER	2.8
1	B	80	GLN	2.7
1	A	86	GLN	2.7
1	A	83	GLN	2.7
1	B	276	HIS	2.7
1	A	119	GLY	2.6
1	C	253	ASP	2.6
1	B	166	SER	2.6
1	B	88	GLN	2.6
1	A	255	SER	2.6
1	A	257	GLU	2.6
1	C	47	SER	2.6
1	A	85	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	115	ASN	2.5
1	B	89	ALA	2.4
1	A	81	GLN	2.4
1	C	274	GLU	2.3
1	C	286	ASP	2.3
1	A	278	ASP	2.3
1	B	255	SER	2.2
1	C	166	SER	2.2
1	C	150	VAL	2.2
1	A	254	THR	2.2
1	A	151	LEU	2.2
1	C	48	GLY	2.1
1	A	52	LEU	2.1
1	B	280	ALA	2.1
1	B	115	ASN	2.1
1	C	226	GLN	2.1
1	A	253	ASP	2.1
1	A	276	HIS	2.0
1	B	66	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	YCM	B	185	10/11	0.97	0.09	32,38,49,52	0
1	YCM	A	185	10/11	0.98	0.07	30,38,49,52	0
1	YCM	C	185	10/11	0.99	0.06	35,38,50,53	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	403	5/5	0.80	0.13	77,80,96,106	0
3	SO4	C	402	5/5	0.80	0.11	71,81,103,118	0
3	SO4	C	403	5/5	0.87	0.18	49,52,93,110	0
3	SO4	B	404	5/5	0.91	0.12	61,67,78,81	0
3	SO4	B	402	5/5	0.93	0.09	70,71,80,81	0
3	SO4	A	403	5/5	0.95	0.08	59,61,66,74	0
2	CA	B	401	1/1	0.99	0.02	34,34,34,34	0
3	SO4	A	402	5/5	0.99	0.06	38,38,41,44	0
2	CA	C	401	1/1	1.00	0.02	34,34,34,34	0
2	CA	A	401	1/1	1.00	0.02	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around CA B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CA C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CA A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.