



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 16, 2020 – 07:46 pm BST

PDB ID : 5D17  
Title : Structure of the C-terminal domain of TnsE at 2.85 resolution  
Authors : Guarne, A.; Caron, J.J.  
Deposited on : 2015-08-03  
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3168 (2.90-2.82)
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RSRZ outliers	127900	3103 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	206	66% 11% 23%
1	B	206	65% 11% 23%
1	C	206	66% 11% 23%
1	D	206	68% 8% 24%
1	E	206	70% 6% 23%
1	F	206	64% 13% 23%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	206	
1	H	206	
1	I	206	
1	K	206	
1	L	206	
1	M	206	

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 14936 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transposon Tn7 transposition protein TnsE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	159	1252	793	217	237	2	3	0	6	0
1	B	158	1251	789	221	236	2	3	0	6	0
1	C	158	1266	797	226	238	2	3	0	6	0
1	D	157	1223	776	215	227	2	3	0	6	0
1	E	158	1234	782	217	230	2	3	0	6	0
1	F	158	1226	777	211	233	2	3	0	6	0
1	G	158	1242	788	216	233	2	3	0	6	0
1	H	156	1236	783	217	231	2	3	0	6	0
1	I	158	1269	799	228	237	2	3	0	6	0
1	K	157	1241	789	214	233	2	3	0	6	0
1	L	158	1209	771	209	224	2	3	0	6	0
1	M	159	1264	796	224	239	2	3	0	6	0

There are 108 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	341	MSE	-	initiating methionine	UNP P05845
A	539	LEU	-	expression tag	UNP P05845
A	540	GLU	-	expression tag	UNP P05845
A	541	HIS	-	expression tag	UNP P05845
A	542	HIS	-	expression tag	UNP P05845

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	543	HIS	-	expression tag	UNP P05845
A	544	HIS	-	expression tag	UNP P05845
A	545	HIS	-	expression tag	UNP P05845
A	546	HIS	-	expression tag	UNP P05845
B	341	MSE	-	initiating methionine	UNP P05845
B	539	LEU	-	expression tag	UNP P05845
B	540	GLU	-	expression tag	UNP P05845
B	541	HIS	-	expression tag	UNP P05845
B	542	HIS	-	expression tag	UNP P05845
B	543	HIS	-	expression tag	UNP P05845
B	544	HIS	-	expression tag	UNP P05845
B	545	HIS	-	expression tag	UNP P05845
B	546	HIS	-	expression tag	UNP P05845
C	341	MSE	-	initiating methionine	UNP P05845
C	539	LEU	-	expression tag	UNP P05845
C	540	GLU	-	expression tag	UNP P05845
C	541	HIS	-	expression tag	UNP P05845
C	542	HIS	-	expression tag	UNP P05845
C	543	HIS	-	expression tag	UNP P05845
C	544	HIS	-	expression tag	UNP P05845
C	545	HIS	-	expression tag	UNP P05845
C	546	HIS	-	expression tag	UNP P05845
D	341	MSE	-	initiating methionine	UNP P05845
D	539	LEU	-	expression tag	UNP P05845
D	540	GLU	-	expression tag	UNP P05845
D	541	HIS	-	expression tag	UNP P05845
D	542	HIS	-	expression tag	UNP P05845
D	543	HIS	-	expression tag	UNP P05845
D	544	HIS	-	expression tag	UNP P05845
D	545	HIS	-	expression tag	UNP P05845
D	546	HIS	-	expression tag	UNP P05845
E	341	MSE	-	initiating methionine	UNP P05845
E	539	LEU	-	expression tag	UNP P05845
E	540	GLU	-	expression tag	UNP P05845
E	541	HIS	-	expression tag	UNP P05845
E	542	HIS	-	expression tag	UNP P05845
E	543	HIS	-	expression tag	UNP P05845
E	544	HIS	-	expression tag	UNP P05845
E	545	HIS	-	expression tag	UNP P05845
E	546	HIS	-	expression tag	UNP P05845
F	341	MSE	-	initiating methionine	UNP P05845
F	539	LEU	-	expression tag	UNP P05845

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	540	GLU	-	expression tag	UNP P05845
F	541	HIS	-	expression tag	UNP P05845
F	542	HIS	-	expression tag	UNP P05845
F	543	HIS	-	expression tag	UNP P05845
F	544	HIS	-	expression tag	UNP P05845
F	545	HIS	-	expression tag	UNP P05845
F	546	HIS	-	expression tag	UNP P05845
G	341	MSE	-	initiating methionine	UNP P05845
G	539	LEU	-	expression tag	UNP P05845
G	540	GLU	-	expression tag	UNP P05845
G	541	HIS	-	expression tag	UNP P05845
G	542	HIS	-	expression tag	UNP P05845
G	543	HIS	-	expression tag	UNP P05845
G	544	HIS	-	expression tag	UNP P05845
G	545	HIS	-	expression tag	UNP P05845
G	546	HIS	-	expression tag	UNP P05845
H	341	MSE	-	initiating methionine	UNP P05845
H	539	LEU	-	expression tag	UNP P05845
H	540	GLU	-	expression tag	UNP P05845
H	541	HIS	-	expression tag	UNP P05845
H	542	HIS	-	expression tag	UNP P05845
H	543	HIS	-	expression tag	UNP P05845
H	544	HIS	-	expression tag	UNP P05845
H	545	HIS	-	expression tag	UNP P05845
H	546	HIS	-	expression tag	UNP P05845
I	341	MSE	-	initiating methionine	UNP P05845
I	539	LEU	-	expression tag	UNP P05845
I	540	GLU	-	expression tag	UNP P05845
I	541	HIS	-	expression tag	UNP P05845
I	542	HIS	-	expression tag	UNP P05845
I	543	HIS	-	expression tag	UNP P05845
I	544	HIS	-	expression tag	UNP P05845
I	545	HIS	-	expression tag	UNP P05845
I	546	HIS	-	expression tag	UNP P05845
K	341	MSE	-	initiating methionine	UNP P05845
K	539	LEU	-	expression tag	UNP P05845
K	540	GLU	-	expression tag	UNP P05845
K	541	HIS	-	expression tag	UNP P05845
K	542	HIS	-	expression tag	UNP P05845
K	543	HIS	-	expression tag	UNP P05845
K	544	HIS	-	expression tag	UNP P05845
K	545	HIS	-	expression tag	UNP P05845

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
K	546	HIS	-	expression tag	UNP P05845
L	341	MSE	-	initiating methionine	UNP P05845
L	539	LEU	-	expression tag	UNP P05845
L	540	GLU	-	expression tag	UNP P05845
L	541	HIS	-	expression tag	UNP P05845
L	542	HIS	-	expression tag	UNP P05845
L	543	HIS	-	expression tag	UNP P05845
L	544	HIS	-	expression tag	UNP P05845
L	545	HIS	-	expression tag	UNP P05845
L	546	HIS	-	expression tag	UNP P05845
M	341	MSE	-	initiating methionine	UNP P05845
M	539	LEU	-	expression tag	UNP P05845
M	540	GLU	-	expression tag	UNP P05845
M	541	HIS	-	expression tag	UNP P05845
M	542	HIS	-	expression tag	UNP P05845
M	543	HIS	-	expression tag	UNP P05845
M	544	HIS	-	expression tag	UNP P05845
M	545	HIS	-	expression tag	UNP P05845
M	546	HIS	-	expression tag	UNP P05845

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O 1 1	0	0
2	B	1	Total O 1 1	0	0
2	C	1	Total O 1 1	0	0
2	D	1	Total O 1 1	0	0
2	E	2	Total O 2 2	0	0
2	F	1	Total O 1 1	0	0
2	G	8	Total O 8 8	0	0
2	H	2	Total O 2 2	0	0
2	I	2	Total O 2 2	0	0
2	K	1	Total O 1 1	0	0

*Continued on next page...*

*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
2	L	1	Total O 1 1	0	0
2	M	2	Total O 2 2	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transposon Tn7 transposition protein TnsE

Chain A:  66% 11% 23%




- Molecule 1: Transposon Tn7 transposition protein TnsE

Chain B:  % 65% 11% 23%



- Molecule 1: Transposon Tn7 transposition protein TnsE


Chain C:  % 66% 11% 23%




- Molecule 1: Transposon Tn7 transposition protein TnsE

Chain D:  % 68% 8% 24%







• Molecule 1: Transposon Tn7 transposition protein TnsE




• Molecule 1: Transposon Tn7 transposition protein TnsE




• Molecule 1: Transposon Tn7 transposition protein TnsE




• Molecule 1: Transposon Tn7 transposition protein TnsE




• Molecule 1: Transposon Tn7 transposition protein TnsE




• Molecule 1: Transposon Tn7 transposition protein TnsE



• Molecule 1: Transposon Tn7 transposition protein TnsE



• Molecule 1: Transposon Tn7 transposition protein TnsE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.33Å 87.12Å 141.62Å 90.00° 95.01° 90.00°	Depositor
Resolution (Å)	39.75 – 2.85 43.21 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.75-2.85) 92.5 (43.21-2.80)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.13 (at 2.81Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.221 , 0.251 0.224 , 0.251	Depositor DCC
$R_{free}$ test set	2000 reflections (3.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	67.4	Xtrriage
Anisotropy	0.313	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 39.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	14936	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	87.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.28 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4175e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.22	0/1272	0.41	0/1718
1	B	0.22	0/1271	0.41	0/1715
1	C	0.21	0/1285	0.42	0/1733
1	D	0.23	0/1243	0.49	2/1680 (0.1%)
1	E	0.22	0/1254	0.42	0/1695
1	F	0.25	0/1246	0.43	0/1686
1	G	0.22	0/1262	0.41	0/1704
1	H	0.21	0/1256	0.42	0/1695
1	I	0.22	0/1289	0.42	0/1737
1	K	0.22	0/1260	0.41	0/1699
1	L	0.22	0/1229	0.48	1/1664 (0.1%)
1	M	0.21	0/1284	0.42	0/1733
All	All	0.22	0/15151	0.43	3/20459 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	517	LEU	CB-CG-CD1	-7.38	98.45	111.00
1	D	517	LEU	CA-CB-CG	6.23	129.63	115.30
1	D	517	LEU	CB-CG-CD1	-6.02	100.77	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1252	0	1200	16	0
1	B	1251	0	1208	16	0
1	C	1266	0	1230	15	0
1	D	1223	0	1177	14	0
1	E	1234	0	1183	11	0
1	F	1226	0	1157	21	0
1	G	1242	0	1197	11	0
1	H	1236	0	1200	16	0
1	I	1269	0	1243	12	0
1	K	1241	0	1209	15	0
1	L	1209	0	1151	13	0
1	M	1264	0	1221	15	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	2	0	0	0	0
2	F	1	0	0	0	0
2	G	8	0	0	0	0
2	H	2	0	0	0	0
2	I	2	0	0	0	0
2	K	1	0	0	0	0
2	L	1	0	0	0	0
2	M	2	0	0	0	0
All	All	14936	0	14376	143	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 143 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:506:ARG:NH2	1:F:531:VAL:O	2.10	0.84
1:G:517:LEU:HB2	1:H:518:GLN:HE21	1.43	0.82
1:I:422:ARG:HH22	1:K:458[B]:LYS:HB2	1.51	0.75
1:F:517:LEU:HG	1:F:518:GLN:N	2.03	0.73
1:I:459[A]:MSE:HE1	1:I:490:ASN:HA	1.71	0.72

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/206 (79%)	160 (98%)	3 (2%)	0	100	100
1	B	162/206 (79%)	161 (99%)	1 (1%)	0	100	100
1	C	162/206 (79%)	159 (98%)	3 (2%)	0	100	100
1	D	161/206 (78%)	158 (98%)	3 (2%)	0	100	100
1	E	162/206 (79%)	159 (98%)	3 (2%)	0	100	100
1	F	162/206 (79%)	161 (99%)	1 (1%)	0	100	100
1	G	162/206 (79%)	161 (99%)	1 (1%)	0	100	100
1	H	160/206 (78%)	159 (99%)	1 (1%)	0	100	100
1	I	162/206 (79%)	159 (98%)	3 (2%)	0	100	100
1	K	161/206 (78%)	156 (97%)	5 (3%)	0	100	100
1	L	162/206 (79%)	161 (99%)	1 (1%)	0	100	100
1	M	163/206 (79%)	162 (99%)	1 (1%)	0	100	100
All	All	1942/2472 (79%)	1916 (99%)	26 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	129/174 (74%)	128 (99%)	1 (1%)	81	93
1	B	131/174 (75%)	128 (98%)	3 (2%)	50	78

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	134/174 (77%)	133 (99%)	1 (1%)	84	94
1	D	125/174 (72%)	124 (99%)	1 (1%)	81	93
1	E	126/174 (72%)	124 (98%)	2 (2%)	62	84
1	F	124/174 (71%)	123 (99%)	1 (1%)	81	93
1	G	128/174 (74%)	127 (99%)	1 (1%)	81	93
1	H	129/174 (74%)	128 (99%)	1 (1%)	81	93
1	I	135/174 (78%)	135 (100%)	0	100	100
1	K	130/174 (75%)	126 (97%)	4 (3%)	40	71
1	L	121/174 (70%)	121 (100%)	0	100	100
1	M	133/174 (76%)	133 (100%)	0	100	100
All	All	1545/2088 (74%)	1530 (99%)	15 (1%)	78	91

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	384	ILE
1	E	477	PHE
1	K	458[B]	LYS
1	D	477	PHE
1	K	458[A]	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	518	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	157/206 (76%)	-0.00	1 (0%) 89 89	52, 79, 120, 142	0
1	B	156/206 (75%)	0.00	3 (1%) 66 64	54, 87, 138, 157	0
1	C	156/206 (75%)	0.17	4 (2%) 56 52	55, 88, 136, 154	0
1	D	155/206 (75%)	0.07	3 (1%) 66 64	53, 93, 132, 144	0
1	E	156/206 (75%)	-0.04	1 (0%) 89 89	59, 95, 132, 147	0
1	F	156/206 (75%)	0.15	3 (1%) 66 64	59, 101, 140, 157	0
1	G	156/206 (75%)	-0.10	1 (0%) 89 89	49, 72, 99, 126	0
1	H	154/206 (74%)	0.11	5 (3%) 47 42	51, 86, 125, 134	0
1	I	156/206 (75%)	0.01	0 100 100	42, 70, 111, 128	0
1	K	155/206 (75%)	0.07	3 (1%) 66 64	41, 84, 127, 147	0
1	L	156/206 (75%)	0.16	5 (3%) 47 42	64, 112, 154, 175	0
1	M	157/206 (76%)	-0.13	1 (0%) 89 89	44, 75, 114, 131	0
All	All	1870/2472 (75%)	0.04	30 (1%) 72 70	41, 85, 134, 175	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	531	VAL	3.5
1	B	519	VAL	3.4
1	H	519	VAL	3.3
1	F	531	VAL	3.3
1	M	465	LEU	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

There are no ligands in this entry.

### 6.5 Other polymers

There are no such residues in this entry.