



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 5, 2025 – 10:08 AM EDT

PDB ID : 9DL0 / pdb\_00009dl0  
Title : Crystal structure of a synthetic Fab (R3H8) in complex with the FRB domain of mTOR  
Authors : O'Leary, K.M.; Slezak, T.; Kossiakoff, A.A.  
Deposited on : 2024-09-10  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.43.1




# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	224	 4% 84% 12% ..
1	H	224	 4% 85% 14% .
2	G	98	 5% 79% 15% 6%
2	I	98	 5% 93% 6% .
3	Y	212	 87% 13%

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Length	Quality of chain	
			5%	20%
3	Z	212	 A horizontal bar chart showing the quality of chain Z. The bar is divided into three segments: a small red segment at the beginning (labeled 5%), a long green segment in the middle (labeled 80%), and a small yellow segment at the end (labeled 20%).	



## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8837 atoms, of which 351 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	218	Total	C	N	O	S	0	0	0
			1618	1024	268	320	6			
1	H	224	Total	C	N	O	S	0	0	0
			1656	1045	275	330	6			

- Molecule 2 is a protein called Serine/threonine-protein kinase mTOR.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	G	92	Total	C	H	N	O	S	173	0	0
			952	497	173	136	139	7			
2	I	98	Total	C	H	N	O	S	178	0	0
			992	517	178	144	146	7			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	2020	SER	-	expression tag	UNP P42345
G	2114	GLY	-	expression tag	UNP P42345
G	2115	GLY	-	expression tag	UNP P42345
G	2116	GLY	-	expression tag	UNP P42345
G	2117	GLY	-	expression tag	UNP P42345
I	2020	SER	-	expression tag	UNP P42345
I	2114	GLY	-	expression tag	UNP P42345
I	2115	GLY	-	expression tag	UNP P42345
I	2116	GLY	-	expression tag	UNP P42345
I	2117	GLY	-	expression tag	UNP P42345

- Molecule 3 is a protein called Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	Y	212	Total	C	N	O	S	0	0	0
			1623	1017	273	328	5			

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	Z	212	Total	C	N	O	S	0	0	0
			1623	1017	273	328	5			

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	50	Total	O	0	0
			50	50		
4	G	29	Total	O	0	0
			29	29		
4	H	56	Total	O	0	0
			56	56		
4	I	44	Total	O	0	0
			44	44		
4	Y	102	Total	O	0	0
			102	102		
4	Z	92	Total	O	0	0
			92	92		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab heavy chain




- Molecule 1: Fab heavy chain



- Molecule 2: Serine/threonine-protein kinase mTOR



- Molecule 2: Serine/threonine-protein kinase mTOR




- Molecule 3: Fab light chain



Chain Y: 

87%

13%




● Molecule 3: Fab light chain

Chain Z: 

5%

80%

20%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.61Å 110.04Å 131.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.31 – 2.00 56.31 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.3 (56.31-2.00) 95.6 (56.31-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.37 (at 2.00Å)	Xtriage
Refinement program	PHENIX (1.21.1)	Depositor
R, $R_{free}$	0.205 , 0.234 0.206 , 0.235	Depositor DCC
$R_{free}$ test set	4447 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.6	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 41.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8837	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/1658	0.55	0/2259
1	H	0.32	0/1697	0.52	0/2312
2	G	0.39	0/801	0.57	0/1078
2	I	0.37	0/836	0.55	0/1121
3	Y	0.41	0/1659	0.64	0/2253
3	Z	0.42	0/1659	0.62	0/2253
All	All	0.37	0/8310	0.58	0/11276

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1618	0	1571	25	0
1	H	1656	0	1610	28	0
2	G	779	173	735	10	0
2	I	814	178	776	9	0
3	Y	1623	0	1582	19	0
3	Z	1623	0	1582	42	0
4	A	50	0	0	1	0
4	G	29	0	0	2	0
4	H	56	0	0	2	0
4	I	44	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	Y	102	0	0	1	0
4	Z	92	0	0	0	0
All	All	8486	351	7856	123	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 123 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:223:LYS:HD3	1:H:223:LYS:H	1.34	0.91
1:A:16:GLN:HG3	1:A:17:PRO:HD2	1.51	0.90
3:Z:3:ILE:HD11	3:Z:94:TRP:CE2	2.08	0.89
1:A:223:LYS:H	1:A:223:LYS:HD2	1.40	0.86
3:Y:184:LYS:O	3:Y:188:GLU:HG3	1.77	0.84

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/224 (96%)	210 (98%)	4 (2%)	0	100	100
1	H	222/224 (99%)	217 (98%)	5 (2%)	0	100	100
2	G	90/98 (92%)	90 (100%)	0	0	100	100
2	I	96/98 (98%)	94 (98%)	2 (2%)	0	100	100
3	Y	210/212 (99%)	203 (97%)	7 (3%)	0	100	100
3	Z	210/212 (99%)	201 (96%)	9 (4%)	0	100	100
All	All	1042/1068 (98%)	1015 (97%)	27 (3%)	0	100	100



There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/186 (97%)	177 (98%)	4 (2%)	47	51
1	H	186/186 (100%)	183 (98%)	3 (2%)	58	64
2	G	80/83 (96%)	80 (100%)	0	100	100
2	I	83/83 (100%)	82 (99%)	1 (1%)	67	73
3	Y	187/187 (100%)	186 (100%)	1 (0%)	86	90
3	Z	187/187 (100%)	185 (99%)	2 (1%)	70	76
All	All	904/912 (99%)	893 (99%)	11 (1%)	67	73

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	I	2094	VAL
3	Y	77	SER
3	Z	200	GLN
3	Z	10	SER
1	H	163	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	184	GLN
3	Y	39	GLN
1	H	212	ASN
3	Y	90	GLN
2	I	2082	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.



## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	218/224 (97%)	0.44	8 (3%) 45 43	30, 46, 69, 92	0
1	H	224/224 (100%)	0.46	9 (4%) 43 41	28, 47, 83, 133	0
2	G	92/98 (93%)	0.30	5 (5%) 32 30	25, 36, 54, 76	0
2	I	98/98 (100%)	0.34	5 (5%) 34 32	24, 35, 61, 69	0
3	Y	212/212 (100%)	0.04	1 (0%) 87 86	27, 38, 58, 76	0
3	Z	212/212 (100%)	0.37	10 (4%) 37 35	23, 40, 83, 92	0
All	All	1056/1068 (98%)	0.33	38 (3%) 46 44	23, 42, 75, 133	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	Z	94	TRP	9.3
3	Z	95	PHE	5.2
1	A	44	PRO	4.2
2	G	2021	ILE	3.5
2	G	2112	SER	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.