



# wwPDB X-ray Structure Validation Summary Report

Jun 19, 2024 – 04:32 AM EDT

PDB ID : 4DOH  
Title : IL20/IL201/IL20R2 Ternary Complex  
Authors : Logsdon, N.J.; Walter, M.R.  
Deposited on : 2012-02-09  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	153	
1	C	153	
2	B	206	
2	D	206	
3	E	221	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	R	221	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	R	302	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8949 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	153	1231	772	224	226	9	0	0	0
1	C	153	1231	772	224	226	9	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	ALA	-	expression tag	UNP Q9NYY1
C	24	ALA	-	expression tag	UNP Q9NYY1

- Molecule 2 is a protein called Interleukin-20 receptor subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	193	1525	977	251	288	9	0	0	0
2	D	193	1525	977	251	288	9	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	29	ALA	-	expression tag	UNP Q6UXL0
B	40	GLN	ASN	engineered mutation	UNP Q6UXL0
B	134	GLN	ASN	engineered mutation	UNP Q6UXL0
B	232	GLU	-	expression tag	UNP Q6UXL0
B	233	GLY	-	expression tag	UNP Q6UXL0
B	234	ARG	-	expression tag	UNP Q6UXL0
D	29	ALA	-	expression tag	UNP Q6UXL0
D	40	GLN	ASN	engineered mutation	UNP Q6UXL0
D	134	GLN	ASN	engineered mutation	UNP Q6UXL0
D	232	GLU	-	expression tag	UNP Q6UXL0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	233	GLY	-	expression tag	UNP Q6UXL0
D	234	ARG	-	expression tag	UNP Q6UXL0

- Molecule 3 is a protein called Interleukin-20 receptor subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	R	204	1655	1055	284	308	8	0	0	0
3	E	204	Total	C	N	O	S	0	0	0
			1655	1055	284	308	8			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	111	ARG	LYS	engineered mutation	UNP Q9UHF4
R	113	ARG	LYS	engineered mutation	UNP Q9UHF4
R	246	ILE	-	expression tag	UNP Q9UHF4
R	247	GLU	-	expression tag	UNP Q9UHF4
R	248	GLY	-	expression tag	UNP Q9UHF4
R	249	ARG	-	expression tag	UNP Q9UHF4
E	111	ARG	LYS	engineered mutation	UNP Q9UHF4
E	113	ARG	LYS	engineered mutation	UNP Q9UHF4
E	246	ILE	-	expression tag	UNP Q9UHF4
E	247	GLU	-	expression tag	UNP Q9UHF4
E	248	GLY	-	expression tag	UNP Q9UHF4
E	249	ARG	-	expression tag	UNP Q9UHF4

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	B	1	Total 14	C 8	N 1	O 5	0	0
4	R	1	Total 14	C 8	N 1	O 5	0	0
4	R	1	Total 14	C 8	N 1	O 5	0	0
4	D	1	Total 14	C 8	N 1	O 5	0	0
4	E	1	Total 14	C 8	N 1	O 5	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	7	Total 7	O 7	0	0
5	B	14	Total 14	O 14	0	0
5	R	6	Total 6	O 6	0	0
5	C	16	Total 16	O 16	0	0
5	D	6	Total 6	O 6	0	0
5	E	8	Total 8	O 8	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-20




- Molecule 1: Interleukin-20




- Molecule 2: Interleukin-20 receptor subunit beta




- Molecule 2: Interleukin-20 receptor subunit beta






• Molecule 3: Interleukin-20 receptor subunit alpha



• Molecule 3: Interleukin-20 receptor subunit alpha



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.11Å 111.76Å 136.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 49.13 – 2.81	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.80) 96.5 (49.13-2.81)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.35 (at 2.81Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.230 , 0.278 0.211 , 0.270	Depositor DCC
$R_{free}$ test set	2006 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	56.5	Xtrriage
Anisotropy	0.312	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 55.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	8949	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/1249	0.66	0/1678
1	C	0.48	0/1249	0.63	0/1678
2	B	0.45	0/1568	0.63	0/2140
2	D	0.46	0/1568	0.65	0/2140
3	E	0.48	0/1701	0.64	0/2322
3	R	0.47	0/1701	0.64	0/2322
All	All	0.47	0/9036	0.64	0/12280

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	D	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	198	GLY	Peptide
2	D	187	GLY	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1231	0	1253	39	0
1	C	1231	0	1253	25	0
2	B	1525	0	1479	58	0
2	D	1525	0	1479	72	0
3	E	1655	0	1618	100	0
3	R	1655	0	1618	90	0
4	B	14	0	13	2	0
4	D	14	0	13	1	0
4	E	14	0	13	4	0
4	R	28	0	26	9	0
5	A	7	0	0	3	0
5	B	14	0	0	0	0
5	C	16	0	0	0	0
5	D	6	0	0	0	0
5	E	8	0	0	1	0
5	R	6	0	0	0	0
All	All	8949	0	8765	377	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 377 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:124:ARG:HG3	1:C:124:ARG:HH11	1.12	1.14
3:E:182:ASN:ND2	4:E:301:NAG:C1	2.14	1.10
3:E:70:THR:HG22	3:E:89:ASN:H	1.18	1.08
2:B:171:ARG:HH22	2:B:198:GLY:HA3	1.19	1.07
1:A:66:ARG:O	1:A:67:ARG:HB3	1.57	1.02

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/153 (99%)	141 (93%)	8 (5%)	2 (1%)	12	36
1	C	151/153 (99%)	142 (94%)	7 (5%)	2 (1%)	12	36
2	B	191/206 (93%)	174 (91%)	14 (7%)	3 (2%)	9	31
2	D	191/206 (93%)	173 (91%)	13 (7%)	5 (3%)	5	18
3	E	202/221 (91%)	178 (88%)	18 (9%)	6 (3%)	4	15
3	R	202/221 (91%)	184 (91%)	14 (7%)	4 (2%)	7	24
All	All	1088/1160 (94%)	992 (91%)	74 (7%)	22 (2%)	7	24

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	R	59	PRO
3	R	122	LYS
1	C	67	ARG
3	E	59	PRO
3	E	122	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/137 (100%)	121 (88%)	16 (12%)	5	16
1	C	137/137 (100%)	123 (90%)	14 (10%)	7	22
2	B	168/177 (95%)	152 (90%)	16 (10%)	8	25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	168/177 (95%)	147 (88%)	21 (12%)	4	14
3	E	186/201 (92%)	163 (88%)	23 (12%)	4	14
3	R	186/201 (92%)	164 (88%)	22 (12%)	5	16
All	All	982/1030 (95%)	870 (89%)	112 (11%)	5	18

5 of 112 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	52	SER
3	E	234	LYS
2	D	52	LEU
3	E	233	GLU
3	E	168	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
3	R	136	GLN
3	E	106	HIS
1	C	40	GLN
3	E	211	ASN
3	E	52	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	E	301	-	14,14,15	1.64	2 (14%)	17,19,21	5.38	3 (17%)
4	NAG	D	301	-	14,14,15	0.55	0	17,19,21	1.99	3 (17%)
4	NAG	R	302	-	14,14,15	0.70	0	17,19,21	1.71	2 (11%)
4	NAG	B	301	-	14,14,15	0.85	0	17,19,21	1.37	3 (17%)
4	NAG	R	301	-	14,14,15	0.59	0	17,19,21	1.46	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	E	301	-	-	4/6/23/26	0/1/1/1
4	NAG	D	301	-	-	1/6/23/26	0/1/1/1
4	NAG	R	302	-	-	4/6/23/26	0/1/1/1
4	NAG	B	301	-	-	3/6/23/26	0/1/1/1
4	NAG	R	301	-	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	301	NAG	O5-C1	-4.98	1.35	1.43
4	E	301	NAG	C1-C2	-2.21	1.49	1.52

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	301	NAG	O5-C1-C2	20.83	143.51	111.29
4	D	301	NAG	C2-N2-C7	-6.33	114.41	122.90

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	301	NAG	C1-O5-C5	-5.48	104.85	112.19
4	R	302	NAG	O5-C1-C2	-4.75	103.95	111.29
4	E	301	NAG	C3-C4-C5	4.34	118.11	110.23

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	R	302	NAG	O7-C7-N2-C2
4	E	301	NAG	O5-C5-C6-O6
4	R	302	NAG	C8-C7-N2-C2
4	R	302	NAG	O5-C5-C6-O6
4	R	302	NAG	C4-C5-C6-O6

There are no ring outliers.

5 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	301	NAG	4	0
4	D	301	NAG	1	0
4	R	302	NAG	7	0
4	B	301	NAG	2	0
4	R	301	NAG	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	153/153 (100%)	-0.21	1 (0%) 87 84	28, 48, 72, 104	0
1	C	153/153 (100%)	-0.31	0 100 100	27, 49, 73, 102	0
2	B	193/206 (93%)	-0.04	2 (1%) 82 77	34, 55, 93, 110	0
2	D	193/206 (93%)	0.20	10 (5%) 27 18	34, 55, 98, 116	0
3	E	204/221 (92%)	0.21	11 (5%) 25 17	41, 67, 114, 134	0
3	R	204/221 (92%)	0.21	9 (4%) 34 24	40, 66, 112, 133	0
All	All	1100/1160 (94%)	0.03	33 (3%) 50 40	27, 57, 101, 134	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	62	GLU	6.6
2	D	117	LEU	6.5
3	R	66	LYS	5.9
3	E	65	VAL	5.3
3	E	62	LEU	5.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	B	301	14/15	0.79	0.37	54,58,61,64	0
4	NAG	R	301	14/15	0.81	0.29	49,51,54,54	0
4	NAG	R	302	14/15	0.84	0.27	49,51,54,54	0
4	NAG	E	301	14/15	0.85	0.23	49,51,54,54	0
4	NAG	D	301	14/15	0.87	0.37	54,58,61,64	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.