



wwPDB X-ray Structure Validation Summary Report ⓘ

May 27, 2025 – 08:47 AM EDT

PDB ID : 9DTF / pdb_00009dtf
Title : Crystal structure of the complex of M. tuberculosis PheRS with cognate precursor tRNA and fragment DDD01008876
Authors : Chang, C.; Michalska, K.; Forte, B.; Baragana, B.; Gilbert, I.H.; Wower, J.; Joachimiak, A.
Deposited on : 2024-09-30
Resolution : 2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1096 (2.46-2.46)
Clashscore	180529	1178 (2.46-2.46)
Ramachandran outliers	177936	1170 (2.46-2.46)
Sidechain outliers	177891	1170 (2.46-2.46)
RSRZ outliers	164620	1096 (2.46-2.46)
RNA backbone	3690	1040 (2.76-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	
1	D	342	
2	B	835	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	835	
3	C	77	
3	F	77	

2 Entry composition i

There are 12 unique types of molecules in this entry. The entry contains 21921 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylalanine-tRNA ligase alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	335	Total	C	N	O	S	0	0	0
			2535	1603	452	471	9			
1	D	338	Total	C	N	O	S	0	8	0
			2662	1679	480	495	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	ALA	-	expression tag	UNP P9WFU3
D	0	ALA	-	expression tag	UNP P9WFU3

- Molecule 2 is a protein called Phenylalanine-tRNA ligase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	834	Total	C	N	O	S	0	12	0
			6329	3976	1150	1181	22			
2	E	806	Total	C	N	O	S	0	10	0
			6083	3821	1108	1135	19			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLN	-	expression tag	UNP P9WFU1
B	-2	SER	-	expression tag	UNP P9WFU1
B	-1	ASN	-	expression tag	UNP P9WFU1
B	0	ALA	-	expression tag	UNP P9WFU1
E	-3	GLN	-	expression tag	UNP P9WFU1
E	-2	SER	-	expression tag	UNP P9WFU1
E	-1	ASN	-	expression tag	UNP P9WFU1
E	0	ALA	-	expression tag	UNP P9WFU1


- Molecule 3 is a RNA chain called tRNA(Phe).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	73	Total 1563	C 695	N 283	O 512	P 73	0	0	0
3	F	73	Total 1563	C 695	N 283	O 512	P 73	0	0	0

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
4	A	1	Total 1	Mg 1	0	0
4	C	2	Total 2	Mg 2	0	0
4	D	1	Total 1	Mg 1	0	0

- Molecule 5 is (5S)-7-benzyl-1,3,7-triazaspiro[4.4]nonane-2,4-dione (CCD ID: A1BCA) (formula: C₁₃H₁₅N₃O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	Total 18	C 13	N 3	O 2	0	0
5	D	1	Total 18	C 13	N 3	O 2	0	0

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	E	1	Total C O 4 2 2	0	0
6	E	1	Total C O 4 2 2	0	0
6	E	1	Total C O 4 2 2	0	0
6	E	1	Total C O 4 2 2	0	0

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Na 1 1	0	0
7	E	1	Total Na 1 1	0	0

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total C O 6 3 3	0	0
8	B	1	Total C O 6 3 3	0	0
8	B	1	Total C O 6 3 3	0	0
8	B	1	Total C O 6 3 3	0	0
8	B	1	Total C O 6 3 3	0	0
8	B	1	Total C O 6 3 3	0	0
8	B	1	Total C O 6 3 3	0	0
8	E	1	Total C O 6 3 3	0	0
8	E	1	Total C O 6 3 3	0	0
8	E	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
8	F	1	6	3	3	0	0

- Molecule 9 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
9	B	1	4	2	1	1	0	0
9	B	1	4	2	1	1	0	0
9	D	1	4	2	1	1	0	0
9	E	1	4	2	1	1	0	0

- Molecule 10 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	C	O	0	0
			10	6	4		
10	D	1	Total	C	O	0	0
			10	6	4		

- Molecule 11 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
11	B	1	15	8	2	4	1	0	0


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	78	Total 78	O 78	0	0
12	B	433	Total 433	O 433	0	0
12	C	86	Total 86	O 86	0	0
12	D	71	Total 71	O 71	0	0
12	E	262	Total 262	O 262	0	0
12	F	34	Total 34	O 34	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phenylalanine-tRNA ligase alpha subunit



- Molecule 1: Phenylalanine-tRNA ligase alpha subunit



- Molecule 2: Phenylalanine-tRNA ligase beta subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	146.69Å 64.06Å 187.92Å 90.00° 110.86° 90.00°	Depositor
Resolution (Å)	48.64 – 2.45 48.64 – 2.45	Depositor EDS
% Data completeness (in resolution range)	90.0 (48.64-2.45) 90.1 (48.64-2.45)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 2.42Å)	Xtrriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.203 , 0.249 0.204 , 0.249	Depositor DCC
R_{free} test set	6125 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	34.0	Xtrriage
Anisotropy	0.089	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.025 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	21921	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, DMS, NA, A1BCA, MG, PGE, GOL, EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.07	0/2594	0.23	0/3538
1	D	0.07	0/2721	0.21	0/3704
2	B	0.07	0/6466	0.22	0/8852
2	E	0.08	0/6213	0.22	0/8508
3	C	0.06	0/1746	0.13	0/2720
3	F	0.06	0/1746	0.14	0/2720
All	All	0.07	0/21486	0.21	0/30042

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2535	0	2449	37	0
1	D	2662	0	2608	27	0
2	B	6329	0	6354	46	0
2	E	6083	0	6096	53	0
3	C	1563	0	792	3	0
3	F	1563	0	792	9	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	2	0	0	0	0
4	D	1	0	0	0	0
5	A	18	0	0	1	0
5	D	18	0	0	0	0
6	A	4	0	3	0	0
6	B	28	0	21	1	0
6	E	16	0	12	0	0
7	A	1	0	0	0	0
7	E	1	0	0	0	0
8	B	42	0	56	2	0
8	E	18	0	24	2	0
8	F	6	0	8	0	0
9	B	8	0	12	2	0
9	D	4	0	6	0	0
9	E	4	0	6	1	0
10	B	10	0	14	1	0
10	D	10	0	14	0	0
11	B	30	0	34	1	0
12	A	78	0	0	1	0
12	B	433	0	0	2	0
12	C	86	0	0	0	0
12	D	71	0	0	1	0
12	E	262	0	0	0	0
12	F	34	0	0	0	0
All	All	21921	0	19301	153	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 153 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:20:U:H5''	1:D:46:GLN:HG2	1.69	0.74
1:A:341:ALA:HA	2:B:512:ARG:HG2	1.76	0.68
3:F:69:U:H2'	3:F:70:G:C8	2.29	0.67
1:A:180:GLN:HB3	1:A:219:LEU:HD22	1.78	0.65
1:D:231:ARG:HE	1:D:247:ARG:HH12	1.44	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/342 (97%)	312 (94%)	16 (5%)	3 (1%)	14	18
1	D	342/342 (100%)	332 (97%)	9 (3%)	1 (0%)	37	45
2	B	844/835 (101%)	823 (98%)	19 (2%)	2 (0%)	44	54
2	E	806/835 (96%)	772 (96%)	32 (4%)	2 (0%)	44	54
All	All	2323/2354 (99%)	2239 (96%)	76 (3%)	8 (0%)	37	45

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	48	LEU
2	B	373	THR
2	E	373[A]	THR
2	E	373[B]	THR
1	A	47	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/268 (94%)	247 (98%)	5 (2%)	50	65
1	D	269/268 (100%)	266 (99%)	3 (1%)	70	81
2	B	661/652 (101%)	658 (100%)	3 (0%)	86	92
2	E	632/652 (97%)	626 (99%)	6 (1%)	75	86
All	All	1814/1840 (99%)	1797 (99%)	17 (1%)	73	86

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	452	HIS
2	E	607	VAL
2	B	564	THR
1	D	104	VAL
1	D	175	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	628	GLN
2	B	687	GLN
2	E	132	ASN
1	D	215	GLN
2	B	35	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	C	72/77 (93%)	7 (9%)	0
3	F	72/77 (93%)	11 (15%)	0
All	All	144/154 (93%)	18 (12%)	0

5 of 18 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	C	6	G
3	C	16	U
3	C	21	A
3	C	22	U
3	C	23	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 39 ligands modelled in this entry, 6 are monoatomic - leaving 33 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	ACT	B	911	-	3,3,3	1.34	0	3,3,3	1.36	0
8	GOL	B	919	-	5,5,5	0.94	0	5,5,5	1.05	0
10	PGE	B	906	-	9,9,9	0.31	0	8,8,8	0.30	0
6	ACT	B	913	-	3,3,3	1.39	0	3,3,3	1.36	0
6	ACT	E	905	-	3,3,3	1.37	0	3,3,3	1.36	0
8	GOL	E	907	-	5,5,5	0.94	0	5,5,5	1.08	0
11	EPE	B	910	-	15,15,15	0.82	1 (6%)	19,20,20	1.69	3 (15%)
6	ACT	B	915	-	3,3,3	1.41	1 (33%)	3,3,3	1.37	0
9	DMS	B	908	-	3,3,3	0.67	0	3,3,3	0.51	0
9	DMS	E	908	-	3,3,3	0.67	0	3,3,3	0.59	0
10	PGE	D	402	-	9,9,9	0.31	0	8,8,8	0.29	0
8	GOL	B	901	-	5,5,5	0.92	0	5,5,5	1.09	0
8	GOL	B	902	-	5,5,5	0.94	0	5,5,5	1.06	0
8	GOL	E	901	-	5,5,5	0.93	0	5,5,5	1.03	0
6	ACT	B	916	-	3,3,3	1.36	0	3,3,3	1.37	0
8	GOL	B	907	-	5,5,5	0.93	0	5,5,5	1.06	0
9	DMS	D	401	-	3,3,3	0.67	0	3,3,3	0.52	0
6	ACT	B	917	-	3,3,3	1.40	1 (33%)	3,3,3	1.37	0
6	ACT	E	909	-	3,3,3	1.40	1 (33%)	3,3,3	1.37	0
8	GOL	B	904	-	5,5,5	0.93	0	5,5,5	1.07	0
5	A1BCA	A	402	-	20,20,20	7.18	12 (60%)	24,29,29	2.44	5 (20%)
6	ACT	A	403	-	3,3,3	1.36	0	3,3,3	1.37	0
6	ACT	B	912	-	3,3,3	1.37	0	3,3,3	1.36	0
6	ACT	E	903	-	3,3,3	1.39	0	3,3,3	1.35	0
8	GOL	E	902	-	5,5,5	0.95	0	5,5,5	1.08	0
8	GOL	B	903	-	5,5,5	0.95	0	5,5,5	1.05	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	GOL	B	914	-	5,5,5	0.95	0	5,5,5	1.09	0
9	DMS	B	905	-	3,3,3	0.67	0	3,3,3	0.53	0
6	ACT	E	904	-	3,3,3	1.37	0	3,3,3	1.37	0
8	GOL	F	101	-	5,5,5	0.94	0	5,5,5	1.06	0
6	ACT	B	918	-	3,3,3	1.35	0	3,3,3	1.38	0
11	EPE	B	909	-	15,15,15	0.79	1 (6%)	19,20,20	1.77	3 (15%)
5	A1BCA	D	403	-	20,20,20	7.23	12 (60%)	24,29,29	2.51	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	B	914	-	-	0/4/4/4	-
8	GOL	B	904	-	-	0/4/4/4	-
5	A1BCA	A	402	-	-	0/4/30/30	0/3/3/3
10	PGE	D	402	-	-	4/7/7/7	-
11	EPE	B	910	-	-	4/9/19/19	0/1/1/1
8	GOL	F	101	-	-	0/4/4/4	-
8	GOL	B	901	-	-	2/4/4/4	-
8	GOL	B	919	-	-	0/4/4/4	-
8	GOL	B	902	-	-	0/4/4/4	-
10	PGE	B	906	-	-	3/7/7/7	-
11	EPE	B	909	-	-	2/9/19/19	0/1/1/1
8	GOL	E	901	-	-	3/4/4/4	-
5	A1BCA	D	403	-	-	0/4/30/30	0/3/3/3
8	GOL	B	907	-	-	0/4/4/4	-
8	GOL	E	907	-	-	0/4/4/4	-
8	GOL	E	902	-	-	1/4/4/4	-
8	GOL	B	903	-	-	0/4/4/4	-

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	403	A1BCA	C18-C07	-17.84	1.31	1.53
5	A	402	A1BCA	C18-C07	-17.64	1.31	1.53
5	D	403	A1BCA	C11-N10	-15.81	1.17	1.47
5	A	402	A1BCA	C11-N10	-15.74	1.17	1.47
5	D	403	A1BCA	C18-N10	14.92	1.60	1.46

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	403	A1BCA	C05-C07-N01	9.51	108.13	100.35
5	A	402	A1BCA	C05-C07-N01	9.36	108.00	100.35
11	B	909	EPE	C5-N4-C3	5.21	120.07	108.84
5	D	403	A1BCA	C07-N01-C02	-5.02	107.95	112.90
5	A	402	A1BCA	C07-N01-C02	-4.91	108.05	112.90

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	B	901	GOL	O1-C1-C2-O2
8	B	901	GOL	O1-C1-C2-C3
11	B	910	EPE	C8-C7-N4-C5
8	E	901	GOL	O1-C1-C2-C3
8	E	901	GOL	O1-C1-C2-O2


There are no ring outliers.

9 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	911	ACT	1	0
10	B	906	PGE	1	0
9	B	908	DMS	2	0
9	E	908	DMS	1	0
8	E	901	GOL	2	0
5	A	402	A1BCA	1	0
8	B	903	GOL	1	0
8	B	914	GOL	1	0
11	B	909	EPE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	335/342 (97%)	0.72	50 (14%) 7 6	19, 47, 122, 214	0
1	D	338/342 (98%)	0.55	32 (9%) 15 15	16, 47, 90, 115	8 (2%)
2	B	834/835 (99%)	-0.12	11 (1%) 74 76	10, 30, 65, 105	12 (1%)
2	E	806/835 (96%)	0.53	110 (13%) 8 7	12, 47, 118, 170	10 (1%)
3	C	73/77 (94%)	0.07	8 (10%) 12 11	25, 50, 176, 226	0
3	F	73/77 (94%)	1.18	23 (31%) 1 1	35, 87, 188, 230	0
All	All	2459/2508 (98%)	0.34	234 (9%) 15 15	10, 39, 112, 230	30 (1%)

The worst 5 of 234 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	90[A]	LEU	8.8
2	E	364[A]	SER	8.3
1	D	91[A]	VAL	8.3
2	E	203[A]	VAL	6.9
1	D	0	ALA	6.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	GOL	E	901	6/6	0.67	0.22	55,58,62,65	0
10	PGE	B	906	10/10	0.74	0.17	31,44,52,58	0
6	ACT	B	915	4/4	0.75	0.18	35,36,41,49	0
7	NA	E	906	1/1	0.75	0.20	43,43,43,43	0
8	GOL	B	903	6/6	0.76	0.20	54,59,61,63	0
5	A1BCA	D	403	18/18	0.78	0.17	31,61,93,95	0
10	PGE	D	402	10/10	0.78	0.17	41,46,54,56	0
6	ACT	A	403	4/4	0.79	0.20	51,52,57,59	0
5	A1BCA	A	402	18/18	0.80	0.16	33,52,75,82	0
6	ACT	E	903	4/4	0.81	0.18	49,53,55,56	0
9	DMS	E	908	4/4	0.81	0.22	26,33,61,64	0
11	EPE	B	909	15/15	0.81	0.17	49,58,71,76	0
8	GOL	E	907	6/6	0.82	0.16	61,65,75,78	0
9	DMS	D	401	4/4	0.82	0.23	46,61,62,74	0
6	ACT	E	905	4/4	0.82	0.22	55,62,65,66	0
8	GOL	B	914	6/6	0.83	0.12	44,52,57,62	0
8	GOL	B	902	6/6	0.84	0.16	45,55,65,70	0
6	ACT	B	918	4/4	0.84	0.13	27,31,40,47	0
6	ACT	B	916	4/4	0.85	0.14	38,48,50,50	0
8	GOL	B	907	6/6	0.85	0.14	44,46,48,51	0
6	ACT	B	917	4/4	0.86	0.23	33,35,43,46	0
6	ACT	B	913	4/4	0.86	0.13	43,46,48,49	0
8	GOL	B	904	6/6	0.87	0.12	47,53,58,59	0
11	EPE	B	910	15/15	0.87	0.15	44,59,70,71	0
6	ACT	E	909	4/4	0.88	0.12	51,53,55,57	0
8	GOL	E	902	6/6	0.88	0.11	40,41,45,48	0
8	GOL	B	919	6/6	0.88	0.27	30,40,46,53	0
6	ACT	B	912	4/4	0.89	0.11	35,44,50,51	0
8	GOL	F	101	6/6	0.89	0.19	44,47,50,52	0
9	DMS	B	908	4/4	0.91	0.12	31,45,51,54	0
9	DMS	B	905	4/4	0.92	0.19	34,42,47,58	0
8	GOL	B	901	6/6	0.92	0.10	25,30,37,37	0
6	ACT	B	911	4/4	0.93	0.11	30,31,32,35	0
6	ACT	E	904	4/4	0.93	0.09	35,41,42,45	0
7	NA	A	404	1/1	0.96	0.05	33,33,33,33	0
4	MG	C	101	1/1	0.97	0.07	35,35,35,35	0
4	MG	D	404	1/1	0.99	0.02	34,34,34,34	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	MG	A	401	1/1	0.99	0.02	26,26,26,26	0
4	MG	C	102	1/1	0.99	0.03	37,37,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.