wwPDB X-ray Structure Validation Summary Report (i)

Oct 16, 2023 – 04:51 PM EDT

PDB ID	:	2DU8
Title	:	Crystal structure of human D-amino acid oxidase
Authors	:	Kawazoe, T.; Tsuge, H.; Fukui, K.
Deposited on	:	2006-07-20
Resolution	:	2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	4% 73% 22% • •
1	B	347	2% 73% 22% • •
1	G	347	4% 78% 17% • •
1	J	347	7% 71% 23% • •

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 11436 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-amino-acid oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	340	2733	1751	479	494	9	0	0	0
1	B	340	2733	1751	479	494	9	0	0	0
1	G	340	2733	1751	479	494	9	0	0	0
1	J	340	2733	1751	479	494	9	0	0	0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).

FAD



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0
2	B	1	53	27	9	15	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	G	1	53	27	9	15	2	0	0
2	J	1	53	27	9	15	2	0	0

- Molecule 3 is BENZOIC ACID (three-letter code: BEZ) (formula: $C_7H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	9	7	2	0	0
3	B	1	9	7	2	0	0
3	G	1	9	7	2	0	0
3	J	1	9	7	2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	71	71	71	0	0
4	B	83	83	83	0	0
4	G	62	62	62	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	J	40	40	40	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: D-amino-acid oxidase

Chain A:



- Molecule 1: D-amino-acid oxidase

Chain B:



- Molecule 1: D-amino-acid oxidase

Chain G:



- Molecule 1: D-amino-acid oxidase

Chain J:







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α, β, γ	150.98Å 183.18Å 51.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.78 – 2.50 46.78 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.1 (46.78-2.50) 99.1 (46.78-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	4.58 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.223 , 0.270 0.223 , 0.219	Depositor DCC
R_{free} test set	2469 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å^2)	49.4	Xtriage
Anisotropy	0.403	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 38.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	11436	wwPDB-VP
Average B, all atoms (Å^2)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.05 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.1757e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, BEZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.57	1/2810 (0.0%)	0.66	1/3824 (0.0%)
1	B	0.60	1/2810 (0.0%)	0.68	2/3824 (0.1%)
1	G	0.58	0/2810	0.67	1/3824 (0.0%)
1	J	0.56	1/2810 (0.0%)	0.63	0/3824
All	All	0.58	3/11240 (0.0%)	0.66	4/15296 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1018	CYS	CB-SG	-6.83	1.70	1.82
1	J	3018	CYS	CB-SG	-5.85	1.72	1.81
1	A	18	CYS	CB-SG	-5.47	1.73	1.81

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	40	THR	N-CA-C	5.23	125.11	111.00
1	G	2040	THR	N-CA-C	5.22	125.10	111.00
1	B	1040	THR	N-CA-C	5.19	125.02	111.00
1	B	1316	LEU	CA-CB-CG	5.04	126.89	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2733	0	2680	64	0
1	B	2733	0	2677	63	0
1	G	2733	0	2677	50	0
1	J	2733	0	2677	69	0
2	A	53	0	31	1	0
2	B	53	0	31	0	0
2	G	53	0	31	0	0
2	J	53	0	31	1	0
3	A	9	0	5	0	0
3	B	9	0	5	0	0
3	G	9	0	5	1	0
3	J	9	0	5	0	0
4	A	71	0	0	4	0
4	B	83	0	0	2	0
4	G	62	0	0	4	0
4	J	40	0	0	1	0
All	All	11436	0	10855	242	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 242 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:ASP:HB3	4:A:403:HOH:O	1.26	1.28
1:J:3180:ASN:HD22	1:J:3307:HIS:HD2	1.09	0.96
1:B:1028:GLN:HB3	1:B:1029:PRO:HD3	1.53	0.91
1:J:3241:ILE:HD12	1:J:3259:ILE:HD11	1.51	0.89
1:G:2241:ILE:HD13	1:G:2255:ASP:HB3	1.56	0.88

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	338/347 (97%)	310 (92%)	26 (8%)	2 (1%)	25	43
1	B	338/347 (97%)	320 (95%)	17 (5%)	1 (0%)	41	61
1	G	338/347 (97%)	319 (94%)	16 (5%)	3 (1%)	17	31
1	J	338/347 (97%)	315 (93%)	21 (6%)	2 (1%)	25	43
All	All	1352/1388 (97%)	1264 (94%)	80 (6%)	8 (1%)	25	43

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	40	THR
1	B	1040	THR
1	G	2040	THR
1	J	3040	THR
1	G	2053	GLN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/299 (98%)	270 (92%)	22 (8%)	13	26
1	B	292/299 (98%)	269 (92%)	23 (8%)	12	24
1	G	292/299 (98%)	272 (93%)	20 (7%)	16	30
1	J	292/299 (98%)	270 (92%)	22 (8%)	13	26
All	All	1168/1196 (98%)	1081 (93%)	87 (7%)	13	27

5 of 87 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	2221	ARG
1	J	3152	LEU
1	G	2270	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	J	3056	LEU
1	J	3205	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 45 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	2196	GLN
1	J	3063	GLN
1	G	2201	GLN
1	G	2307	HIS
1	J	3143	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	BEZ	B	1352	-	9,9,9	0.91	0	11,11,11	0.61	0
3	BEZ	A	352	-	9,9,9	0.88	0	11,11,11	0.56	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	BEZ	J	3352	-	9,9,9	0.87	0	11,11,11	0.63	0
3	BEZ	G	2352	-	9,9,9	0.82	0	11,11,11	0.89	0
2	FAD	A	351	-	53,58,58	0.92	2 (3%)	68,89,89	1.50	15 (22%)
2	FAD	G	2351	-	53,58,58	0.97	5 (9%)	68,89,89	1.46	11 (16%)
2	FAD	B	1351	-	53,58,58	1.08	3 (5%)	68,89,89	1.49	13 (19%)
2	FAD	J	3351	-	53,58,58	0.92	1 (1%)	68,89,89	1.44	14 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BEZ	B	1352	-	-	0/4/4/4	0/1/1/1
3	BEZ	A	352	-	-	0/4/4/4	0/1/1/1
3	BEZ	J	3352	-	-	0/4/4/4	0/1/1/1
3	BEZ	G	2352	-	-	0/4/4/4	0/1/1/1
2	FAD	A	351	-	-	1/30/50/50	0/6/6/6
2	FAD	G	2351	-	-	1/30/50/50	0/6/6/6
2	FAD	B	1351	-	-	1/30/50/50	0/6/6/6
2	FAD	J	3351	-	-	1/30/50/50	0/6/6/6

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1351	FAD	O4B-C1B	3.40	1.45	1.41
2	A	351	FAD	C4X-N5	3.17	1.36	1.30
2	B	1351	FAD	C4X-N5	3.03	1.36	1.30
2	J	3351	FAD	C4X-N5	3.01	1.36	1.30
2	G	2351	FAD	C4X-N5	2.51	1.35	1.30

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	1351	FAD	N3A-C2A-N1A	-5.43	120.20	128.68
2	G	2351	FAD	N3A-C2A-N1A	-5.30	120.40	128.68
2	A	351	FAD	N3A-C2A-N1A	-4.68	121.36	128.68
2	J	3351	FAD	N3A-C2A-N1A	-4.59	121.51	128.68
2	G	2351	FAD	C4X-C10-N10	3.79	122.02	116.48

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	G	2351	FAD	O4B-C4B-C5B-O5B
2	A	351	FAD	O4B-C4B-C5B-O5B
2	B	1351	FAD	O4B-C4B-C5B-O5B
2	J	3351	FAD	O4B-C4B-C5B-O5B

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	2352	BEZ	1	0
2	A	351	FAD	1	0
2	J	3351	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand FAD A 351



Bond lengths



Bond angles



Torsions



Rings

Ligand FAD G 2351



Bond lengths



Bond angles



Torsions



Rings

Ligand FAD B 1351



Bond lengths



Bond angles



Torsions



Rings



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	340/347 (97%)	0.18	13 (3%)	40	43	33, 49, 72, 82	0
1	B	340/347 (97%)	0.13	8 (2%)	59	62	32, 47, 72, 89	0
1	G	340/347 (97%)	0.23	14 (4%)	37	40	30, 48, 70, 87	0
1	J	340/347 (97%)	0.44	23 (6%)	17	17	35, 54, 79, 98	0
All	All	1360/1388 (97%)	0.24	58 (4%)	35	38	30, 50, 74, 98	0

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	2339	LEU	11.3
1	J	3339	LEU	10.1
1	J	3340	SER	9.3
1	J	3300	PRO	9.1
1	J	3301	SER	8.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	BEZ	G	2352	9/9	0.85	0.27	52,57,60,61	0
3	BEZ	B	1352	9/9	0.91	0.22	51,57,59,59	0
3	BEZ	J	3352	9/9	0.93	0.20	62,64,67,67	0
3	BEZ	A	352	9/9	0.95	0.30	60,61,62,62	0
2	FAD	A	351	53/53	0.96	0.14	35,39,43,43	0
2	FAD	G	2351	53/53	0.97	0.15	32,37,40,42	0
2	FAD	B	1351	53/53	0.98	0.13	27,33,37,40	0
2	FAD	J	3351	53/53	0.98	0.13	42,45,47,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 351:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around FAD G 2351:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FAD B 1351:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FAD J 3351:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.