wwPDB X-ray Structure Validation Summary Report (i)

May 15, 2020 – 10:47 am BST

PDB ID	:	4E4L
Title	:	JAK1 kinase (JH1 domain) in complex with compound 30
Authors	:	Eigenbrot, C.
Deposited on	:	2012-03-13
Resolution	:	2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	% 84% 8% 8%
1	B	302	2% 84% 8% 7%
1	D	302	2% 82% 12% 6%
1	E	302	% 83% 10% 7%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 9949 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase JAK1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	279	2301	1466	393	425	2	15	0	3	0
1	B	281	2337	1488	399	433	2	15	0	5	0
1	E	282	2333	1484	399	433	2	15	0	4	0
1	D	283	2334	1483	400	434	2	15	0	4	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	853	GLY	-	EXPRESSION TAG	UNP P23458
B	853	GLY	-	EXPRESSION TAG	UNP P23458
E	853	GLY	-	EXPRESSION TAG	UNP P23458
D	853	GLY	-	EXPRESSION TAG	UNP P23458

- Molecule 2 is 1-[4-methyl-1-(methylsulfonyl)piperidin-4-yl]-1,6-dihydroimidazo[4,5-d]pyrrolo[2,3-b]pyridine (three-letter code: 0NH) (formula: $C_{15}H_{19}N_5O_2S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	23	15	5	2	1	0	0
			Total	C	N	O	S		
2	A	1	23	15	5	2	1	0	0
			Total	C	N	O	S		
2	B	1	23	15	5	2	1	0	0
			Total	C	N	O	S		
2	D	1	23	15	5	2	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	143	143	143	0	0
			Total	O		
3	B	140	140	140	0	0
			Total	O		
3	E	132	132	132	0	0
			Total	O		
3	D	137	137	137	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase JAK1

Chain A:



- Molecule 1: Tyrosine-protein kinase JAK1

Chain B:



- Molecule 1: Tyrosine-protein kinase JAK1

Chain E:



- Molecule 1: Tyrosine-protein kinase JAK1

Chain D:


GLY
ASP
ILE
VAL
SER
GLU
LYS
LYS
PRO
ALA
THR
E864
H885
F886
L910
K911
PRO
GLU
SER
GLY
GLY
ASN
H918
I919
H934
E935
N936
T945
GLU
ASP
G948
G949
N950
Y967
L977
Q980
R1002
D1003
R1007
L1010
V1011
L1024
T1030
D1031
Y1035
T1036
V1037
L1053
S1056
K1057
F1058
W1065
L1075
L1089
K1090
P1094
T1102
T1107
K1112
P1118
M1128
W1132
S1137
N1144
K1154

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	42.84Å 171.76Å 87.82Å 90.00° 92.15° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 47.95 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.5 (50.00-2.00) 94.6 (47.95-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$< I/\sigma(I) >$ [1]	2.67 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.190 , 0.228 0.194 , 0.232	Depositor DCC
R_{free} test set	3725 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å^2)	26.1	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 51.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.107 for h,-k,-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	9949	wwPDB-VP
Average B, all atoms (Å^2)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 53.86 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.9670e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 0NH, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.51	0/2315	0.62	0/3112
1	B	0.50	0/2352	0.60	0/3162
1	D	0.49	0/2348	0.61	0/3156
1	E	0.49	0/2348	0.61	0/3158
All	All	0.50	0/9363	0.61	0/12588

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2301	0	2301	17	0
1	B	2337	0	2326	22	0
1	D	2334	0	2324	28	0
1	E	2333	0	2325	29	0
2	A	46	0	38	1	0
2	B	23	0	19	1	0
2	D	23	0	19	1	0
3	A	143	0	0	1	0
3	B	140	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	137	0	0	5	0
3	E	132	0	0	6	0
All	All	9949	0	9352	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 90 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:900:ASN:HB2	3:E:1432:HOH:O	1.66	0.94
1:A:1053:LEU:O	1:B:1102:THR:HG21	1.75	0.87
1:B:1000:VAL:HB	1:B:1058[B]:PHE:CE2	2.19	0.78
1:B:1000:VAL:HB	1:B:1058[B]:PHE:CD2	2.20	0.77
1:E:885:HIS:HD2	1:E:886:PHE:CZ	2.07	0.72

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	274/302 (91%)	271 (99%)	3 (1%)	0	100 100
1	B	278/302 (92%)	276 (99%)	2 (1%)	0	100 100
1	D	279/302 (92%)	275 (99%)	3 (1%)	1 (0%)	34 30
1	E	278/302 (92%)	274 (99%)	4 (1%)	0	100 100
All	All	1109/1208 (92%)	1096 (99%)	12 (1%)	1 (0%)	51 49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	949	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/267 (95%)	251 (99%)	2 (1%)	81	86
1	B	257/267 (96%)	256 (100%)	1 (0%)	91	93
1	D	256/267 (96%)	253 (99%)	3 (1%)	71	76
1	E	257/267 (96%)	256 (100%)	1 (0%)	91	93
All	All	1023/1068 (96%)	1016 (99%)	7 (1%)	84	88

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	1056	SER
1	D	1137	SER
1	D	885	HIS
1	A	1102	THR
1	D	950	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	936	ASN
1	D	936	ASN
1	E	900	ASN
1	B	885	HIS
1	E	885	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	PTR	D	1035	1	15,16,17	0.87	1 (6%)	19,22,24	0.79	0
1	PTR	D	1034	1	15,16,17	0.75	0	19,22,24	0.86	0
1	PTR	A	1034	1	15,16,17	0.71	0	19,22,24	0.96	1 (5%)
1	PTR	B	1034	1	15,16,17	0.77	0	19,22,24	0.89	1 (5%)
1	PTR	A	1035	1	15,16,17	0.91	1 (6%)	19,22,24	0.76	0
1	PTR	E	1034	1	15,16,17	0.70	0	19,22,24	0.92	1 (5%)
1	PTR	E	1035	1	15,16,17	0.96	1 (6%)	19,22,24	0.86	0
1	PTR	B	1035	1	15,16,17	0.87	0	19,22,24	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	D	1035	1	-	0/10/11/13	0/1/1/1
1	PTR	D	1034	1	-	0/10/11/13	0/1/1/1
1	PTR	A	1034	1	-	0/10/11/13	0/1/1/1
1	PTR	B	1034	1	-	0/10/11/13	0/1/1/1
1	PTR	A	1035	1	-	0/10/11/13	0/1/1/1
1	PTR	E	1034	1	-	0/10/11/13	0/1/1/1
1	PTR	E	1035	1	-	0/10/11/13	0/1/1/1
1	PTR	B	1035	1	-	0/10/11/13	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	1035	PTR	P-OH	2.60	1.63	1.59
1	A	1035	PTR	P-OH	2.53	1.63	1.59
1	D	1035	PTR	P-OH	2.44	1.63	1.59

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	E	1034	PTR	OH-P-O1P	-2.28	100.70	109.31
1	A	1034	PTR	OH-P-O1P	-2.23	100.89	109.31
1	B	1034	PTR	OH-P-O1P	-2.13	101.29	109.31

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	1035	PTR	1	0
1	B	1035	PTR	1	0

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	0NH	D	1201	-	20,26,26	5.21	14 (70%)	22,41,41	4.35	16 (72%)
2	0NH	A	1201	-	20,26,26	5.38	11 (55%)	22,41,41	3.14	10 (45%)
2	0NH	B	1201	-	20,26,26	5.22	12 (60%)	22,41,41	3.15	11 (50%)
2	0NH	A	1202	-	20,26,26	5.35	11 (55%)	22,41,41	3.09	11 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0NH	D	1201	-	-	5/10/24/24	0/4/4/4
2	0NH	A	1201	-	-	5/10/24/24	0/4/4/4
2	0NH	B	1201	-	-	5/10/24/24	0/4/4/4
2	0NH	A	1202	-	-	5/10/24/24	0/4/4/4

The worst 5 of 48 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1202	0NH	C24-C15	-12.96	1.39	1.52
2	A	1201	0NH	C18-C19	-12.63	1.33	1.53
2	A	1201	0NH	C24-C15	-12.37	1.40	1.52
2	B	1201	0NH	C18-C19	-11.86	1.34	1.53
2	B	1201	0NH	C24-C15	-11.73	1.41	1.52

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	D	1201	0NH	C6-N5-C4	9.39	122.56	112.17
2	B	1201	0NH	C6-N5-C4	9.29	122.45	112.17
2	A	1201	0NH	C6-N5-C4	9.24	122.39	112.17
2	A	1202	0NH	C6-N5-C4	7.76	120.76	112.17
2	D	1201	0NH	C1-C2-C3	7.68	124.50	110.87

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	1201	0NH	C3-C2-N12-C13
2	D	1201	0NH	C7-C2-N12-C13
2	D	1201	0NH	C6-N5-S8-O9
2	D	1201	0NH	C6-N5-S8-O10
2	D	1201	0NH	C6-N5-S8-C11

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1201	0NH	1	0
2	A	1201	0NH	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1201	0NH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 0NH D 1201



Ligand 0NH A 1201



Bond lengths



Bond angles



Torsions



Rings

Ligand 0NH B 1201



Bond lengths



Bond angles



Torsions



Rings

Ligand 0NH A 1202



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	277/302 (91%)	-0.26	2 (0%)	87	87	12, 28, 49, 71	6 (2%)
1	B	279/302 (92%)	-0.32	5 (1%)	68	66	12, 27, 51, 76	9 (3%)
1	D	281/302 (93%)	-0.19	6 (2%)	63	62	14, 31, 56, 78	9 (3%)
1	E	280/302 (92%)	-0.22	4 (1%)	75	74	12, 30, 54, 79	8 (2%)
All	All	1117/1208 (92%)	-0.25	17 (1%)	73	72	12, 29, 53, 79	32 (2%)

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	885	HIS	4.0
1	E	1031	ASP	3.7
1	D	885	HIS	3.2
1	E	885	HIS	3.1
1	D	1154	LYS	3.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	PTR	B	1034	16/17	0.89	0.12	30,43,58,60	0
1	PTR	E	1035	16/17	0.89	0.11	34,38,49,54	0
1	PTR	D	1035	16/17	0.92	0.11	34,40,51,53	0
1	PTR	E	1034	16/17	0.92	0.14	38,51,67,72	0
1	PTR	D	1034	16/17	0.92	0.12	35,54,71,75	0
1	PTR	A	1034	16/17	0.93	0.10	31,43,61,61	0
1	PTR	A	1035	16/17	0.94	0.10	25,32,42,45	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	PTR	B	1035	16/17	0.95	0.09	28,35,44,44	0

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	0NH	D	1201	23/23	0.98	0.09	7,15,19,23	0
2	0NH	A	1201	23/23	0.98	0.10	7,12,17,18	0
2	0NH	B	1201	23/23	0.98	0.08	10,15,19,24	0
2	0NH	A	1202	23/23	0.98	0.09	8,12,16,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 0NH D 1201:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around 0NH A 1201:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around 0NH B 1201:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around 0NH A 1202:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.