Full wwPDB X-ray Structure Validation Report (i)

Jun 13, 2024 – 07:17 AM EDT

PDB ID	:	4G21
Title	:	Structural basis for the accommodation of bis- and tris-aromatic derivatives in Vitamin D Nuclear Receptor
Authors	:	Ciesielski, F.; Sato, Y.; Moras, D.; Rochel, N.
Deposited on	:	2012-07-11
Resolution	:	2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	% 49% 28% • 20%
2	B	15	7% 33% 27% 13% 27%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2017 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vitamin D3 receptor A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	239	1858	1183	315	348	12	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	154	HIS	-	expression tag	UNP Q9PTN2
A	155	MET	-	expression tag	UNP Q9PTN2

- Molecule 2 is a protein called Nuclear receptor coactivator 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	11	101	64	23	14	0	0	0

- Molecule 3 is 3-(5'-{[3,4-bis(hydroxymethyl)benzyl]oxy}-2'-ethyl-2-propylbiphenyl-4-yl)pentan-3-ol (three-letter code: 0VP) (formula: $C_{31}H_{40}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	34	30	4	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	24	24	24	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Vitamin D3 receptor A

Chain A:



• Molecule 2: Nuclear receptor coactivator 1

Chain B:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α, β, γ	66.29Å 66.29Å 263.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.90 19.46 – 2.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.90) 93.8 (19.46-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.12 (at 2.88Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.232 , 0.291 0.221 , 0.257	Depositor DCC
R_{free} test set	445 reflections (5.41%)	wwPDB-VP
Wilson B-factor (Å^2)	62.1	Xtriage
Anisotropy	0.577	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 57.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	2017	wwPDB-VP
Average B, all atoms (Å^2)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.70% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 0VP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.30	0/1895	0.52	0/2568
2	B	0.32	0/102	0.53	0/134
All	All	0.30	0/1997	0.52	0/2702

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1858	0	1805	81	0
2	B	101	0	110	14	0
3	A	34	0	35	3	0
4	A	24	0	0	4	0
All	All	2017	0	1950	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:686:ARG:HG3	2:B:687:HIS:H	1.37	0.90
1:A:155:MET:HA	1:A:397:ILE:HG12	1.60	0.84
1:A:339:GLU:HB3	1:A:340:PRO:HD3	1.60	0.83
1:A:369:PRO:O	1:A:374:VAL:HG21	1.84	0.78
2:B:692:ARG:HE	2:B:695:GLN:NE2	1.90	0.70
1:A:328:VAL:HG13	3:A:501:0VP:H23	1.74	0.69
1:A:338:LEU:O	1:A:342:VAL:HG23	1.93	0.69
1:A:383:LEU:O	1:A:387:LEU:HD23	1.92	0.69
1:A:309:LEU:HD21	1:A:349:LYS:HG2	1.77	0.67
1:A:258:LEU:HD12	1:A:331:ALA:HB2	1.77	0.67
1:A:337:LEU:HB2	1:A:426:GLN:NE2	2.12	0.64
1:A:157:SER:OG	1:A:160:GLN:HG3	1.98	0.64
2:B:686:ARG:HA	2:B:686:ARG:HH21	1.62	0.64
1:A:433:GLN:HB3	1:A:435:GLU:OE2	1.99	0.63
2:B:686:ARG:HG3	2:B:687:HIS:N	2.11	0.61
1:A:157:SER:H	1:A:160:GLN:NE2	1.99	0.61
1:A:337:LEU:O	1:A:340:PRO:HD2	2.01	0.60
1:A:252:LEU:HD22	1:A:332:GLY:HA2	1.85	0.59
1:A:435:GLU:CD	1:A:435:GLU:H	2.06	0.58
1:A:309:LEU:HD22	1:A:309:LEU:N	2.18	0.58
1:A:396:ARG:HH11	1:A:397:ILE:HD11	1.69	0.58
1:A:179:TYR:CD2	1:A:316:CYS:HB3	2.40	0.57
1:A:293:SER:HB3	1:A:371:ARG:HD2	1.87	0.56
1:A:397:ILE:HD12	1:A:397:ILE:N	2.20	0.55
1:A:335:LEU:N	1:A:335:LEU:HD12	2.22	0.55
2:B:692:ARG:O	2:B:695:GLN:HG2	2.06	0.55
1:A:386:ARG:HB2	1:A:386:ARG:NH1	2.22	0.55
1:A:296:ILE:O	1:A:300:MET:HG3	2.07	0.54
1:A:403:ARG:N	1:A:403:ARG:HD2	2.22	0.53
1:A:159:GLU:HA	1:A:162:GLN:NE2	2.24	0.52
1:A:313:SER:HB2	1:A:323:TYR:O	2.10	0.52
1:A:296:ILE:O	1:A:296:ILE:HD13	2.09	0.52
1:A:302:ARG:HG3	3:A:501:0VP:H36	1.91	0.52
1:A:305:GLN:NE2	4:A:619:HOH:O	2.42	0.52
1:A:386:ARG:HB2	1:A:386:ARG:HH11	1.76	0.51
1:A:386:ARG:O	1:A:390:VAL:HG23	2.11	0.51
1:A:309:LEU:HD21	1:A:349:LYS:CG	2.42	0.50
1:A:157:SER:H	1:A:160:GLN:HE21	1.58	0.50
1:A:252:LEU:HA	1:A:330:LYS:O	2.12	0.49
1:A:292:LYS:NZ	2:B:687:HIS:HE1	2.10	0.49
1:A:393:ALA:O	1:A:396:ARG:HB3	2.11	0.49
1:A:266:ILE:O	1:A:270:ILE:HG13	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:LEU:HD22	1:A:332:GLY:CA	2.43	0.49
1:A:441:THR:OG1	1:A:444:VAL:HG23	2.12	0.49
1:A:356:GLU:HA	1:A:359:VAL:HG12	1.94	0.48
1:A:408:LYS:O	1:A:411:GLN:HB3	2.13	0.48
1:A:172:HIS:HE2	1:A:355:GLU:CD	2.17	0.48
1:A:284:ALA:O	1:A:288:ILE:HG13	2.14	0.47
1:A:428:ARG:HG2	1:A:432:PHE:CZ	2.50	0.47
1:A:292:LYS:HD2	2:B:687:HIS:CE1	2.49	0.47
1:A:168:VAL:HG22	1:A:359:VAL:HG11	1.96	0.46
1:A:270:ILE:O	1:A:274:LYS:HG3	2.14	0.46
2:B:686:ARG:HH21	2:B:686:ARG:CA	2.28	0.46
1:A:339:GLU:HB3	1:A:340:PRO:CD	2.39	0.45
1:A:404:LEU:HD12	1:A:404:LEU:N	2.32	0.45
1:A:308:SER:HB3	1:A:311:ASP:OD1	2.17	0.45
1:A:285:GLU:H	1:A:285:GLU:CD	2.21	0.45
1:A:434:PRO:HA	1:A:437:SER:OG	2.16	0.45
1:A:154:HIS:N	4:A:604:HOH:O	2.49	0.44
1:A:389:ASP:O	1:A:390:VAL:C	2.55	0.44
1:A:405:LEU:O	1:A:406:TYR:C	2.55	0.44
1:A:314:TRP:CZ2	1:A:345:GLN:HG3	2.53	0.44
1:A:397:ILE:HD12	1:A:397:ILE:H	1.83	0.44
1:A:310:GLU:OE1	1:A:310:GLU:N	2.48	0.44
2:B:692:ARG:HE	2:B:695:GLN:HE21	1.61	0.44
1:A:258:LEU:HD12	1:A:331:ALA:CB	2.46	0.43
1:A:158:ASP:O	1:A:162:GLN:HG3	2.18	0.43
1:A:329:THR:C	1:A:331:ALA:H	2.22	0.43
1:A:377:HIS:HB2	4:A:605:HOH:O	2.18	0.43
1:A:412:LYS:HA	1:A:415:ASP:OD2	2.19	0.43
2:B:695:GLN:O	2:B:696:GLU:C	2.57	0.43
1:A:157:SER:N	1:A:160:GLN:HE21	2.17	0.43
2:B:686:ARG:CG	2:B:687:HIS:H	2.08	0.43
1:A:335:LEU:HD12	1:A:335:LEU:H	1.82	0.43
1:A:157:SER:CB	1:A:160:GLN:HE21	2.32	0.43
1:A:286:ASP:O	1:A:290:LEU:HG	2.19	0.43
1:A:296:ILE:CG2	1:A:297:GLU:N	2.82	0.42
1:A:397:ILE:H	1:A:397:ILE:CD1	2.32	0.42
1:A:261:LEU:HD21	3:A:501:0VP:C5	2.49	0.42
2:B:688:LYS:HB3	2:B:688:LYS:HE2	1.79	0.42
1:A:254:MET:HA	4:A:623:HOH:O	2.18	0.42
1:A:376:ASP:C	1:A:378:VAL:H	2.22	0.42
1:A:395:ILE:HG23	1:A:405:LEU:HD23	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:342:VAL:O	1:A:346:VAL:HG23	2.20	0.41
1:A:428:ARG:HG2	1:A:432:PHE:CE2	2.54	0.41
2:B:692:ARG:NE	2:B:695:GLN:NE2	2.61	0.41
1:A:397:ILE:N	1:A:397:ILE:CD1	2.82	0.41
1:A:397:ILE:HG22	1:A:398:GLN:OE1	2.21	0.41
1:A:157:SER:N	1:A:160:GLN:NE2	2.67	0.41
1:A:396:ARG:HH11	1:A:397:ILE:CD1	2.34	0.41
1:A:351:LEU:O	1:A:352:LYS:C	2.59	0.40
2:B:696:GLU:CD	2:B:696:GLU:H	2.24	0.40
1:A:395:ILE:HG21	1:A:405:LEU:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/300 (78%)	208 (88%)	24 (10%)	3 (1%)	12	37
2	B	9/15 (60%)	9 (100%)	0	0	100	100
All	All	244/315 (78%)	217 (89%)	24 (10%)	3 (1%)	13	40

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	330	LYS
1	A	328	VAL
1	A	377	HIS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	201/272 (74%)	196 (98%)	5 (2%)	47 78
2	B	11/14 (79%)	9 (82%)	2 (18%)	1 5
All	All	212/286 (74%)	205 (97%)	7 (3%)	38 72

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	158	ASP
1	A	296	ILE
1	A	341	LEU
1	A	392	GLN
1	A	403	ARG
2	B	686	ARG
2	B	692	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	160	GLN
1	A	162	GLN
1	A	305	GLN
1	A	411	GLN
2	B	687	HIS
2	B	695	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	0VP	A	501	-	34,36,37	4.23	22 (64%)	46,50,51	1.82	6 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	0VP	A	501	-	-	2/29/29/30	0/3/3/3

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	0VP	C7-C11	10.49	1.55	1.40
3	A	501	0VP	C13-C16	7.65	1.52	1.39
3	A	501	0VP	C3-C2	6.77	1.53	1.40
3	A	501	0VP	C4-C3	6.37	1.49	1.39
3	A	501	0VP	C18-C19	6.21	1.49	1.40
3	A	501	0VP	C10-C5	6.20	1.65	1.50
3	A	501	0VP	C17-C14	6.10	1.49	1.39
3	A	501	0VP	C17-C18	5.65	1.48	1.39
3	A	501	0VP	C12-C16	5.50	1.47	1.39
3	A	501	0VP	C6-C5	4.92	1.48	1.38
3	A	501	0VP	C4-C5	4.87	1.47	1.39
3	A	501	0VP	C9-C7	4.71	1.47	1.40
3	A	501	0VP	C20-C19	4.10	1.46	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	0VP	C21-C20	3.67	1.44	1.38
3	A	501	0VP	C21-C14	3.66	1.45	1.38
3	A	501	0VP	C24-C16	3.53	1.56	1.53
3	A	501	0VP	C13-C9	3.47	1.44	1.38
3	A	501	0VP	C8-C19	3.05	1.59	1.51
3	A	501	0VP	O1-C10	2.75	1.52	1.43
3	A	501	0VP	C1-C2	2.34	1.43	1.39
3	A	501	0VP	C7-C18	-2.09	1.45	1.49
3	A	501	0VP	C12-C11	2.05	1.42	1.39

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	501	0VP	C10-C5-C6	-9.02	99.85	120.64
3	A	501	0VP	C15-C11-C7	3.22	125.72	122.13
3	A	501	0VP	O1-C10-C5	-3.17	99.85	109.16
3	A	501	0VP	C10-O1-C14	3.14	125.09	117.62
3	A	501	0VP	C7-C18-C19	2.39	125.72	122.28
3	A	501	0VP	C52-C2-C1	-2.04	115.81	120.23

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	0VP	C17-C14-O1-C10
3	A	501	0VP	C21-C14-O1-C10

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	0VP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 0VP A 501



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	239/300 (79%)	-0.27	4 (1%)	70	69	39, 63, 87, 91	0
2	B	11/15 (73%)	0.30	1 (9%)	9	6	58, 69, 88, 93	0
All	All	250/315 (79%)	-0.25	5 (2%)	65	63	39, 63, 87, 93	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	686	ARG	4.0
1	A	155	MET	2.8
1	A	308	SER	2.5
1	A	251	ARG	2.1
1	A	312	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	0VP	A	501	34/35	0.94	0.24	40,53,56,60	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 0VP A 501:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.