wwPDB X-ray Structure Validation Summary Report (i)

Mar 10, 2024 – 10:35 AM EDT

PDB ID	:	4G5O
Title	:	Structure of LGN GL4/Galphai3(Q147L) complex
Authors	:	Jia, M.; Li, J.; Zhu, J.; Wen, W.; Zhang, M.; Wang, W.
Deposited on	:	2012-07-18
Resolution	:	2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	.% 86% 11% • •
1	B	330	2% 85% 10% • 5%
1	C	330	.% 82% 13% • •
1	D	330	2% 86% 12% • •
2	E	26	77% 8% 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	26	50% 31% • 15%
2	G	26	4% 65% 19% 15%
2	H	26	4% 65% 23% 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CIT	A	402	-	X	-	-
4	CIT	C	402	-	X	-	-

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 11190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Guanine nucleotide-binding protein G(k) subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	320	2553	1622	424	491	16	0	1	0
1	B	315	2522	1602	417	487	16	0	1	0
1	C	316	2533	1614	418	485	16	0	0	0
1	D	324	2586	1644	430	496	16	0	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	147	LEU	GLN	engineered mutation	UNP P08754
B	147	LEU	GLN	engineered mutation	UNP P08754
C	147	LEU	GLN	engineered mutation	UNP P08754
D	147	LEU	GLN	engineered mutation	UNP P08754

- Molecule 2 is a protein called G-protein-signaling modulator 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	22	177	113	33	30	1	0	0	0
2	F	22	184	117	34	32	1	0	0	0
2	G	22	188	119	34	34	1	0	0	0
2	H	23	186	119	35	31	1	0	0	0

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	28	10	5	11	2	0	0
			Total	C	N	O	P		
3	B	1	28	10	5	11	2	0	0
			Total	C	N	O	P		
3	C	1	28	10	5	11	2	0	0
			Total	C	N	O	P		
3	D	1	28	10	5	11	2	0	0

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	13	6	7	0	0
			Total	C	O		
4	A	1	13	6	7	0	0
			Total	C	O		
4	C	1	13	6	7	0	0
			Total	C	O		
4	D	1	13	6	7	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
5	A	1	5	4	1	0	0
			Total	O	S		
5	A	1	5	4	1	0	0
			Total	O	S		
5	A	1	5	4	1	0	0
			Total	O	S		
5	A	1	5	4	1	0	0
			Total	O	S		
5	B	1	5	4	1	0	0
			Total	O	S		
5	B	1	5	4	1	0	0
			Total	O	S		
5	B	1	5	4	1	0	0
			Total	O	S		
5	B	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total O 2 2	0	0
6	B	1	Total O 1 1	0	0
6	C	4	Total O 4 4	0	0
6	D	3	Total O 3 3	0	0
6	F	1	Total O 1 1	0	0
6	G	1	Total O 1 1	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Guanine nucleotide-binding protein G(k) subunit alpha

Chain A:



• Molecule 1: Guanine nucleotide-binding protein G(k) subunit alpha

Chain B:



• Molecule 1: Guanine nucleotide-binding protein G(k) subunit alpha

Chain C:



• Molecule 1: Guanine nucleotide-binding protein G(k) subunit alpha

Chain D:



• Molecule 2: G-protein-signaling modulator 2

Chain E:



• Molecule 2: G-protein-signaling modulator 2

Chain F:

• Molecule 2: G-protein-signaling modulator 2

Chain G:

• Molecule 2: G-protein-signaling modulator 2

Chain H:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α, β, γ	207.33Å 207.33Å 236.54Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.90 43.04 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.90) 99.8 (43.04-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.69 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.208 , 0.244 0.216 , 0.252	Depositor DCC
R_{free} test set	3375 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å^2)	61.6	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 37.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	11190	wwPDB-VP
Average B, all atoms (Å^2)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.12% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CIT, GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.52	2/2601 (0.1%)	0.63	0/3511
1	B	0.52	0/2569	0.61	0/3467
1	C	0.53	2/2576 (0.1%)	0.61	0/3470
1	D	0.53	2/2636 (0.1%)	0.60	0/3556
2	E	0.58	0/178	0.79	0/237
2	F	0.61	0/185	0.81	0/245
2	G	0.42	0/189	0.60	0/250
2	H	0.45	0/187	0.62	0/248
All	All	0.53	6/11121 (0.1%)	0.62	0/14984

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	211	TRP	CD2-CE2	5.66	1.48	1.41
1	A	211	TRP	CD2-CE2	5.58	1.48	1.41
1	C	211	TRP	CD2-CE2	5.40	1.47	1.41
1	A	258	TRP	CD2-CE2	5.36	1.47	1.41
1	C	131	TRP	CD2-CE2	5.27	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	207	GLU	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2553	0	2500	25	0
1	B	2522	0	2469	27	0
1	C	2533	0	2489	25	0
1	D	2586	0	2528	21	0
2	E	177	0	176	0	0
2	F	184	0	189	9	0
2	G	188	0	193	9	0
2	H	186	0	189	6	0
3	A	28	0	12	1	0
3	B	28	0	12	1	0
3	C	28	0	12	1	0
3	D	28	0	12	0	0
4	A	26	0	10	0	0
4	C	13	0	5	0	0
4	D	13	0	5	1	0
5	A	20	0	0	0	0
5	B	20	0	0	0	0
5	C	20	0	0	1	0
5	D	25	0	0	0	0
6	A	2	0	0	0	0
6	B	1	0	0	0	0
6	C	4	0	0	3	0
6	D	3	0	0	0	0
6	F	1	0	0	4	0
6	G	1	0	0	0	0
All	All	11190	0	10801	116	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 116 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:622:GLU:OE2	2:G:630:ARG:NE	1.76	1.16
2:F:635:ARG:HB3	6:F:701:HOH:O	1.63	0.97
2:G:622:GLU:OE1	2:G:630:ARG:NH1	2.07	0.87
2:G:622:GLU:CD	2:G:630:ARG:HH11	1.77	0.86
2:F:640:ARG:HD2	6:F:701:HOH:O	1.80	0.82

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/330 (97%)	302 (95%)	16 (5%)	1 (0%)	41	71
1	B	312/330 (94%)	299 (96%)	13 (4%)	0	100	100
1	C	308/330 (93%)	292 (95%)	15 (5%)	1 (0%)	41	71
1	D	323/330 (98%)	306 (95%)	17 (5%)	0	100	100
2	E	20/26 (77%)	17 (85%)	3 (15%)	0	100	100
2	F	20/26 (77%)	19 (95%)	1 (5%)	0	100	100
2	G	20/26 (77%)	20 (100%)	0	0	100	100
2	H	21/26 (81%)	20 (95%)	1 (5%)	0	100	100
All	All	1343/1424 (94%)	1275 (95%)	66 (5%)	2 (0%)	51	82

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	208	ARG
1	C	118	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/290 (95%)	271 (98%)	5 (2%)	59	85
1	B	275/290 (95%)	269 (98%)	6 (2%)	52	81
1	C	276/290 (95%)	270 (98%)	6 (2%)	52	81
1	D	278/290 (96%)	269 (97%)	9 (3%)	39	73
2	E	18/25 (72%)	16 (89%)	2 (11%)	6	19
2	F	20/25 (80%)	19 (95%)	1 (5%)	24	57
2	G	21/25 (84%)	20 (95%)	1 (5%)	25	58
2	H	19/25 (76%)	19 (100%)	0	100	100
All	All	1183/1260 (94%)	1153 (98%)	30 (2%)	47	78

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	201	VAL
2	E	642	LEU
1	D	75	SER
2	G	642	LEU
1	D	260	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	79	GLN
1	D	106	GLN
2	F	632	GLN
1	A	204	GLN
1	A	79	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

25 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	CIT	A	402	-	12,12,12	1.45	1 (8%)	17,17,17	5.74	10 (58%)
4	CIT	D	402	-	12,12,12	1.47	2 (16%)	17,17,17	4.89	8 (47%)
5	SO4	B	405	-	4,4,4	0.34	0	6,6,6	0.11	0
5	SO4	D	405	-	4,4,4	0.34	0	6,6,6	0.11	0
5	SO4	D	403	-	4,4,4	0.45	0	6,6,6	0.15	0
5	SO4	D	404	-	4,4,4	0.40	0	6,6,6	0.11	0
5	SO4	B	402	-	4,4,4	0.43	0	6,6,6	0.28	0
5	SO4	B	404	-	4,4,4	0.42	0	6,6,6	0.16	0
4	CIT	A	403	-	12,12,12	1.14	1 (8%)	17,17,17	5.26	10 (58%)
4	CIT	C	402	-	12,12,12	1.57	2 (16%)	17,17,17	5.07	8 (47%)
3	GDP	A	401	-	24,30,30	0.88	1 (4%)	30,47,47	1.44	7 (23%)
3	GDP	D	401	-	24,30,30	1.05	2 (8%)	30,47,47	1.48	4 (13%)
3	GDP	C	401	-	24,30,30	0.98	2 (8%)	30,47,47	1.18	2 (6%)
5	SO4	A	405	-	4,4,4	0.38	0	6,6,6	0.22	0
3	GDP	B	401	-	24,30,30	1.05	1 (4%)	30,47,47	1.64	6 (20%)
5	SO4	C	406	-	4,4,4	0.37	0	6,6,6	0.14	0
5	SO4	A	406	-	4,4,4	0.39	0	6,6,6	0.14	0
5	SO4	C	405	-	4,4,4	0.35	0	6,6,6	0.14	0
5	SO4	D	406	-	4,4,4	0.37	0	6,6,6	0.28	0
5	SO4	C	403	-	4,4,4	0.38	0	6,6,6	0.13	0
5	SO4	A	404	-	4,4,4	0.39	0	6,6,6	0.17	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	SO4	A	407	-	4,4,4	0.35	0	6,6,6	0.24	0
5	SO4	C	404	-	4,4,4	0.36	0	6,6,6	0.26	0
5	SO4	B	403	-	4,4,4	0.38	0	6,6,6	0.08	0
5	SO4	D	407	-	4,4,4	0.35	0	6,6,6	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIT	A	403	-	-	4/16/16/16	-
3	GDP	C	401	-	-	0/12/32/32	0/3/3/3
4	CIT	A	402	-	-	8/16/16/16	-
4	CIT	D	402	-	-	3/16/16/16	-
3	GDP	B	401	-	-	0/12/32/32	0/3/3/3
4	CIT	C	402	-	-	9/16/16/16	-
3	GDP	A	401	-	-	2/12/32/32	0/3/3/3
3	GDP	D	401	-	-	3/12/32/32	0/3/3/3

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	GDP	O4'-C1'	3.26	1.45	1.41
3	D	401	GDP	O4'-C1'	3.17	1.45	1.41
4	D	402	CIT	C3-C6	3.10	1.56	1.53
4	C	402	CIT	C2-C3	-2.78	1.50	1.53
4	C	402	CIT	C3-C6	2.55	1.56	1.53

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	A	402	CIT	O7-C3-C6	-17.14	84.79	108.86
4	C	402	CIT	O7-C3-C6	-13.76	89.54	108.86
4	A	403	CIT	O7-C3-C6	-12.98	90.64	108.86
4	C	402	CIT	C2-C3-C6	-11.48	85.44	110.11
4	D	402	CIT	O7-C3-C6	-11.11	93.26	108.86

There are no chirality outliers.

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	GDP	PA-O3A-PB-O2B
3	A	401	GDP	PA-O3A-PB-O3B
3	D	401	GDP	PA-O3A-PB-O2B
3	D	401	GDP	PA-O3A-PB-O3B
4	A	402	CIT	C1-C2-C3-O7

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	402	CIT	1	0
3	A	401	GDP	1	0
3	C	401	GDP	1	0
3	B	401	GDP	1	0
5	C	406	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand GDP A 401



Bond lengths



Bond angles



Torsions



Rings

Ligand GDP D 401



Bond lengths



Bond angles



Torsions



Rings

Ligand GDP C 401



Ligand GDP B 401



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	320/330 (96%)	-0.12	3 (0%)	84	84	42, 63, 101, 125	0
1	B	315/330 (95%)	-0.20	6 (1%)	66	65	39, 59, 90, 113	0
1	C	316/330 (95%)	-0.21	4 (1%)	77	77	39, 65, 113, 134	0
1	D	324/330 (98%)	-0.16	7 (2%)	62	59	39, 63, 104, 134	0
2	E	22/26 (84%)	-0.19	0	100	100	59, 72, 88, 97	0
2	F	22/26 (84%)	-0.06	0	100	100	51, 67, 86, 97	0
2	G	22/26 (84%)	-0.02	1 (4%)	33	29	52, 69, 84, 98	0
2	H	23/26 (88%)	0.31	1 (4%)	35	31	62, 80, 113, 136	0
All	All	1364/1424 (95%)	-0.16	22 (1%)	72	71	39, 64, 104, 136	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	117	GLY	5.6
1	B	207	GLU	4.8
1	D	316	THR	4.6
1	D	116	GLU	4.4
1	C	354	TYR	4.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	SO4	D	406	5/5	0.69	0.24	128,130,133,138	0
5	SO4	D	404	5/5	0.79	0.30	137,142,150,151	0
5	SO4	B	405	5/5	0.80	0.35	138,143,145,147	0
5	SO4	A	404	5/5	0.80	0.20	117,124,135,135	0
5	SO4	B	404	5/5	0.80	0.31	120,124,130,139	0
5	SO4	C	405	5/5	0.81	0.27	138,144,149,151	0
5	SO4	D	405	5/5	0.82	0.30	150,153,158,160	0
5	SO4	C	406	5/5	0.83	0.23	120,129,136,137	0
5	SO4	B	403	5/5	0.85	0.21	123,130,139,141	0
5	SO4	C	404	5/5	0.86	0.22	116,123,127,127	0
5	SO4	A	407	5/5	0.87	0.28	106,115,122,127	0
5	SO4	B	402	5/5	0.88	0.18	97,111,117,119	0
5	SO4	D	407	5/5	0.89	0.37	136,137,139,140	0
5	SO4	C	403	5/5	0.90	0.16	104,109,111,115	0
5	SO4	A	405	5/5	0.90	0.15	110,110,114,119	0
5	SO4	A	406	5/5	0.93	0.18	103,106,107,108	0
5	SO4	D	403	5/5	0.96	0.13	99,105,106,106	0
4	CIT	A	402	13/13	0.96	0.18	55,60,64,72	0
4	CIT	A	403	13/13	0.96	0.18	55,58,62,64	0
4	CIT	C	402	13/13	0.96	0.16	45,48,59,60	0
4	CIT	D	402	13/13	0.96	0.17	45,46,54,55	0
3	GDP	D	401	28/28	0.98	0.15	39,46,51,54	0
3	GDP	A	401	28/28	0.98	0.19	46,57,61,61	0
3	GDP	B	401	28/28	0.99	0.18	40,47,51,52	0
3	GDP	C	401	28/28	0.99	0.14	46,52,55,57	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GDP D 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around GDP A 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around GDP B 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around GDP C 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.