



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:49 PM UTC

PDB ID : 9G65 / pdb_00009g65
Title : Crystal structure of the engineered photoenzyme SpEnT1.3
Authors : Hardy, F.J.; Roberts, G.W.
Deposited on : 2024-07-18
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5140 atoms, of which 2472 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Engineered photoenzyme SpEnT1.3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	314	4920	1584	2448	413	455	20	0	7	0

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	21	ALA	GLU	engineered mutation	UNP Q7SIG4
A	23	CYS	PRO	engineered mutation	UNP Q7SIG4
A	37	ALA	GLU	engineered mutation	UNP Q7SIG4
A	120	ALA	ASN	engineered mutation	UNP Q7SIG4
A	121	GLY	ASP	engineered mutation	UNP Q7SIG4
A	144	PHE	TYR	engineered mutation	UNP Q7SIG4
A	146	ILE	ARG	engineered mutation	UNP Q7SIG4
A	148	LEU	MET	engineered mutation	UNP Q7SIG4
A	173	ALA	PHE	engineered mutation	UNP Q7SIG4
A	175	ALA	ASN	engineered mutation	UNP Q7SIG4
A	195	GLN	THR	engineered mutation	UNP Q7SIG4
A	225	LYS	GLU	engineered mutation	UNP Q7SIG4
A	226	LEU	GLY	engineered mutation	UNP Q7SIG4
A	229	ALA	ASP	engineered mutation	UNP Q7SIG4
A	244	A1IHG	TRP	engineered mutation	UNP Q7SIG4
A	271	CYS	SER	engineered mutation	UNP Q7SIG4
A	272	ALA	ASN	engineered mutation	UNP Q7SIG4
A	287	LEU	HIS	engineered mutation	UNP Q7SIG4
A	315	GLY	-	expression tag	UNP Q7SIG4
A	316	SER	-	expression tag	UNP Q7SIG4
A	317	LEU	-	expression tag	UNP Q7SIG4
A	318	GLU	-	expression tag	UNP Q7SIG4
A	319	TRP	-	expression tag	UNP Q7SIG4
A	320	SER	-	expression tag	UNP Q7SIG4
A	321	HIS	-	expression tag	UNP Q7SIG4
A	322	PRO	-	expression tag	UNP Q7SIG4
A	323	GLN	-	expression tag	UNP Q7SIG4

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	324	PHE	-	expression tag	UNP Q7SIG4
A	325	GLU	-	expression tag	UNP Q7SIG4
A	326	LYS	-	expression tag	UNP Q7SIG4

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 3 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			24	6	14	4		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mg	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	177	Total	O	0	0
			177	177		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Engineered photoenzyme SpEnT1.3



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	87.24Å 78.51Å 43.16Å 90.00° 95.13° 90.00°	Depositor
Resolution (Å)	43.45 – 1.80 43.45 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.6 (43.45-1.80) 99.8 (43.45-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.12 (at 1.80Å)	Xtriage
Refinement program	PHENIX 1.21_5207	Depositor
R, R_{free}	0.196 , 0.232 0.200 , 0.239	Depositor DCC
R_{free} test set	1328 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	35.5	Xtriage
Anisotropy	0.298	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5140	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1IHG, PGE, MG, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.22	0/2535	0.45	0/3428

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2472	2448	2425	36	1
2	A	7	10	10	1	0
3	A	10	14	14	0	0
4	A	2	0	0	0	0
5	A	177	0	0	8	1
All	All	2668	2472	2449	37	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:204:ASP:OD1	5:A:601:HOH:O	1.85	0.95
1:A:296:GLU:OE2	5:A:602:HOH:O	2.12	0.67
1:A:50:ARG:NH1	5:A:605:HOH:O	2.25	0.62
1:A:23:CYS:HB3	1:A:33:ILE:HD13	1.84	0.57
1:A:206:LYS:NZ	5:A:609:HOH:O	2.41	0.54
1:A:7:GLU:HG2	1:A:264:ARG:HB3	1.88	0.54
1:A:90[A]:MET:CE	1:A:146:ILE:HD11	2.42	0.50
1:A:225:LYS:H	1:A:225:LYS:HD3	1.77	0.49
1:A:90[B]:MET:HE3	1:A:135:PRO:HG2	1.94	0.49
1:A:17[A]:ILE:HD11	1:A:58:LYS:CD	2.44	0.48
1:A:220:ILE:HG13	1:A:228:ALA:HB2	1.95	0.48
1:A:182:MET:HE3	1:A:186:ARG:HB2	1.95	0.48
1:A:146:ILE:HG22	1:A:148:LEU:H	1.79	0.47
1:A:17[A]:ILE:HD11	1:A:58:LYS:HD2	1.97	0.47
1:A:182:MET:HG3	1:A:188:TYR:HB2	1.96	0.47
1:A:100:ASP:OD1	1:A:102:THR:HG23	2.14	0.46
1:A:90[A]:MET:HE1	1:A:146:ILE:HD11	1.97	0.46
1:A:19:GLY:O	1:A:20:ALA:C	2.59	0.45
1:A:133:THR:HB	1:A:174:PRO:HB2	1.99	0.45
1:A:214:LYS:NZ	5:A:615:HOH:O	2.45	0.45
1:A:146:ILE:CG2	1:A:148:LEU:HG	2.47	0.45
1:A:225:LYS:HD3	1:A:225:LYS:N	2.32	0.45
1:A:90[B]:MET:HE1	1:A:146:ILE:HD11	1.99	0.44
1:A:234:ASP:OD1	1:A:234:ASP:C	2.61	0.43
1:A:275:PHE:CZ	1:A:282:ILE:HD11	2.53	0.43
1:A:19:GLY:O	1:A:36:PRO:HD2	2.20	0.42
2:A:501:PEG:H11	5:A:680:HOH:O	2.19	0.42
1:A:31:PHE:CE2	1:A:33:ILE:HD11	2.55	0.42
1:A:72:ILE:HB	1:A:90[A]:MET:HB2	2.01	0.41
1:A:127:GLU:OE2	5:A:603:HOH:O	2.21	0.41
1:A:195:GLN:HB3	1:A:196:PRO:HD3	2.02	0.41
1:A:21:ALA:O	1:A:22:GLY:C	2.64	0.41
1:A:224:HIS:HB2	5:A:611:HOH:O	2.21	0.41
1:A:46:GLY:HA3	1:A:73:PRO:HD2	2.02	0.40
1:A:110:ASP:OD1	1:A:110:ASP:C	2.63	0.40
1:A:79:ASP:HB3	1:A:82:ALA:O	2.22	0.40
1:A:297:TRP:CG	1:A:298:GLN:H	2.38	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:THR:O	5:A:605:HOH:O[1_556]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/326 (98%)	302 (95%)	16 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/266 (98%)	259 (99%)	3 (1%)	65	60

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	223	THR
1	A	273[A]	LEU
1	A	273[B]	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	98	GLN
1	A	183	ASN
1	A	224	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	A1IHG	A	244	1	21,22,23	2.53	11 (52%)	26,31,33	2.77	10 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	A1IHG	A	244	1	-	0/5/6/8	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	244	A1IHG	CZ-S16	5.01	1.84	1.76
1	A	244	A1IHG	C15-S16	4.82	1.84	1.76
1	A	244	A1IHG	C10-C09	4.34	1.57	1.48
1	A	244	A1IHG	CE2-C09	3.12	1.54	1.48
1	A	244	A1IHG	CE1-CD1	2.65	1.43	1.38
1	A	244	A1IHG	O-C	2.57	1.29	1.20
1	A	244	A1IHG	C11-C10	2.53	1.43	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	244	A1IHG	CD1-CG	2.28	1.43	1.38
1	A	244	A1IHG	C12-C11	2.24	1.42	1.38
1	A	244	A1IHG	CE1-CZ	2.19	1.43	1.39
1	A	244	A1IHG	O17-C09	2.08	1.26	1.22

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	244	A1IHG	C10-C09-CE2	7.85	131.08	117.44
1	A	244	A1IHG	C15-S16-CZ	5.13	113.10	100.34
1	A	244	A1IHG	O17-C09-C10	-4.95	113.84	120.99
1	A	244	A1IHG	O17-C09-CE2	-4.10	115.08	120.99
1	A	244	A1IHG	C15-C10-C09	-3.60	118.98	123.54
1	A	244	A1IHG	C11-C10-C09	3.31	124.42	119.26
1	A	244	A1IHG	CZ-CE2-C09	-2.85	119.93	123.54
1	A	244	A1IHG	C14-C15-C10	2.53	122.30	119.16
1	A	244	A1IHG	CE1-CZ-CE2	2.51	122.27	119.16
1	A	244	A1IHG	CD2-CE2-C09	2.43	123.57	119.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PGE	A	502	-	9,9,9	0.32	0	8,8,8	0.55	0
2	PEG	A	501	-	6,6,6	0.23	0	5,5,5	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	A	502	-	-	5/7/7/7	-
2	PEG	A	501	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	PGE	O3-C5-C6-O4
3	A	502	PGE	O2-C3-C4-O3
2	A	501	PEG	O1-C1-C2-O2
3	A	502	PGE	C1-C2-O2-C3
3	A	502	PGE	C3-C4-O3-C5
2	A	501	PEG	C1-C2-O2-C3
3	A	502	PGE	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	PEG	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	313/326 (96%)	-0.05	3 (0%) 79 80	19, 45, 68, 122	6 (1%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	314	PHE	4.2
1	A	148	LEU	2.7
1	A	1	MET	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	A1IHG	A	244	20/21	0.93	0.09	37,54,72,75	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PGE	A	502	10/10	0.78	0.17	41,53,64,64	24
2	PEG	A	501	7/7	0.84	0.13	42,57,68,70	0
4	MG	A	503	1/1	0.98	0.04	32,32,32,32	0
4	MG	A	504	1/1	0.99	0.05	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.