



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2025 – 04:19 pm BST

PDB ID : 9G88 / pdb\_00009g88  
Title : Carotenoid cleavage oxygenase from Moesziomyces aphidis bound to acetate  
Authors : Plewka, J.; Schober, L.; Magiera-Mularz, K.; Rudroff, F.; Winkler, M.  
Deposited on : 2024-07-23  
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.44


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2247 (1.40-1.40)
Clashscore	180529	2446 (1.40-1.40)
Ramachandran outliers	177936	2398 (1.40-1.40)
Sidechain outliers	177891	2397 (1.40-1.40)
RSRZ outliers	164620	2246 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	556	 88% 6% • 6%
1	B	556	 88% 6% • 6%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9789 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lignostilbene dioxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	525	Total	C	N	O	S	0	3	0
			4235	2704	727	788	16			
1	B	525	Total	C	N	O	S	0	2	0
			4224	2698	724	786	16			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP W3VHW6
A	-6	TRP	-	expression tag	UNP W3VHW6
A	-5	SER	-	expression tag	UNP W3VHW6
A	-4	HIS	-	expression tag	UNP W3VHW6
A	-3	PRO	-	expression tag	UNP W3VHW6
A	-2	GLN	-	expression tag	UNP W3VHW6
A	-1	PHE	-	expression tag	UNP W3VHW6
A	0	GLU	-	expression tag	UNP W3VHW6
A	1	LYS	-	expression tag	UNP W3VHW6
B	-7	MET	-	initiating methionine	UNP W3VHW6
B	-6	TRP	-	expression tag	UNP W3VHW6
B	-5	SER	-	expression tag	UNP W3VHW6
B	-4	HIS	-	expression tag	UNP W3VHW6
B	-3	PRO	-	expression tag	UNP W3VHW6
B	-2	GLN	-	expression tag	UNP W3VHW6
B	-1	PHE	-	expression tag	UNP W3VHW6
B	0	GLU	-	expression tag	UNP W3VHW6
B	1	LYS	-	expression tag	UNP W3VHW6

- Molecule 2 is ACETATE ION (CCD ID: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is FE (II) ION (CCD ID: FE2) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe	0	0
			1	1		
3	B	1	Total	Fe	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	698	Total	O	0	0
			698	698		
4	B	622	Total	O	0	0
			622	622		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lignostilbene dioxygenase



- Molecule 1: Lignostilbene dioxygenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.16Å 85.96Å 93.87Å 90.00° 100.34° 90.00°	Depositor
Resolution (Å)	46.17 – 1.40 46.17 – 1.40	Depositor EDS
% Data completeness (in resolution range)	97.5 (46.17-1.40) 97.5 (46.17-1.40)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.41 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, $R_{free}$	0.168 , 0.193 0.168 , 0.193	Depositor DCC
$R_{free}$ test set	10868 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.5	Xtriage
Anisotropy	0.178	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 30.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	9789	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, FE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.78	0/4367	1.18	14/5944 (0.2%)
1	B	0.75	1/4356 (0.0%)	1.17	14/5929 (0.2%)
All	All	0.77	1/8723 (0.0%)	1.17	28/11873 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	1
All	All	0	5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	266	PRO	CA-CB	5.86	1.57	1.53

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	412	PHE	CA-CB-CG	7.65	121.45	113.80
1	A	73	ARG	CG-CD-NE	-6.93	96.76	112.00
1	B	252	ASP	CA-CB-CG	6.77	119.37	112.60
1	B	517	GLU	CB-CA-C	6.54	124.13	110.32
1	A	324	GLU	CB-CG-CD	6.30	123.31	112.60
1	B	164	ARG	CB-CG-CD	-6.24	96.94	111.30
1	B	503	PHE	CA-CB-CG	6.24	120.04	113.80
1	B	73	ARG	CG-CD-NE	6.16	125.56	112.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	410	ASN	CA-CB-CG	5.96	118.56	112.60
1	A	410	ASN	CA-CB-CG	5.86	118.46	112.60
1	A	164	ARG	CD-NE-CZ	5.83	132.57	124.40
1	A	77	ASP	CA-CB-CG	5.83	118.43	112.60
1	A	252	ASP	CA-CB-CG	5.64	118.24	112.60
1	A	340	PHE	CA-CB-CG	5.58	119.39	113.80
1	A	412	PHE	CA-CB-CG	5.52	119.32	113.80
1	B	272	GLU	CB-CG-CD	5.52	121.98	112.60
1	A	509	ILE	CB-CA-C	5.51	120.33	111.29
1	A	416	ARG	CG-CD-NE	5.45	123.99	112.00
1	B	340	PHE	CA-CB-CG	5.43	119.23	113.80
1	B	396	ASP	CA-CB-CG	5.41	118.01	112.60
1	A	373	LYS	CB-CG-CD	5.41	123.73	111.30
1	A	161	HIS	CA-CB-CG	-5.39	108.41	113.80
1	B	359	THR	CA-CB-OG1	-5.29	101.67	109.60
1	B	77	ASP	CA-CB-CG	5.22	117.82	112.60
1	A	109	LYS	CB-CG-CD	5.18	123.22	111.30
1	B	73	ARG	CA-CB-CG	5.10	124.30	114.10
1	A	73	ARG	CA-CB-CG	-5.07	103.96	114.10
1	B	509	ILE	CB-CA-C	5.03	119.54	111.29

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	164	ARG	Sidechain
1	A	238	ARG	Sidechain
1	A	416	ARG	Sidechain
1	A	459	ARG	Sidechain
1	B	238	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4235	0	4057	19	0
1	B	4224	0	4049	16	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	4	0	3	0	0
2	B	4	0	3	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	698	0	0	11	0
4	B	622	0	0	4	0
All	All	9789	0	8112	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:THR:CG2	4:A:900:HOH:O	2.02	1.04
1:A:156:THR:HG22	4:A:1202:HOH:O	1.64	0.96
1:B:143:HIS:HD2	1:B:145:CYS:H	1.29	0.80
1:B:24:TYR:N	4:B:701:HOH:O	2.13	0.80
4:A:705:HOH:O	1:B:58:MET:HG2	1.94	0.66
1:A:308:HIS:HE1	4:A:1255:HOH:O	1.85	0.60
1:A:156:THR:HG21	4:A:900:HOH:O	1.86	0.58
1:A:139:PRO:HA	1:A:142:HIS:CD2	2.38	0.58
1:A:515:PRO:HG3	4:A:1319:HOH:O	2.04	0.58
1:A:308:HIS:HD2	4:A:1180:HOH:O	1.88	0.56
1:B:143:HIS:CD2	1:B:145:CYS:H	2.17	0.55
1:B:139:PRO:HA	1:B:142:HIS:CD2	2.43	0.54
1:B:124:VAL:HG11	1:B:146:VAL:HG11	1.93	0.51
1:B:279:HIS:HD2	4:B:724:HOH:O	1.95	0.49
1:B:279:HIS:HE1	4:B:877:HOH:O	1.94	0.49
1:A:480:ASN:HB3	1:B:480:ASN:HB3	1.96	0.47
1:A:514[A]:ILE:HD13	4:A:1079:HOH:O	2.14	0.46
1:B:389:PRO:HB2	1:B:412:PHE:CD1	2.50	0.46
1:B:137:ARG:HA	1:B:151:GLU:OE2	2.16	0.45
1:A:314:PRO:HA	1:A:333:SER:O	2.17	0.45
1:A:494:LYS:HE2	4:A:1254:HOH:O	2.16	0.45
1:B:77:ASP:O	1:B:78:TYR:C	2.60	0.44
1:A:109:LYS:HB2	4:A:737:HOH:O	2.17	0.44
1:A:77:ASP:O	1:A:78:TYR:C	2.61	0.44
1:A:72:TYR:CD2	1:A:514[B]:ILE:HD12	2.53	0.43
1:A:213:PHE:HA	1:A:226:TYR:O	2.18	0.43
1:A:389:PRO:HB2	1:A:412:PHE:CD1	2.54	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:308:HIS:HD2	4:B:1124:HOH:O	2.02	0.43
1:B:160:MET:HB2	1:B:514[A]:ILE:HD13	2.01	0.42
1:B:213:PHE:HA	1:B:226:TYR:O	2.19	0.42
1:A:156:THR:HG23	1:A:509:ILE:HG21	2.00	0.42
1:A:100:VAL:O	1:A:114[B]:GLN:HA	2.21	0.41
1:A:68:HIS:HD2	4:A:963:HOH:O	2.03	0.41
1:B:58:MET:HE3	1:B:58:MET:HB2	1.75	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	526/556 (95%)	513 (98%)	12 (2%)	1 (0%)	44	20
1	B	525/556 (94%)	512 (98%)	13 (2%)	0	100	100
All	All	1051/1112 (94%)	1025 (98%)	25 (2%)	1 (0%)	48	22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	509	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	452/475 (95%)	445 (98%)	7 (2%)	60	32
1	B	451/475 (95%)	445 (99%)	6 (1%)	65	38
All	All	903/950 (95%)	890 (99%)	13 (1%)	62	35

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	ARG
1	A	123	LYS
1	A	242	LEU
1	A	355	GLU
1	A	391	GLU
1	A	416	ARG
1	A	420	ASN
1	B	123	LYS
1	B	242	LEU
1	B	391	GLU
1	B	461	LYS
1	B	503	PHE
1	B	517	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	HIS
1	A	148	GLN
1	A	173	ASN
1	A	308	HIS
1	A	541	GLN
1	B	114	GLN
1	B	143	HIS
1	B	148	GLN
1	B	173	ASN
1	B	279	HIS
1	B	308	HIS
1	B	336	HIS
1	B	397	ASN
1	B	410	ASN
1	B	422	ASN
1	B	541	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ACT	A	601	3	3,3,3	0.79	0	3,3,3	0.95	0
2	ACT	B	601	3	3,3,3	1.22	0	3,3,3	1.11	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	525/556 (94%)	-0.45	2 (0%)	89 90	6, 16, 29, 49	3 (0%)
1	B	525/556 (94%)	-0.09	7 (1%)	74 76	8, 20, 36, 74	2 (0%)
All	All	1050/1112 (94%)	-0.27	9 (0%)	81 83	6, 18, 33, 74	5 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	419	GLY	3.4
1	B	412	PHE	3.1
1	A	25	VAL	2.6
1	B	418	ASP	2.5
1	B	416	ARG	2.4
1	B	414	PRO	2.3
1	B	420	ASN	2.2
1	B	356	ALA	2.2
1	A	50	PHE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ACT	A	601	4/4	0.97	0.07	15,16,17,18	0
2	ACT	B	601	4/4	0.97	0.08	17,19,21,22	0
3	FE2	B	602	1/1	0.99	0.04	17,17,17,17	0
3	FE2	A	602	1/1	1.00	0.04	14,14,14,14	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








**Electron density around ACT B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FE2 B 602:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FE2 A 602:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.