



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:08 PM UTC

PDB ID : 9GEI / pdb_00009gei
Title : Entamoeba histolytica Gal/GalNAc lectin heavy chain residues 808-992 in the presence of monoclonal antibody CP33-H/L-LA22
Authors : Gerard, S.F.; Higgins, M.K.
Deposited on : 2024-08-07
Resolution : 3.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3685 (3.08-3.00)
Clashscore	190562	4007 (3.08-3.00)
Ramachandran outliers	187476	3834 (3.08-3.00)
Sidechain outliers	187428	3836 (3.08-3.00)
RSRZ outliers	180081	3684 (3.08-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	188	
1	C	188	
2	B	253	
2	D	253	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6268 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Galactose/N-acetyl-D-galactosamine lectin heavy subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	174	Total	C	N	O	S	0	0	0
			1408	881	228	284	15			
1	C	174	Total	C	N	O	S	0	0	0
			1408	881	228	284	15			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	972	SER	CYS	conflict	UNP P32022
A	993	GLY	THR	conflict	UNP P32022
C	972	SER	CYS	conflict	UNP P32022
C	993	GLY	THR	conflict	UNP P32022


- Molecule 2 is a protein called CP33-H/L-LA22.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	226	Total	C	N	O	S	0	0	0
			1726	1076	300	343	7			
2	D	226	Total	C	N	O	S	0	0	0
			1726	1076	300	343	7			



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Galactose/N-acetyl-D-galactosamine lectin heavy subunit 1




- Molecule 1: Galactose/N-acetyl-D-galactosamine lectin heavy subunit 1



- Molecule 2: CP33-H/L-LA22



- Molecule 2: CP33-H/L-LA22



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	134.10Å 151.98Å 60.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.99 – 3.03 75.99 – 3.03	Depositor EDS
% Data completeness (in resolution range)	65.6 (75.99-3.03) 65.6 (75.99-3.03)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 3.01Å)	Xtriage
Refinement program	BUSTER 2.10.4 (26-JUL-2023)	Depositor
R, R_{free}	0.241 , 0.277 0.243 , 0.290	Depositor DCC
R_{free} test set	779 reflections (3.17%)	wwPDB-VP
Wilson B-factor (Å ²)	85.6	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 87.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6268	wwPDB-VP
Average B, all atoms (Å ²)	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/1432	1.00	0/1931
1	C	0.66	1/1432 (0.1%)	0.98	0/1931
2	B	0.61	0/1766	0.94	1/2397 (0.0%)
2	D	0.59	0/1766	0.93	1/2397 (0.0%)
All	All	0.63	1/6396 (0.0%)	0.96	2/8656 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	971	GLU	CA-C	5.21	1.57	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	208	GLY	N-CA-C	5.69	117.36	111.95
2	D	208	GLY	N-CA-C	5.57	117.24	111.95

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1408	0	1354	9	0
1	C	1408	0	1354	10	0
2	B	1726	0	1644	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1726	0	1644	22	0
All	All	6268	0	5996	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:821:LYS:HB2	1:A:823:GLN:HE21	1.44	0.82
1:C:821:LYS:HB2	1:C:823:GLN:HE21	1.45	0.79
1:C:828:LYS:HE2	1:C:838:LYS:HE3	1.69	0.74
1:C:971:GLU:HG3	1:C:972:SER:H	1.57	0.69
2:B:179:GLN:HB2	2:B:189:LEU:HD11	1.75	0.68
2:D:43:ALA:HB3	2:D:46:LYS:HB2	1.75	0.68
2:D:179:GLN:HB2	2:D:189:LEU:HD11	1.75	0.68
2:B:43:ALA:HB3	2:B:46:LYS:HB2	1.75	0.67
2:D:171:ILE:HG22	2:D:234:TYR:HB3	1.79	0.64
2:D:16:GLN:HE22	2:D:19:ARG:HH21	1.45	0.63
1:A:858:CYS:HB2	1:A:881:ALA:HB1	1.83	0.60
2:B:171:ILE:HG22	2:B:234:TYR:HB3	1.84	0.59
2:D:6:LYS:HE3	2:D:8:MET:HE3	1.83	0.59
2:B:91:PRO:HA	2:B:122:VAL:HB	1.85	0.58
1:C:858:CYS:HB2	1:C:881:ALA:HB1	1.85	0.58
2:B:16:GLN:HE22	2:B:19:ARG:HH21	1.51	0.58
2:D:91:PRO:HA	2:D:122:VAL:HB	1.85	0.58
2:B:4:GLU:HB2	2:B:6:LYS:HZ2	1.68	0.58
2:D:38:HIS:CE1	2:D:53:ARG:HG3	2.39	0.57
2:D:16:GLN:NE2	2:D:19:ARG:HH21	2.04	0.55
1:A:928:LEU:HD22	1:A:932:ARG:HA	1.89	0.54
1:C:928:LEU:HD22	1:C:932:ARG:HA	1.89	0.54
2:D:148:GLN:H	2:D:242:GLN:NE2	2.06	0.54
1:A:884:HIS:CD2	1:A:915:THR:HG23	2.43	0.54
1:A:918:ILE:HD12	1:A:926:CYS:HB3	1.89	0.54
2:B:16:GLN:NE2	2:B:19:ARG:HH21	2.07	0.53
1:A:945:THR:HG23	2:B:105:SER:HB3	1.91	0.52
2:B:171:ILE:HG21	2:B:232:GLN:HB2	1.93	0.51
2:D:154:SER:HA	2:D:247:GLU:O	2.11	0.51
2:D:107:PRO:HB3	2:D:234:TYR:HA	1.93	0.51
2:B:4:GLU:HB2	2:B:6:LYS:NZ	2.24	0.51
1:C:918:ILE:HD12	1:C:926:CYS:HB3	1.92	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:971:GLU:HG3	1:C:972:SER:N	2.27	0.50
2:D:22:ARG:NH2	2:D:83:TYR:CE1	2.79	0.50
2:B:107:PRO:HB3	2:B:234:TYR:HA	1.93	0.49
2:D:225:PHE:HE2	2:D:248:ILE:HG12	1.78	0.49
2:B:153:LEU:HD11	2:B:246:VAL:HG13	1.94	0.49
2:D:94:THR:HG23	2:D:121:THR:HA	1.95	0.48
2:D:22:ARG:HD2	2:D:85:GLN:HG2	1.95	0.48
2:B:14:VAL:HG22	2:B:121:THR:HB	1.95	0.47
2:B:94:THR:HG23	2:B:121:THR:HA	1.97	0.47
2:D:14:VAL:HG22	2:D:121:THR:HB	1.95	0.47
1:A:884:HIS:NE2	1:A:915:THR:HG21	2.33	0.44
2:D:171:ILE:HG21	2:D:232:GLN:HB2	1.99	0.44
1:C:970:ILE:HG13	1:C:973:LYS:HB2	1.99	0.44
2:D:175:LEU:HD11	2:D:230:CYS:HB2	1.99	0.43
1:A:928:LEU:HA	1:A:935:VAL:HG12	2.01	0.43
1:C:928:LEU:HA	1:C:935:VAL:HG12	2.01	0.43
1:A:884:HIS:NE2	1:A:915:THR:CG2	2.83	0.42
2:B:220:LEU:HD13	2:B:248:ILE:HG12	2.01	0.42
2:D:225:PHE:CE2	2:D:248:ILE:HG12	2.54	0.41
1:C:980:THR:HG23	1:C:987:THR:HB	2.01	0.41
2:D:153:LEU:HD11	2:D:246:VAL:HG13	2.02	0.41
2:D:100:ALA:HB1	2:D:111:MET:HB3	2.02	0.41
2:B:100:ALA:HB1	2:B:111:MET:HB3	2.02	0.41
2:D:50:TRP:HZ2	2:D:53:ARG:HB2	1.85	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	171/188 (91%)	163 (95%)	8 (5%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	171/188 (91%)	160 (94%)	11 (6%)	0	100	100
2	B	222/253 (88%)	209 (94%)	13 (6%)	0	100	100
2	D	222/253 (88%)	209 (94%)	13 (6%)	0	100	100
All	All	786/882 (89%)	741 (94%)	45 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/176 (93%)	156 (95%)	8 (5%)	22	53
1	C	164/176 (93%)	152 (93%)	12 (7%)	13	39
2	B	188/203 (93%)	177 (94%)	11 (6%)	18	47
2	D	188/203 (93%)	176 (94%)	12 (6%)	16	44
All	All	704/758 (93%)	661 (94%)	43 (6%)	17	46

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	848	GLU
1	A	928	LEU
1	A	930	GLU
1	A	945	THR
1	A	970	ILE
1	A	973	LYS
1	A	990	CYS
1	A	991	LYS
2	B	4	GLU
2	B	24	SER
2	B	31	ARG
2	B	41	ARG
2	B	101	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	111	MET
2	B	175	LEU
2	B	189	LEU
2	B	197	GLN
2	B	242	GLN
2	B	245	LYS
1	C	808	CYS
1	C	821	LYS
1	C	848	GLU
1	C	911	GLU
1	C	928	LEU
1	C	930	GLU
1	C	945	THR
1	C	953	LYS
1	C	970	ILE
1	C	973	LYS
1	C	990	CYS
1	C	992	ASN
2	D	4	GLU
2	D	22	ARG
2	D	24	SER
2	D	41	ARG
2	D	85	GLN
2	D	101	ARG
2	D	111	MET
2	D	175	LEU
2	D	180	GLN
2	D	197	GLN
2	D	220	LEU
2	D	224	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	823	GLN
1	A	847	ASN
2	B	60	GLN
2	B	62	HIS
2	B	85	GLN
2	B	221	GLN
1	C	823	GLN
1	C	936	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	968	GLN
1	C	975	GLN
2	D	60	GLN
2	D	85	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	174/188 (92%)	0.10	2 (1%) 78 56	64, 76, 146, 153	0
1	C	174/188 (92%)	0.27	2 (1%) 78 56	63, 88, 166, 170	0
2	B	226/253 (89%)	0.28	2 (0%) 81 60	65, 98, 126, 131	0
2	D	226/253 (89%)	0.46	7 (3%) 51 29	92, 115, 142, 156	0
All	All	800/882 (90%)	0.29	13 (1%) 70 47	63, 100, 146, 170	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	248	ILE	4.1
2	D	240	PHE	2.8
1	A	973	LYS	2.6
2	D	225	PHE	2.4
1	C	970	ILE	2.3
2	D	123	SER	2.2
2	B	123	SER	2.1
2	B	225	PHE	2.1
2	D	48	LEU	2.1
1	C	982	LYS	2.1
2	D	179	GLN	2.1
1	A	970	ILE	2.0
2	D	15	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.