



wwPDB EM Validation Summary Report ⓘ

Mar 24, 2026 – 09:24 PM UTC

PDB ID : 9H9I / pdb_00009h9i
EMDB ID : EMD-51965
Title : Complex 2 (HEAD) 30S-IF1-IF3-tRNA-GE81112
Authors : Schedlbauer, A.; Han, X.; van Bakel, W.; Kaminishi, T.; Ochoa-Lizarralde, B.; Iturrioz, I.; Capuni, R.; Parry, R.; Zegarra, R.; Gil-Carton, D.; Lopez-Alonso, J.P.; Barragan Sanz, K.; Brandi, L.; Gualerzi, C.O.; Fucini, P.; Connell, S.R.
Deposited on : 2024-10-31
Resolution : 3.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	15020 (2.70 - 3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	1541	
2	2	4	
3	3	77	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	B	241	
5	C	233	
6	G	179	
7	I	130	
8	J	103	
9	M	118	
10	N	101	
11	S	92	

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 18959 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 16S RNA (head domain).

Mol	Chain	Residues	Atoms					AltConf	Trace
1	1	456	Total	C	N	O	P	0	0
			9758	4357	1772	3173	456		

- Molecule 2 is a RNA chain called mRNA (5'-R(P*AP*UP*GP*U)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
2	2	4	Total	C	N	O	P	0	0
			85	38	14	29	4		

- Molecule 3 is a RNA chain called t-RNA (5'-R(P*UP*CP*GP*GP*GP*CP*UP*CP*AP*U P*AP*AP*CP*CP*CP*GP*A)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
3	3	17	Total	C	N	O	P	0	0
			361	161	65	118	17		

- Molecule 4 is a protein called Small ribosomal subunit protein uS2.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	B	224	Total	C	N	O	S	0	0
			1753	1109	315	321	8		

- Molecule 5 is a protein called Small ribosomal subunit protein uS3.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	C	211	Total	C	N	O	S	0	0
			1653	1046	310	293	4		

- Molecule 6 is a protein called Small ribosomal subunit protein uS7.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	G	142	Total	C	N	O	S	0	0
			1116	694	216	203	3		

- Molecule 7 is a protein called Small ribosomal subunit protein uS9.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	I	127	Total	C	N	O	S	0	0
			1021	634	206	178	3		

- Molecule 8 is a protein called Small ribosomal subunit protein uS10.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	J	100	Total	C	N	O	S	0	0
			803	502	154	146	1		

- Molecule 9 is a protein called Small ribosomal subunit protein uS13.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	M	115	Total	C	N	O	S	0	0
			891	552	179	157	3		

- Molecule 10 is a protein called Small ribosomal subunit protein uS14.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	N	100	Total	C	N	O	S	0	0
			805	499	164	139	3		

- Molecule 11 is a protein called Small ribosomal subunit protein uS19.


Mol	Chain	Residues	Atoms					AltConf	Trace
11	S	83	Total	C	N	O	S	0	0
			663	424	126	111	2		

- Molecule 12 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
12	1	49	Total	Mg	0
			49	49	

- Molecule 13 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
13	B	1	Total	Zn	0
			1	1	




• Molecule 2: mRNA (5'-R(P*AP*UP*GP*U)-3')

Chain 2: 25% 75%




• Molecule 3: t-RNA (5'-R(P*UP*CP*GP*GP*GP*CP*UP*CP*AP*UP*AP*AP*CP*CP*CP*GP*A)-3')

Chain 3: 17% 5% 78%





• Molecule 4: Small ribosomal subunit protein uS2

Chain B: 10% 67% 26% 7%




• Molecule 5: Small ribosomal subunit protein uS3

Chain C: 




- Molecule 6: Small ribosomal subunit protein uS7

Chain G: 



- Molecule 7: Small ribosomal subunit protein uS9

Chain I: 



- Molecule 8: Small ribosomal subunit protein uS10

Chain J: 




- Molecule 9: Small ribosomal subunit protein uS13

Chain M: 



- Molecule 10: Small ribosomal subunit protein uS14

Chain N: 



- Molecule 11: Small ribosomal subunit protein uS19

Chain S: 67% 23% 10%

MET	P2	I11	K17	K18	V19	E20	V23	E24	T39	I40	F41	P42	N43	M44	L47	A50	V51	H52	Q56	H57	D64	E65	M66	H83	A84	ALA	ASP	LYS	LYS	ALA	LYS	LYS	LYS
-----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	34701	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	44.08	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	3250	Depositor
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.114	Depositor
Minimum map value	-0.027	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.017	Depositor
Map size (\AA)	427.39203, 427.39203, 427.39203	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.113, 1.113, 1.113	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5MC, 2MG, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	1	0.14	0/10842	0.25	0/16904
2	2	0.42	0/94	0.64	0/144
3	3	0.34	0/402	0.50	0/624
4	B	0.28	0/1784	0.60	2/2403 (0.1%)
5	C	0.15	0/1680	0.35	0/2263
6	G	0.13	0/1129	0.32	0/1514
7	I	0.15	0/1033	0.42	0/1375
8	J	0.43	0/813	0.62	0/1100
9	M	0.45	0/900	0.67	2/1204 (0.2%)
10	N	0.33	0/817	0.66	0/1088
11	S	0.16	0/680	0.40	0/915
All	All	0.21	0/20174	0.38	4/29534 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	M	17	ILE	N-CA-C	-7.02	103.46	110.62
9	M	6	GLY	CA-C-O	-6.33	117.87	122.23
4	B	72	THR	N-CA-C	-6.16	106.36	114.31
4	B	63	ARG	N-CA-C	-5.05	105.77	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	9758	0	4927	100	0
2	2	85	0	43	3	0
3	3	361	0	186	2	0
4	B	1753	0	1780	81	0
5	C	1653	0	1727	23	0
6	G	1116	0	1175	41	0
7	I	1021	0	1070	22	0
8	J	803	0	842	28	0
9	M	891	0	952	25	0
10	N	805	0	844	19	0
11	S	663	0	688	19	0
12	1	49	0	0	0	0
13	B	1	0	0	0	0
All	All	18959	0	14234	329	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 329 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:96:TRP:CH2	4:B:100:MET:SD	2.42	1.13
4:B:27:MET:HB3	4:B:189:THR:HB	1.37	1.04
4:B:96:TRP:CZ3	4:B:100:MET:SD	2.51	1.03
4:B:64:LYS:HG2	4:B:225:ARG:HG3	1.41	0.99
4:B:49:MET:HB2	4:B:201:PRO:HD3	1.40	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	B	222/241 (92%)	216 (97%)	6 (3%)	0	100	100
5	C	209/233 (90%)	205 (98%)	4 (2%)	0	100	100
6	G	140/179 (78%)	138 (99%)	2 (1%)	0	100	100
7	I	125/130 (96%)	120 (96%)	5 (4%)	0	100	100
8	J	98/103 (95%)	94 (96%)	4 (4%)	0	100	100
9	M	113/118 (96%)	106 (94%)	7 (6%)	0	100	100
10	N	98/101 (97%)	98 (100%)	0	0	100	100
11	S	81/92 (88%)	81 (100%)	0	0	100	100
All	All	1086/1197 (91%)	1058 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	B	186/199 (94%)	186 (100%)	0	100	100
5	C	172/190 (90%)	172 (100%)	0	100	100
6	G	119/147 (81%)	119 (100%)	0	100	100
7	I	105/107 (98%)	105 (100%)	0	100	100
8	J	88/90 (98%)	88 (100%)	0	100	100
9	M	93/96 (97%)	93 (100%)	0	100	100
10	N	83/84 (99%)	83 (100%)	0	100	100
11	S	72/79 (91%)	72 (100%)	0	100	100
All	All	918/992 (92%)	918 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
4	B	120	GLN
4	B	122	GLN
4	B	170	HIS
5	C	69	HIS
7	I	32	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	1	455/1541 (29%)	69 (15%)	7 (1%)
2	2	3/4 (75%)	1 (33%)	0
3	3	16/77 (20%)	1 (6%)	0
All	All	474/1622 (29%)	71 (14%)	7 (1%)

5 of 71 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	1	934	C
1	1	935	A
1	1	960	U
1	1	966	2MG
1	1	968	A

5 of 7 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	1	1196	A
1	1	1279	G
1	1	1335	U
1	1	1286	U
1	1	1183	U

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection.

RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	5MC	1	967	1	19,22,23	0.78	1 (5%)	26,32,35	0.49	0
1	2MG	1	966	1	23,26,27	0.38	0	33,38,41	0.42	0
1	2MG	1	1207	1	23,26,27	0.70	0	33,38,41	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	5MC	1	967	1	-	0/7/25/26	0/2/2/2
1	2MG	1	966	1	-	0/9/27/28	0/3/3/3
1	2MG	1	1207	1	-	0/9/27/28	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	1	967	5MC	C5-C4	-3.11	1.41	1.44

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	1	967	5MC	1	0
1	1	1207	2MG	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 50 ligands modelled in this entry, 50 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-51965. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 171




Y Index: 223




Z Index: 244


6.3.2 Raw map



X Index: 170



Y Index: 223




Z Index: 222


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.017. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 321 nm³; this corresponds to an approximate mass of 290 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.312 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.312 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.20	-	-
Author-provided FSC curve	3.19	4.44	3.31
Unmasked-calculated*	8.33	17.92	9.75

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 8.33 differs from the reported value 3.2 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-51965 and PDB model 9H9I. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.017 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.017).

9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 95% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.017) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.9540	<div></div> 0.4490
1	<div></div> 1.0000	<div></div> 0.4920
2	<div></div> 0.7650	<div></div> 0.2580
3	<div></div> 0.8560	<div></div> 0.4060
B	<div></div> 0.7490	<div></div> 0.2610
C	<div></div> 0.9490	<div></div> 0.4640
G	<div></div> 0.8380	<div></div> 0.3440
I	<div></div> 0.9650	<div></div> 0.4670
J	<div></div> 0.9740	<div></div> 0.4340
M	<div></div> 0.9790	<div></div> 0.4310
N	<div></div> 0.9790	<div></div> 0.4840
S	<div></div> 0.9910	<div></div> 0.4750

1.0

0.0

<0.0