



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 2, 2021 – 09:25 PM EDT

PDB ID : 3HB9
Title : Crystal Structure of *S. aureus* Pyruvate Carboxylase A610T Mutant
Authors : Tong, L.; Yu, L.P.C.
Deposited on : 2009-05-04
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1150	 4% 79% 17% ..
1	B	1150	 4% 75% 16% 7%
1	C	1150	 3% 74% 16% 7%
1	D	1150	 3% 70% 14% 14%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 33840 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate carboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1133	8942	5665	1505	1742	30	0	0	0
1	B	1074	8467	5366	1427	1645	29	0	0	0
1	C	1067	8443	5351	1421	1641	30	0	0	0
1	D	993	7870	4995	1326	1521	28	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	610	THR	ALA	engineered mutation	UNP Q99UY8
B	610	THR	ALA	engineered mutation	UNP Q99UY8
C	610	THR	ALA	engineered mutation	UNP Q99UY8
D	610	THR	ALA	engineered mutation	UNP Q99UY8

- Molecule 2 is 5-(HEXAHYDRO-2-OXO-1H-THIENO[3,4-D]IMIDAZOL-6-YL)PENTANAL (three-letter code: BTI) (formula: C₁₀H₁₆N₂O₂S).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mn	0	0
			1	1		
4	B	1	Total	Mn	0	0
			1	1		
4	C	1	Total	Mn	0	0
			1	1		
4	D	1	Total	Mn	0	0
			1	1		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyruvate carboxylase





- Molecule 1: Pyruvate carboxylase






• Molecule 1: Pyruvate carboxylase





● Molecule 1: Pyruvate carboxylase



ILE	
THR	
GLU	
A1142	
M1145	
GLU	
THR	
ILE	
GLN	
ALA	
PRO	
PHE	
ASP	
GLY	
VAL	
ILE	
LYS	
GLN	
VAL	
THR	
VAL	
ASN	
ASN	
GLY	
ASP	
THR	
ILE	
ALA	
THR	
GLY	
ASP	
LEU	
LEU	
ILE	
GLU	
ILE	
GLU	
LYS	
ALA	
THR	
ASP	

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	96.57Å 256.76Å 126.48Å 90.00° 109.65° 90.00°	Depositor
Resolution (Å)	30.00 – 2.90 29.58 – 2.82	Depositor EDS
% Data completeness (in resolution range)	92.8 (30.00-2.90) 90.1 (29.58-2.82)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 2.80Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.220 , 0.266 0.213 , 0.256	Depositor DCC
R_{free} test set	6262 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	51.8	Xtrriage
Anisotropy	0.306	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 30.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.035 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	33840	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, MN, BTI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/9110	0.50	0/12324
1	B	0.33	0/8627	0.49	0/11678
1	C	0.33	0/8605	0.49	0/11630
1	D	0.33	0/8021	0.50	0/10848
All	All	0.33	0/34363	0.49	0/46480

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8942	0	8872	118	0
1	B	8467	0	8413	115	0
1	C	8443	0	8352	118	0
1	D	7870	0	7808	102	0
2	A	15	0	15	1	0
2	B	15	0	15	1	0
2	C	30	0	31	1	0
3	A	27	0	12	0	0
3	C	27	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
All	All	33840	0	33530	444	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 444 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1024:ARG:HG3	1:B:1024:ARG:HH11	1.13	1.14
1:D:1085:ARG:HG2	1:D:1085:ARG:HH11	1.21	1.05
1:C:44:ASN:HD22	1:C:45:ARG:H	1.00	0.99
1:B:1000:GLY:H	1:B:1001:PRO:HD3	1.28	0.94
1:C:700:SER:H	1:C:736:HIS:HD2	1.16	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1127/1150 (98%)	1044 (93%)	74 (7%)	9 (1%)	19	51
1	B	1070/1150 (93%)	995 (93%)	60 (6%)	15 (1%)	11	36
1	C	1063/1150 (92%)	983 (92%)	69 (6%)	11 (1%)	15	45
1	D	987/1150 (86%)	928 (94%)	52 (5%)	7 (1%)	22	54
All	All	4247/4600 (92%)	3950 (93%)	255 (6%)	42 (1%)	15	45

5 of 42 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	386	ASP
1	A	1094	VAL
1	B	522	PRO
1	B	1001	PRO
1	B	1093	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	978/988 (99%)	911 (93%)	67 (7%)	15	41
1	B	929/988 (94%)	863 (93%)	66 (7%)	14	40
1	C	918/988 (93%)	855 (93%)	63 (7%)	15	41
1	D	860/988 (87%)	799 (93%)	61 (7%)	14	40
All	All	3685/3952 (93%)	3428 (93%)	257 (7%)	15	41

5 of 257 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	715	ARG
1	D	798	VAL
1	B	607	TRP
1	B	551	LYS
1	D	863	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 128 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	736	HIS
1	D	858	ASN
1	B	622	ASN
1	B	617	ASN
1	D	898	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BTI	A	2000	1	16,16,16	1.69	2 (12%)	21,21,21	2.22	4 (19%)
2	BTI	B	2000	1	16,16,16	1.69	2 (12%)	21,21,21	2.09	4 (19%)
3	ADP	C	2100	-	24,29,29	1.02	1 (4%)	29,45,45	1.34	4 (13%)
3	ADP	A	2100	-	24,29,29	1.01	1 (4%)	29,45,45	1.37	4 (13%)
2	BTI	C	1183	-	16,16,16	1.64	2 (12%)	21,21,21	2.31	5 (23%)
2	BTI	C	2000	1	16,16,16	1.66	2 (12%)	21,21,21	2.14	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BTI	A	2000	1	-	3/5/27/27	0/2/2/2
2	BTI	B	2000	1	-	3/5/27/27	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	C	2100	-	-	2/12/32/32	0/3/3/3
3	ADP	A	2100	-	-	2/12/32/32	0/3/3/3
2	BTI	C	1183	-	-	4/5/27/27	0/2/2/2
2	BTI	C	2000	1	-	4/5/27/27	0/2/2/2

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2000	BTI	O3-C3	4.83	1.33	1.23
2	B	2000	BTI	O3-C3	4.79	1.33	1.23
2	C	1183	BTI	O3-C3	4.73	1.33	1.23
2	C	2000	BTI	O3-C3	4.58	1.33	1.23
2	B	2000	BTI	C2-S1	-3.73	1.76	1.82

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2000	BTI	C2-C4-N2	-6.91	106.94	113.13
2	C	2000	BTI	C6-C5-N3	-6.29	105.04	113.03
2	C	1183	BTI	C2-C4-N2	-6.06	107.69	113.13
2	B	2000	BTI	C6-C5-N3	-5.91	105.52	113.03
2	C	2000	BTI	C2-C4-N2	-5.72	108.00	113.13

There are no chirality outliers.

5 of 18 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	2000	BTI	C11-C10-C9-C8
2	A	2000	BTI	S1-C2-C7-C8
2	A	2000	BTI	C4-C2-C7-C8
2	B	2000	BTI	S1-C2-C7-C8
2	B	2000	BTI	C4-C2-C7-C8


There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2000	BTI	1	0
2	B	2000	BTI	1	0
2	C	2000	BTI	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1133/1150 (98%)	-0.10	51 (4%) 33 29	19, 45, 109, 169	0
1	B	1074/1150 (93%)	0.08	46 (4%) 35 31	29, 60, 127, 178	0
1	C	1067/1150 (92%)	-0.02	37 (3%) 44 38	29, 61, 102, 147	0
1	D	993/1150 (86%)	-0.10	31 (3%) 49 44	17, 44, 138, 192	0
All	All	4267/4600 (92%)	-0.04	165 (3%) 39 35	17, 55, 121, 192	0

The worst 5 of 165 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	492	PRO	8.9
1	B	240	ASP	6.6
1	D	490	ILE	6.5
1	D	494	LEU	5.8
1	C	421	TYR	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ADP	A	2100	27/27	0.88	0.20	99,100,103,103	0
4	MN	A	2002	1/1	0.91	0.13	63,63,63,63	0
3	ADP	C	2100	27/27	0.93	0.17	78,78,78,79	0
4	MN	B	2002	1/1	0.94	0.13	77,77,77,77	0
2	BTI	C	2000	15/15	0.95	0.17	45,45,57,57	0
2	BTI	C	1183	15/15	0.95	0.16	48,48,49,49	0
2	BTI	B	2000	15/15	0.95	0.15	48,48,51,51	0
4	MN	C	2002	1/1	0.95	0.11	75,75,75,75	0
2	BTI	A	2000	15/15	0.96	0.15	50,51,53,53	0
4	MN	D	2002	1/1	0.97	0.20	57,57,57,57	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.