wwPDB X-ray Structure Validation Summary Report (i)

Aug 27, 2023 – 02:34 PM EDT

PDB ID	:	3HHG
Title	:	Structure of CrgA, a LysR-type transcriptional regulator from Neisseria meningitidis.
Authors	:	Sainsbury, S.; Ren, J.; Owens, R.J.; Stuart, D.I.; Oxford Protein Production Facility (OPPF)
Deposited on	:	2009-05-15
Resolution	:	3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	2%, 62%, 32%, • •
1	B	306	2%, 65%, 30%, • •
1	C	306	3%, 59%, 35%, • •
1	D	306	4%, 61%, 33%, • •
1	E	306	6%, 66%, 29%, • •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	306	2% 61% 33%
1	G	306	8% 66% 28%
1	H	306	6% 64% 31%

2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 18304 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator, LysR family.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	293	2278	1436	399	434	9	0	0	0
1	B	295	2294	1444	403	438	9	0	0	0
1	C	294	2286	1440	401	436	9	0	0	0
1	D	294	2286	1440	401	436	9	0	0	0
1	E	297	2310	1455	405	440	10	0	0	0
1	F	293	2278	1436	399	434	9	0	0	0
1	G	294	2286	1441	400	435	10	0	0	0
1	H	294	2286	1441	400	435	10	0	0	0

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	300	LYS	-	expression tag	UNP Q9JXW7
A	301	HIS	-	expression tag	UNP Q9JXW7
A	302	HIS	-	expression tag	UNP Q9JXW7
A	303	HIS	-	expression tag	UNP Q9JXW7
A	304	HIS	-	expression tag	UNP Q9JXW7
A	305	HIS	-	expression tag	UNP Q9JXW7
A	306	HIS	-	expression tag	UNP Q9JXW7
B	300	LYS	-	expression tag	UNP Q9JXW7
B	301	HIS	-	expression tag	UNP Q9JXW7
B	302	HIS	-	expression tag	UNP Q9JXW7
B	303	HIS	-	expression tag	UNP Q9JXW7
B	304	HIS	-	expression tag	UNP Q9JXW7
B	305	HIS	-	expression tag	UNP Q9JXW7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	306	HIS	-	expression tag	UNP Q9JXW7
C	300	LYS	-	expression tag	UNP Q9JXW7
C	301	HIS	-	expression tag	UNP Q9JXW7
C	302	HIS	-	expression tag	UNP Q9JXW7
C	303	HIS	-	expression tag	UNP Q9JXW7
C	304	HIS	-	expression tag	UNP Q9JXW7
C	305	HIS	-	expression tag	UNP Q9JXW7
C	306	HIS	-	expression tag	UNP Q9JXW7
D	300	LYS	-	expression tag	UNP Q9JXW7
D	301	HIS	-	expression tag	UNP Q9JXW7
D	302	HIS	-	expression tag	UNP Q9JXW7
D	303	HIS	-	expression tag	UNP Q9JXW7
D	304	HIS	-	expression tag	UNP Q9JXW7
D	305	HIS	-	expression tag	UNP Q9JXW7
D	306	HIS	-	expression tag	UNP Q9JXW7
E	300	LYS	-	expression tag	UNP Q9JXW7
E	301	HIS	-	expression tag	UNP Q9JXW7
E	302	HIS	-	expression tag	UNP Q9JXW7
E	303	HIS	-	expression tag	UNP Q9JXW7
E	304	HIS	-	expression tag	UNP Q9JXW7
E	305	HIS	-	expression tag	UNP Q9JXW7
E	306	HIS	-	expression tag	UNP Q9JXW7
F	300	LYS	-	expression tag	UNP Q9JXW7
F	301	HIS	-	expression tag	UNP Q9JXW7
F	302	HIS	-	expression tag	UNP Q9JXW7
F	303	HIS	-	expression tag	UNP Q9JXW7
F	304	HIS	-	expression tag	UNP Q9JXW7
F	305	HIS	-	expression tag	UNP Q9JXW7
F	306	HIS	-	expression tag	UNP Q9JXW7
G	300	LYS	-	expression tag	UNP Q9JXW7
G	301	HIS	-	expression tag	UNP Q9JXW7
G	302	HIS	-	expression tag	UNP Q9JXW7
G	303	HIS	-	expression tag	UNP Q9JXW7
G	304	HIS	-	expression tag	UNP Q9JXW7
G	305	HIS	-	expression tag	UNP Q9JXW7
G	306	HIS	-	expression tag	UNP Q9JXW7
H	300	LYS	-	expression tag	UNP Q9JXW7
H	301	HIS	-	expression tag	UNP Q9JXW7
H	302	HIS	-	expression tag	UNP Q9JXW7
H	303	HIS	-	expression tag	UNP Q9JXW7
H	304	HIS	-	expression tag	UNP Q9JXW7
H	305	HIS	-	expression tag	UNP Q9JXW7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	306	HIS	-	expression tag	UNP Q9JXW7

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Transcriptional regulator, LysR family

Chain A:



• Molecule 1: Transcriptional regulator, LysR family

Chain B:



• Molecule 1: Transcriptional regulator, LysR family

Chain C:





- Molecule 1: Transcriptional regulator, LysR family

Chain D:



- Molecule 1: Transcriptional regulator, LysR family

Chain E:



- Molecule 1: Transcriptional regulator, LysR family

Chain F:





• Molecule 1: Transcriptional regulator, LysR family

Chain G:



• Molecule 1: Transcriptional regulator, LysR family

Chain H:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	104.64Å 119.09Å 250.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.88 – 3.20 29.81 – 3.20	Depositor EDS
% Data completeness (in resolution range)	86.5 (29.88-3.20) 81.7 (29.81-3.20)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.88 (at 3.18Å)	Xtriage
Refinement program	REFMAC 5.5.0047	Depositor
R, R_{free}	0.214 , 0.286 0.228 , 0.295	Depositor DCC
R_{free} test set	2165 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å^2)	93.8	Xtriage
Anisotropy	0.278	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.23 , 45.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	18304	wwPDB-VP
Average B, all atoms (Å^2)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 35.28 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.9808e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.34	0/2317	0.52	0/3137
1	B	0.35	0/2333	0.52	0/3159
1	C	0.34	0/2325	0.51	0/3148
1	D	0.33	0/2325	0.52	0/3148
1	E	0.35	0/2349	0.52	0/3180
1	F	0.34	0/2317	0.51	0/3137
1	G	0.32	0/2325	0.48	0/3147
1	H	0.34	0/2325	0.47	0/3147
All	All	0.34	0/18616	0.51	0/25203

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2278	0	2286	92	0
1	B	2294	0	2298	91	0
1	C	2286	0	2292	118	0
1	D	2286	0	2292	103	0
1	E	2310	0	2321	92	0
1	F	2278	0	2286	95	0
1	G	2286	0	2298	94	0
1	H	2286	0	2298	105	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	18304	0	18371	715	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 715 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:9:THR:HG22	1:G:74:LEU:HD21	1.19	1.12
1:H:168:ALA:HB2	1:H:236:ILE:HG22	1.30	1.10
1:E:167:ILE:HD13	1:E:253:LEU:HD13	1.32	1.10
1:F:82:THR:HG21	1:F:287:ASP:OD2	1.56	1.05
1:G:168:ALA:HB2	1:G:236:ILE:HG22	1.42	1.01

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	291/306 (95%)	271 (93%)	19 (6%)	1 (0%)	41	74
1	B	293/306 (96%)	275 (94%)	17 (6%)	1 (0%)	41	74
1	C	292/306 (95%)	279 (96%)	12 (4%)	1 (0%)	41	74
1	D	292/306 (95%)	269 (92%)	23 (8%)	0	100	100
1	E	295/306 (96%)	280 (95%)	15 (5%)	0	100	100
1	F	291/306 (95%)	277 (95%)	13 (4%)	1 (0%)	41	74
1	G	292/306 (95%)	273 (94%)	18 (6%)	1 (0%)	41	74
1	H	292/306 (95%)	268 (92%)	23 (8%)	1 (0%)	41	74
All	All	2338/2448 (96%)	2192 (94%)	140 (6%)	6 (0%)	41	74

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	20	PHE
1	C	197	GLY
1	F	196	PRO
1	B	196	PRO
1	A	196	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/260 (95%)	237 (96%)	11 (4%)	28	64
1	B	250/260 (96%)	237 (95%)	13 (5%)	23	59
1	C	249/260 (96%)	238 (96%)	11 (4%)	28	64
1	D	249/260 (96%)	236 (95%)	13 (5%)	23	59
1	E	252/260 (97%)	243 (96%)	9 (4%)	35	69
1	F	248/260 (95%)	239 (96%)	9 (4%)	35	69
1	G	249/260 (96%)	240 (96%)	9 (4%)	35	69
1	H	249/260 (96%)	242 (97%)	7 (3%)	43	74
All	All	1994/2080 (96%)	1912 (96%)	82 (4%)	30	66

5 of 82 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	19	SER
1	G	134	ASN
1	F	57	LEU
1	F	247	ASP
1	H	9	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	26	GLN
1	H	92	GLN
1	C	180	GLN
1	C	88	HIS
1	H	116	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	293/306 (95%)	-0.20	6 (2%)	65	51	43, 71, 123, 139	0
1	B	295/306 (96%)	-0.24	5 (1%)	70	57	41, 72, 126, 135	0
1	C	294/306 (96%)	-0.12	10 (3%)	45	29	49, 78, 124, 140	0
1	D	294/306 (96%)	0.00	13 (4%)	34	21	50, 78, 129, 138	0
1	E	297/306 (97%)	-0.02	19 (6%)	19	11	38, 76, 131, 145	0
1	F	293/306 (95%)	-0.14	5 (1%)	70	57	48, 76, 126, 140	0
1	G	294/306 (96%)	0.32	24 (8%)	11	6	54, 81, 129, 137	0
1	H	294/306 (96%)	0.17	19 (6%)	18	11	56, 80, 128, 136	0
All	All	2354/2448 (96%)	-0.03	101 (4%)	35	22	38, 77, 128, 145	0

The worst 5 of 101 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	35	SER	5.6
1	G	150	ASP	5.2
1	E	38	VAL	4.9
1	D	30	ALA	4.7
1	E	37	ILE	4.5

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.