wwPDB X-ray Structure Validation Summary Report (i)

Nov 1, 2023 – 01:10 PM JST

PDB ID	:	5HPY
Title	:	Crystal Structure of RhoA.GDP.MgF3-in complex with human Myosin 9b RhoGAP domain
Authors	:	Yi, F.S.; Ren, J.Q.; Feng, W.
Deposited on	:	2016-01-21
Resolution	:	2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	9%, 70%, 11%, •, 17%
1	D	232	11%, 68%, 14%, 18%
2	B	185	2%, 79%, 17%, •, •
2	F	185	2%, 82%, 14%, •, •

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5968 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Unconventional myosin-IXb.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	192	1501	966	257	269	9	0	0	0
1	D	190	1496	964	254	267	11	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	185	GLY	-	expression tag	UNP Q13459
A	186	PRO	-	expression tag	UNP Q13459
A	187	GLY	-	expression tag	UNP Q13459
A	188	SER	-	expression tag	UNP Q13459
A	189	GLU	-	expression tag	UNP Q13459
A	190	PHE	-	expression tag	UNP Q13459
D	185	GLY	-	expression tag	UNP Q13459
D	186	PRO	-	expression tag	UNP Q13459
D	187	GLY	-	expression tag	UNP Q13459
D	188	SER	-	expression tag	UNP Q13459
D	189	GLU	-	expression tag	UNP Q13459
D	190	PHE	-	expression tag	UNP Q13459

- Molecule 2 is a protein called Transforming protein RhoA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	178	1398	884	234	270	10	0	1	0
2	F	178	1370	866	229	265	10	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P61586
B	-2	PRO	-	expression tag	UNP P61586
B	-1	GLY	-	expression tag	UNP P61586
B	0	SER	-	expression tag	UNP P61586
B	1	GLU	-	expression tag	UNP P61586
B	2	PHE	-	expression tag	UNP P61586
B	25	ASN	PHE	engineered mutation	UNP P61586
F	-3	GLY	-	expression tag	UNP P61586
F	-2	PRO	-	expression tag	UNP P61586
F	-1	GLY	-	expression tag	UNP P61586
F	0	SER	-	expression tag	UNP P61586
F	1	GLU	-	expression tag	UNP P61586
F	2	PHE	-	expression tag	UNP P61586
F	25	ASN	PHE	engineered mutation	UNP P61586

- Molecule 3 is TRIFLUOROMAGNESATE (three-letter code: MGF) (formula: F_3Mg).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	F	Mg		
3	B	1	4	3	1	0	0
			Total	F	Mg		
3	F	1	4	3	1	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
4	B	1	1	1	0	0
			Total	Mg		
4	F	2	2	2	0	0

- Molecule 5 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	B	1	28	10	5	11	2	0	0
			Total	C	N	O	P		
5	F	1	28	10	5	11	2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	28	28	28	0	0
			Total	O		
6	B	45	45	45	0	0
			Total	O		
6	D	20	20	20	0	0
			Total	O		
6	F	43	43	43	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Unconventional myosin-IXb

Chain A:



• Molecule 1: Unconventional myosin-IXb

Chain D:



• Molecule 2: Transforming protein RhoA

Chain B:



• Molecule 2: Transforming protein RhoA

Chain F:


GLY
PRO
GLY
SER
GLU
PHE
A3
I4
V24
D28
E47
Q52
P71
L72
S73
Y74
S85
I86
D87
E93
P96
T100
P101
N123
D124
E125
A132
K133
M134
K140
P141
E142
R145
M157
K162
T163
K164
D165
R168
E169
E172
A177

Q180
ALA

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α, β, γ	114.06Å 90.76Å 93.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.70 – 2.40 48.70 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.70-2.40) 99.8 (48.70-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.83 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.211 , 0.246 0.210 , 0.245	Depositor DCC
R_{free} test set	1869 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å^2)	51.5	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 57.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	5968	wwPDB-VP
Average B, all atoms (Å^2)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.37 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7810e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GDP, MGF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.42	0/1527	0.57	0/2067
1	D	0.41	0/1523	0.55	0/2061
2	B	0.50	0/1428	0.61	0/1936
2	F	0.48	0/1397	0.59	0/1898
All	All	0.45	0/5875	0.58	0/7962

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1501	0	1526	20	0
1	D	1496	0	1535	18	0
2	B	1398	0	1374	20	0
2	F	1370	0	1325	18	0
3	B	4	0	0	0	0
3	F	4	0	0	0	0
4	B	1	0	0	0	0
4	F	2	0	0	0	0
5	B	28	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	F	28	0	12	0	0
6	A	28	0	0	1	0
6	B	45	0	0	0	0
6	D	20	0	0	1	0
6	F	43	0	0	3	0
All	All	5968	0	5784	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 73 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:351:ALA:HB1	1:A:373:ILE:HG13	1.56	0.87
1:D:351:ALA:HB1	1:D:373:ILE:HG13	1.54	0.86
2:B:118:LYS:HB3	2:B:121:LEU:HD12	1.76	0.67
1:D:284:THR:N	6:D:501:HOH:O	2.29	0.66
1:D:383:GLU:OE1	1:D:386:ARG:NH1	2.28	0.65

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/232 (80%)	179 (96%)	6 (3%)	1 (0%)	29	41
1	D	184/232 (79%)	179 (97%)	4 (2%)	1 (0%)	29	41
2	B	177/185 (96%)	173 (98%)	4 (2%)	0	100	100
2	F	176/185 (95%)	172 (98%)	4 (2%)	0	100	100
All	All	723/834 (87%)	703 (97%)	18 (2%)	2 (0%)	41	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	197	HIS
1	D	298	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/200 (80%)	154 (97%)	5 (3%)	40	60
1	D	162/200 (81%)	157 (97%)	5 (3%)	40	60
2	B	153/160 (96%)	151 (99%)	2 (1%)	69	84
2	F	147/160 (92%)	145 (99%)	2 (1%)	67	82
All	All	621/720 (86%)	607 (98%)	14 (2%)	50	70

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	212	VAL
1	D	283	MET
2	F	145	ARG
1	D	371	LEU
2	F	142	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	GDP	B	203	4,3	24,30,30	1.15	1 (4%)	30,47,47	1.17	3 (10%)
3	MGF	B	201	6,5	0,3,3	-	-	-		
5	GDP	F	202	4,3	24,30,30	0.96	1 (4%)	30,47,47	1.29	4 (13%)
3	MGF	F	201	6,5	0,3,3	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GDP	B	203	4,3	-	2/12/32/32	0/3/3/3
5	GDP	F	202	4,3	-	1/12/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	203	GDP	C6-N1	-2.81	1.33	1.37
5	F	202	GDP	C6-N1	-2.15	1.34	1.37

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	F	202	GDP	PA-O3A-PB	-3.65	120.31	132.83
5	B	203	GDP	O3B-PB-O3A	3.12	115.10	104.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	F	202	GDP	C3'-C2'-C1'	2.90	105.35	100.98
5	B	203	GDP	PA-O3A-PB	-2.47	124.34	132.83
5	B	203	GDP	C8-N7-C5	2.25	107.27	102.99

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	203	GDP	PA-O3A-PB-O3B
5	B	203	GDP	PA-O3A-PB-O1B
5	F	202	GDP	PB-O3A-PA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	192/232 (82%)	0.56	21 (10%)	5	5	43, 74, 97, 109	0
1	D	190/232 (81%)	0.71	26 (13%)	3	2	44, 76, 100, 122	0
2	B	178/185 (96%)	0.32	4 (2%)	62	60	34, 59, 89, 104	0
2	F	178/185 (96%)	0.43	4 (2%)	62	60	39, 62, 100, 111	0
All	All	738/834 (88%)	0.51	55 (7%)	14	13	34, 67, 97, 122	0

The worst 5 of 55 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	252	THR	4.8
1	D	199	GLY	4.6
1	D	313	HIS	4.4
1	A	361	SER	4.2
1	A	252	THR	4.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	MG	F	204	1/1	0.95	0.13	49,49,49,49	0
5	GDP	F	202	28/28	0.95	0.18	39,56,62,68	0
4	MG	B	202	1/1	0.96	0.14	37,37,37,37	0
4	MG	F	203	1/1	0.96	0.09	42,42,42,42	0
5	GDP	B	203	28/28	0.98	0.18	34,50,58,67	0
3	MGF	F	201	4/4	0.98	0.16	36,40,41,47	0
3	MGF	B	201	4/4	0.99	0.13	37,38,39,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GDP F 202:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)







Electron density around GDP B 203:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.