



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 05:46 AM EDT

PDB ID : 9HQ1 / pdb\_00009hq1  
Title : XusB lipoprotein bound to ferric salmochelin  
Authors : Silale, A.; Soo, Y.L.; van den Berg, B.  
Deposited on : 2024-12-16  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	472	
1	B	472	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7471 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DUF4374 domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	426	Total	C	N	O	S	0	1	0
			3307	2090	547	660	10			
1	B	426	Total	C	N	O	S	0	0	0
			3298	2084	545	659	10			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	465	LEU	-	expression tag	UNP Q8A622
A	466	GLU	-	expression tag	UNP Q8A622
A	467	HIS	-	expression tag	UNP Q8A622
A	468	HIS	-	expression tag	UNP Q8A622
A	469	HIS	-	expression tag	UNP Q8A622
A	470	HIS	-	expression tag	UNP Q8A622
A	471	HIS	-	expression tag	UNP Q8A622
A	472	HIS	-	expression tag	UNP Q8A622
B	465	LEU	-	expression tag	UNP Q8A622
B	466	GLU	-	expression tag	UNP Q8A622
B	467	HIS	-	expression tag	UNP Q8A622
B	468	HIS	-	expression tag	UNP Q8A622
B	469	HIS	-	expression tag	UNP Q8A622
B	470	HIS	-	expression tag	UNP Q8A622
B	471	HIS	-	expression tag	UNP Q8A622
B	472	HIS	-	expression tag	UNP Q8A622

- Molecule 2 is Salmochelin S4 (CCD ID: A1IWM) (formula: C<sub>42</sub>H<sub>47</sub>N<sub>3</sub>O<sub>25</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			70	42	3	25		

- Molecule 3 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe	0	0
			1	1		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Ca	0	0
			2	2		
4	B	4	Total	Ca	0	0
			4	4		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Cl	0	0
			2	2		
5	B	2	Total	Cl	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	384	Total 384	O 384	0	0
6	B	401	Total 401	O 401	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DUF4374 domain-containing protein



- Molecule 1: DUF4374 domain-containing protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.44Å 108.72Å 93.83Å 90.00° 96.66° 90.00°	Depositor
Resolution (Å)	54.36 – 1.80 54.36 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (54.36-1.80) 99.7 (54.36-1.80)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.66 (at 1.80Å)	Xtriage
Refinement program	PHENIX 1.21.1_5286	Depositor
R, $R_{free}$	0.193 , 0.229 0.193 , 0.227	Depositor DCC
$R_{free}$ test set	4149 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.3	Xtriage
Anisotropy	0.213	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 32.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7471	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, CA, FE, A1IWM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/3384	0.48	0/4603
1	B	0.28	0/3373	0.47	0/4591
All	All	0.28	0/6757	0.48	0/9194

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3307	0	3165	21	0
1	B	3298	0	3153	12	0
2	A	70	0	0	0	0
3	A	1	0	0	0	0
4	A	2	0	0	0	0
4	B	4	0	0	0	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	384	0	0	3	0
6	B	401	0	0	2	0
All	All	7471	0	6318	32	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:35:THR:HG23	1:B:436:GLU:HG3	1.82	0.62
1:A:171:TYR:CD2	1:A:234[B]:ARG:HG2	2.36	0.60
1:A:77:ILE:HB	1:A:446:GLY:HA3	1.85	0.58
1:B:410:GLN:O	1:B:417:GLY:HA2	2.05	0.56
1:B:396:THR:HG21	1:B:411:TYR:HB3	1.88	0.55
1:B:261:ARG:NH2	6:B:611:HOH:O	2.35	0.53
1:A:69:GLY:HA3	1:A:105:TRP:CG	2.48	0.49
1:B:69:GLY:HA3	1:B:105:TRP:CG	2.50	0.47
1:A:234[B]:ARG:HD3	1:A:236:TRP:CZ2	2.51	0.46
1:B:238:ILE:HD11	1:B:244:LEU:HB2	1.97	0.45
1:A:346:ARG:NH2	1:A:403:ASP:OD2	2.49	0.45
1:A:83:LYS:HD3	1:A:144:ASN:HA	1.97	0.45
1:B:138:ALA:HB2	1:B:160:SER:O	2.16	0.44
1:A:222:ASN:ND2	1:A:225:GLU:HG3	2.33	0.44
1:B:286:ASP:HB2	6:B:899:HOH:O	2.17	0.44
1:A:39:GLN:HB3	1:A:63:GLU:HG3	2.00	0.43
1:A:284:GLU:H	1:A:284:GLU:CD	2.26	0.43
1:A:408:ARG:O	1:A:419:THR:HA	2.19	0.43
1:B:174:ILE:HG22	1:B:184:VAL:HG13	2.01	0.42
1:B:183:TYR:CE1	1:B:267:ALA:HB1	2.54	0.42
1:A:233:LEU:HG	1:A:234[B]:ARG:HG3	2.02	0.42
1:B:77:ILE:HB	1:B:446:GLY:HA3	2.02	0.42
1:A:143:TYR:CE1	1:A:151:PRO:HB3	2.55	0.42
1:A:143:TYR:CD1	1:A:148:PHE:HA	2.56	0.41
1:A:441:ARG:HA	6:A:881:HOH:O	2.20	0.41
1:A:287:ARG:NH2	6:A:630:HOH:O	2.53	0.41
1:A:398:TRP:CZ3	1:A:408:ARG:HB2	2.56	0.41
1:A:183:TYR:CE1	1:A:208:ARG:HD3	2.56	0.41
1:A:79:GLU:OE1	6:A:601:HOH:O	2.22	0.40
1:A:143:TYR:CZ	1:A:151:PRO:HB3	2.57	0.40
1:A:114:THR:O	1:A:126:LYS:NZ	2.49	0.40
1:B:411:TYR:CE1	1:B:441:ARG:HG3	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	421/472 (89%)	407 (97%)	14 (3%)	0	100	100
1	B	422/472 (89%)	407 (96%)	14 (3%)	1 (0%)	43	31
All	All	843/944 (89%)	814 (97%)	28 (3%)	1 (0%)	48	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	388	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	352/389 (90%)	351 (100%)	1 (0%)	86	86
1	B	351/389 (90%)	349 (99%)	2 (1%)	78	77
All	All	703/778 (90%)	700 (100%)	3 (0%)	84	83

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	444	THR
1	B	119	SER
1	B	415	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	GLN
1	A	56	ASN
1	A	316	ASN
1	A	328	HIS
1	A	337	ASN
1	A	410	GLN
1	A	412	ASN
1	B	81	ASN
1	B	337	ASN
1	B	352	HIS
1	B	413	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 11 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	A1IWM	A	501	3	75,75,75	2.05	22 (29%)	110,110,110	1.92	33 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1IWM	A	501	3	-	2/63/103/103	0/5/6/6

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	A1IWM	C26-C25	5.06	1.65	1.53
2	A	501	A1IWM	C17-N16	4.87	1.45	1.34
2	A	501	A1IWM	O52-C47	-4.36	1.38	1.43
2	A	501	A1IWM	O05-C03	4.24	1.42	1.33
2	A	501	A1IWM	C01-C02	4.21	1.64	1.52
2	A	501	A1IWM	O15-C13	3.95	1.41	1.33
2	A	501	A1IWM	C61-N60	3.82	1.43	1.34
2	A	501	A1IWM	O10-C08	3.61	1.40	1.33
2	A	501	A1IWM	C19-C17	3.55	1.57	1.50
2	A	501	A1IWM	C24-C23	3.41	1.44	1.39
2	A	501	A1IWM	C39-N38	3.27	1.41	1.34
2	A	501	A1IWM	O52-C51	-2.90	1.37	1.44
2	A	501	A1IWM	C50-C49	2.85	1.59	1.52
2	A	501	A1IWM	C65-C64	2.77	1.43	1.40
2	A	501	A1IWM	C02-N60	2.55	1.51	1.45
2	A	501	A1IWM	C53-C51	2.48	1.60	1.51
2	A	501	A1IWM	C49-C48	2.30	1.58	1.52
2	A	501	A1IWM	C19-C20	2.24	1.45	1.41
2	A	501	A1IWM	C50-C51	2.23	1.57	1.53
2	A	501	A1IWM	C06-C07	2.21	1.58	1.52
2	A	501	A1IWM	C12-N16	2.14	1.50	1.45
2	A	501	A1IWM	C23-C25	2.04	1.54	1.51

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	A1IWM	C23-C25-C26	6.47	127.81	112.93
2	A	501	A1IWM	O52-C47-C45	5.87	117.96	107.88
2	A	501	A1IWM	O30-C29-C31	5.01	118.86	106.44
2	A	501	A1IWM	O10-C08-O09	-4.29	116.28	124.14
2	A	501	A1IWM	O30-C25-C23	-4.16	100.74	107.88
2	A	501	A1IWM	C51-O52-C47	3.36	119.02	112.53
2	A	501	A1IWM	C46-C41-C42	-3.31	115.53	119.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	A1IWM	C41-C42-C43	3.25	121.97	119.99
2	A	501	A1IWM	O10-C08-C07	3.21	119.13	111.58
2	A	501	A1IWM	O62-C61-N60	-3.19	116.40	122.47
2	A	501	A1IWM	C27-C26-C25	3.07	115.91	109.41
2	A	501	A1IWM	C41-C39-N38	2.90	122.96	116.67
2	A	501	A1IWM	O18-C17-N16	-2.87	117.01	122.47
2	A	501	A1IWM	C24-C19-C20	-2.86	116.05	119.39
2	A	501	A1IWM	C07-N38-C39	2.83	128.36	121.56
2	A	501	A1IWM	O40-C39-N38	-2.76	117.22	122.47
2	A	501	A1IWM	C63-C61-N60	2.73	122.60	116.67
2	A	501	A1IWM	C50-C49-C48	2.65	115.47	110.83
2	A	501	A1IWM	C06-C07-N38	2.63	117.12	111.31
2	A	501	A1IWM	C45-C47-C48	-2.61	106.93	112.93
2	A	501	A1IWM	O52-C51-C53	2.51	112.66	106.44
2	A	501	A1IWM	C11-C12-N16	2.42	116.67	111.31
2	A	501	A1IWM	O10-C11-C12	2.37	114.92	108.26
2	A	501	A1IWM	O35-C26-C27	-2.36	104.81	110.38
2	A	501	A1IWM	C27-C28-C29	-2.30	106.07	110.23
2	A	501	A1IWM	C44-C43-C42	-2.25	118.89	120.45
2	A	501	A1IWM	C24-C19-C17	2.23	123.53	117.21
2	A	501	A1IWM	C19-C17-N16	2.21	121.47	116.67
2	A	501	A1IWM	C41-C46-C45	2.16	124.16	120.92
2	A	501	A1IWM	C06-C07-C08	-2.14	104.47	109.88
2	A	501	A1IWM	C46-C41-C39	2.03	122.97	117.21
2	A	501	A1IWM	C49-C50-C51	2.03	113.91	110.23
2	A	501	A1IWM	O59-C42-C41	-2.02	117.51	121.11

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	A1IWM	C28-C29-C31-O32
2	A	501	A1IWM	O30-C29-C31-O32

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	426/472 (90%)	0.04	11 (2%) 57 57	11, 24, 40, 53	1 (0%)
1	B	426/472 (90%)	0.06	22 (5%) 33 31	16, 23, 44, 63	0
All	All	852/944 (90%)	0.05	33 (3%) 43 43	11, 24, 42, 63	1 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	418	VAL	6.1
1	B	417	GLY	4.2
1	B	415	ASN	3.7
1	B	411	TYR	3.4
1	A	388	ASN	3.3
1	B	310	THR	3.3
1	B	315	THR	3.3
1	A	418	VAL	3.2
1	B	412	ASN	3.2
1	B	118	GLY	3.2
1	B	414	GLY	3.0
1	B	169	GLN	3.0
1	A	437	TYR	3.0
1	A	427	ASN	2.9
1	A	293	GLY	2.8
1	B	119	SER	2.7
1	A	310	THR	2.7
1	A	435	ASN	2.6
1	B	416	GLU	2.5
1	A	315	THR	2.5
1	B	440	THR	2.5
1	B	388	ASN	2.5
1	B	168	SER	2.4
1	B	167	ARG	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	413	GLN	2.4
1	B	396	THR	2.3
1	A	316	ASN	2.3
1	B	120	ASN	2.3
1	B	316	ASN	2.3
1	B	410	GLN	2.3
1	A	438	THR	2.2
1	A	426	THR	2.1
1	B	122	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	A1IWM	A	501	70/70	0.86	0.11	26,35,40,42	70
5	CL	A	505	1/1	0.93	0.10	40,40,40,40	0
4	CA	B	502	1/1	0.95	0.11	42,42,42,42	0
4	CA	B	504	1/1	0.96	0.11	49,49,49,49	0
4	CA	A	504	1/1	0.97	0.12	43,43,43,43	0
5	CL	B	506	1/1	0.97	0.07	47,47,47,47	0
4	CA	B	503	1/1	0.98	0.16	41,41,41,41	0
4	CA	B	501	1/1	0.98	0.07	27,27,27,27	0
4	CA	A	503	1/1	0.99	0.03	24,24,24,24	0
5	CL	A	506	1/1	0.99	0.03	25,25,25,25	0
5	CL	B	505	1/1	0.99	0.04	26,26,26,26	0
3	FE	A	502	1/1	0.99	0.03	27,27,27,27	0




The following is a graphical depiction of the model fit to experimental electron density of all



instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around A1IWM A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.