



wwPDB EM Validation Summary Report ⓘ

Mar 14, 2026 – 07:49 AM UTC

PDB ID : 9HY6 / pdb_00009hy6
EMDB ID : EMD-52500
Title : Structure of E.coli ribosome with nascent chain at linker length of 31 amino acids, with mRNA, P-site and A-site tRNAs
Authors : Mitropoulou, A.; Wlodarski, T.; Plessa, E.; Cabrita, L.D.; Christodoulou, J.
Deposited on : 2025-01-09
Resolution : 2.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



























Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	8728 (2.10 - 3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	77	<div><div></div><div>27%57%14%</div></div>
1	v	77	<div><div></div><div>35%49%14%</div></div>
2	X	9	<div><div></div><div>11%33%56%</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	z	148	
4	0	78	
5	1	63	
6	2	59	
7	3	57	
8	4	55	
9	6	46	
10	7	65	
11	8	50	
12	a	120	
13	b	2904	
14	c	273	
15	d	209	
16	e	201	
17	f	179	
18	g	177	
19	h	149	
20	j	142	
21	k	123	
22	l	144	
23	m	136	
24	n	127	
25	o	117	
26	p	115	
27	q	118	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
28	r	103	
29	s	110	
30	t	100	
31	u	104	
32	w	94	
33	y	85	

2 Entry composition [i](#)

There are 33 unique types of molecules in this entry. The entry contains 92670 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called pro-tRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	v	76	Total	C	N	O	P	0	0
			1626	724	292	534	76		
1	A	76	Total	C	N	O	P	0	0
			1626	724	292	534	76		

- Molecule 2 is a RNA chain called mRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	X	9	Total	C	N	O	P	0	0
			189	84	32	64	9		

- Molecule 3 is a protein called Gelation factor.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	z	20	Total	C	N	O	0	0
			181	126	30	25		

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
z	1	MET	-	initiating methionine	UNP P13466
z	2	HIS	-	expression tag	UNP P13466
z	3	HIS	-	expression tag	UNP P13466
z	4	HIS	-	expression tag	UNP P13466
z	5	HIS	-	expression tag	UNP P13466
z	6	HIS	-	expression tag	UNP P13466
z	7	HIS	-	expression tag	UNP P13466
z	8	ALA	-	expression tag	UNP P13466
z	9	SER	-	expression tag	UNP P13466
z	130	GLU	-	expression tag	UNP P13466
z	131	LEU	-	expression tag	UNP P13466
z	132	PHE	-	expression tag	UNP P13466
z	133	SER	-	expression tag	UNP P13466

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
z	134	THR	-	expression tag	UNP P13466
z	135	PRO	-	expression tag	UNP P13466
z	136	VAL	-	expression tag	UNP P13466
z	137	TRP	-	expression tag	UNP P13466
z	138	ILE	-	expression tag	UNP P13466
z	139	TRP	-	expression tag	UNP P13466
z	140	TRP	-	expression tag	UNP P13466
z	141	TRP	-	expression tag	UNP P13466
z	142	PRO	-	expression tag	UNP P13466
z	143	ARG	-	expression tag	UNP P13466
z	144	ILE	-	expression tag	UNP P13466
z	145	ARG	-	expression tag	UNP P13466
z	146	GLY	-	expression tag	UNP P13466
z	147	PRO	-	expression tag	UNP P13466
z	148	PRO	-	expression tag	UNP P13466

- Molecule 4 is a protein called Large ribosomal subunit protein bL28.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	0	77	Total	C	N	O	S	0	0
			625	388	129	106	2		

- Molecule 5 is a protein called Large ribosomal subunit protein uL29.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	1	61	Total	C	N	O	S	0	0
			495	305	97	92	1		

- Molecule 6 is a protein called Large ribosomal subunit protein uL30.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	2	57	Total	C	N	O	S	0	0
			439	276	86	75	2		

- Molecule 7 is a protein called Large ribosomal subunit protein bL32.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	3	55	Total	C	N	O	S	0	0
			434	263	92	78	1		

- Molecule 8 is a protein called Large ribosomal subunit protein bL33.

Mol	Chain	Residues	Atoms				AltConf	Trace
8	4	50	Total	C	N	O	0	0
			409	263	75	71		

- Molecule 9 is a protein called Large ribosomal subunit protein bL34.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	6	46	Total	C	N	O	S	0	0
			377	228	90	57	2		

- Molecule 10 is a protein called Large ribosomal subunit protein bL35.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	7	64	Total	C	N	O	S	0	0
			504	323	105	74	2		

- Molecule 11 is a protein called 50S ribosomal protein L36.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	8	38	Total	C	N	O	S	0	0
			301	185	65	47	4		

- Molecule 12 is a RNA chain called 5S rRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	a	118	Total	C	N	O	P	0	0
			2529	1126	464	821	118		

- Molecule 13 is a RNA chain called 23S rRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	b	2901	Total	C	N	O	P	0	0
			62281	27784	11464	20132	2901		

- Molecule 14 is a protein called Large ribosomal subunit protein uL2.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	c	271	Total	C	N	O	S	0	0
			2082	1288	423	364	7		

- Molecule 15 is a protein called 50S ribosomal protein L3.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	d	209	Total	C	N	O	S	0	0
			1565	979	288	294	4		

- Molecule 16 is a protein called Large ribosomal subunit protein uL4.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	e	201	Total	C	N	O	S	0	0
			1552	974	283	290	5		

- Molecule 17 is a protein called Large ribosomal subunit protein uL5.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	f	177	Total	C	N	O	S	0	0
			1410	899	249	256	6		

- Molecule 18 is a protein called Large ribosomal subunit protein uL6.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	g	176	Total	C	N	O	S	0	0
			1323	832	243	246	2		

- Molecule 19 is a protein called Large ribosomal subunit protein bL9.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	h	39	Total	C	N	O	S	0	0
			287	184	51	51	1		

- Molecule 20 is a protein called Large ribosomal subunit protein uL13.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	j	142	Total	C	N	O	S	0	0
			1129	714	212	199	4		

- Molecule 21 is a protein called Large ribosomal subunit protein uL14.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	k	122	Total	C	N	O	S	0	0
			938	587	180	165	6		

- Molecule 22 is a protein called Large ribosomal subunit protein uL15.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	l	143	Total	C	N	O	S	0	0
			1043	649	206	186	2		

- Molecule 23 is a protein called 50S ribosomal protein L16.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	m	136	Total	C	N	O	S	0	0
			1074	686	205	177	6		

- Molecule 24 is a protein called Large ribosomal subunit protein bL17.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	n	120	Total	C	N	O	S	0	0
			960	593	196	166	5		

- Molecule 25 is a protein called Large ribosomal subunit protein uL18.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	o	116	Total	C	N	O		0	0
			892	552	178	162			

- Molecule 26 is a protein called Large ribosomal subunit protein bL19.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	p	113	Total	C	N	O	S	0	0
			908	570	177	160	1		

- Molecule 27 is a protein called Large ribosomal subunit protein bL20.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	q	117	Total	C	N	O		0	0
			947	604	192	151			

- Molecule 28 is a protein called Large ribosomal subunit protein bL21.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	r	102	Total	C	N	O	S	0	0
			810	513	152	143	2		

- Molecule 29 is a protein called Large ribosomal subunit protein uL22.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	s	109	Total	C	N	O	S	0	0
			845	526	162	154	3		

- Molecule 30 is a protein called Large ribosomal subunit protein uL23.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	t	93	Total	C	N	O	S	0	0
			738	466	139	131	2		

- Molecule 31 is a protein called Large ribosomal subunit protein uL24.

Mol	Chain	Residues	Atoms					AltConf	Trace
31	u	102	Total	C	N	O		0	0
			779	492	146	141			

- Molecule 32 is a protein called Large ribosomal subunit protein bL25.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	w	94	Total	C	N	O	S	0	0
			753	479	137	134	3		


- Molecule 33 is a protein called Large ribosomal subunit protein bL27.

Mol	Chain	Residues	Atoms					AltConf	Trace
33	y	82	Total	C	N	O	S	0	0
			619	383	127	108	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: pro-tRNA




• Molecule 1: pro-tRNA




• Molecule 2: mRNA




• Molecule 3: Gelation factor




- Molecule 4: Large ribosomal subunit protein bL28

Chain 0:  85% 14%





- Molecule 5: Large ribosomal subunit protein uL29

Chain 1:  13% 70% 27%




- Molecule 6: Large ribosomal subunit protein uL30

Chain 2:  8% 80% 17%




- Molecule 7: Large ribosomal subunit protein bL32

Chain 3:  70% 25%




- Molecule 8: Large ribosomal subunit protein bL33

Chain 4:  65% 25% 9%




- Molecule 9: Large ribosomal subunit protein bL34

Chain 6:  98%




- Molecule 10: Large ribosomal subunit protein bL35

Chain 7:  94% 5%







- Molecule 11: 50S ribosomal protein L36




- Molecule 12: 5S rRNA





- Molecule 13: 23S rRNA









- Molecule 15: 50S ribosomal protein L3





- Molecule 16: Large ribosomal subunit protein uL4



- Molecule 17: Large ribosomal subunit protein uL5




- Molecule 18: Large ribosomal subunit protein uL6





- Molecule 19: Large ribosomal subunit protein bL9







- Molecule 20: Large ribosomal subunit protein uL13



- Molecule 21: Large ribosomal subunit protein uL14



- Molecule 22: Large ribosomal subunit protein uL15




- Molecule 23: 50S ribosomal protein L16




- Molecule 24: Large ribosomal subunit protein bL17





- Molecule 25: Large ribosomal subunit protein uL18








- Molecule 26: Large ribosomal subunit protein bL19





- Molecule 27: Large ribosomal subunit protein bL20




- Molecule 28: Large ribosomal subunit protein bL21




- Molecule 29: Large ribosomal subunit protein uL22



- Molecule 30: Large ribosomal subunit protein uL23





- Molecule 31: Large ribosomal subunit protein uL24





• Molecule 32: Large ribosomal subunit protein bL25



• Molecule 33: Large ribosomal subunit protein bL27



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	195415	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	22.8	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2400	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.034	Depositor
Minimum map value	-0.007	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.006	Depositor
Map size (\AA)	397.44, 397.44, 397.44	wwPDB
Map dimensions	480, 480, 480	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.828, 0.828, 0.828	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/1817	0.46	0/2832
1	v	0.44	0/1817	0.44	0/2832
2	X	0.55	0/209	0.80	0/323
3	z	1.97	3/192 (1.6%)	2.13	5/267 (1.9%)
4	0	0.39	0/635	0.52	0/848
5	1	0.31	0/496	0.46	0/660
6	2	0.35	0/443	0.40	0/593
7	3	0.47	0/440	0.72	0/588
8	4	0.48	0/416	0.66	0/554
9	6	0.39	0/380	0.48	0/498
10	7	0.46	0/513	0.57	0/676
11	8	0.71	0/302	0.77	0/397
12	a	0.38	0/2828	0.36	0/4410
13	b	0.51	0/69757	0.72	12/108827 (0.0%)
14	c	0.37	0/2121	0.48	0/2852
15	d	0.40	0/1586	0.48	0/2134
16	e	0.33	0/1571	0.43	1/2113 (0.0%)
17	f	0.40	0/1434	0.62	0/1926
18	g	0.35	0/1343	0.52	0/1816
19	h	0.28	0/290	0.63	0/392
20	j	0.42	0/1152	0.47	1/1551 (0.1%)
21	k	0.40	0/947	0.47	0/1268
22	l	0.63	0/1052	0.82	1/1401 (0.1%)
23	m	0.36	0/1093	0.50	0/1460
24	n	0.40	0/973	0.52	0/1301
25	o	0.42	0/902	0.60	0/1209
26	p	0.38	0/920	0.43	0/1231
27	q	0.42	0/960	0.47	0/1278
28	r	0.53	0/823	0.65	0/1100
29	s	0.38	0/852	0.44	0/1142
30	t	0.39	0/744	0.53	0/994
31	u	0.39	0/787	0.57	0/1051
32	w	0.34	0/766	0.43	0/1025
33	y	0.50	0/627	0.62	0/829

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
All	All	0.49	3/101188 (0.0%)	0.67	20/152378 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	z	0	5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	z	131	LEU	C-N	-16.50	1.11	1.33
3	z	130	GLU	C-N	14.88	1.59	1.33
3	z	130	GLU	N-CA	7.46	1.55	1.46

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	z	130	GLU	O-C-N	-26.09	87.89	122.59
13	b	2832	U	C4'-C3'-O3'	7.79	121.09	109.40
13	b	228	C	C4'-C3'-O3'	6.93	119.80	109.40
3	z	130	GLU	CA-C-N	6.82	133.23	123.14
3	z	130	GLU	C-N-CA	6.82	133.23	123.14

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	z	129	VAL	Mainchain
3	z	130	GLU	Mainchain
3	z	131	LEU	Mainchain
3	z	143	ARG	Sidechain
3	z	145	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1626	0	821	59	0
1	v	1626	0	821	58	0
2	X	189	0	99	9	0
3	z	181	0	173	33	0
4	0	625	0	652	7	0
5	1	495	0	526	13	0
6	2	439	0	482	6	0
7	3	434	0	445	11	0
8	4	409	0	440	8	0
9	6	377	0	418	1	0
10	7	504	0	572	2	0
11	8	301	0	341	22	0
12	a	2529	0	1281	14	0
13	b	62281	0	31323	454	0
14	c	2082	0	2154	30	0
15	d	1565	0	1616	24	0
16	e	1552	0	1619	25	0
17	f	1410	0	1444	42	0
18	g	1323	0	1371	27	0
19	h	287	0	307	13	0
20	j	1129	0	1162	21	0
21	k	938	0	1012	25	0
22	l	1043	0	1123	11	0
23	m	1074	0	1157	20	0
24	n	960	0	1000	18	0
25	o	892	0	923	17	0
26	p	908	0	956	22	0
27	q	947	0	1019	16	0
28	r	810	0	834	5	0
29	s	845	0	909	34	0
30	t	738	0	807	9	0
31	u	779	0	831	25	0
32	w	753	0	780	4	0
33	y	619	0	642	22	0
All	All	92670	0	60060	959	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 959 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
13:b:892:A:H8	13:b:893:C:C6	1.26	1.52
13:b:892:A:C8	13:b:893:C:C6	1.96	1.49
13:b:892:A:C8	13:b:893:C:C5	2.13	1.37
13:b:370:G:OP2	13:b:370:G:C8	1.81	1.32
13:b:851:C:O2'	13:b:852:U:H5'	1.41	1.20

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	z	18/148 (12%)	15 (83%)	1 (6%)	2 (11%)	0	0
4	0	75/78 (96%)	74 (99%)	1 (1%)	0	100	100
5	1	59/63 (94%)	56 (95%)	3 (5%)	0	100	100
6	2	55/59 (93%)	53 (96%)	2 (4%)	0	100	100
7	3	53/57 (93%)	49 (92%)	4 (8%)	0	100	100
8	4	48/55 (87%)	45 (94%)	3 (6%)	0	100	100
9	6	44/46 (96%)	43 (98%)	1 (2%)	0	100	100
10	7	62/65 (95%)	59 (95%)	3 (5%)	0	100	100
11	8	36/50 (72%)	26 (72%)	7 (19%)	3 (8%)	0	0
14	c	269/273 (98%)	258 (96%)	11 (4%)	0	100	100
15	d	207/209 (99%)	197 (95%)	10 (5%)	0	100	100
16	e	199/201 (99%)	196 (98%)	3 (2%)	0	100	100
17	f	175/179 (98%)	164 (94%)	10 (6%)	1 (1%)	21	42
18	g	174/177 (98%)	159 (91%)	15 (9%)	0	100	100
19	h	37/149 (25%)	32 (86%)	5 (14%)	0	100	100
20	j	140/142 (99%)	138 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
21	k	120/123 (98%)	116 (97%)	4 (3%)	0	100	100
22	l	141/144 (98%)	132 (94%)	9 (6%)	0	100	100
23	m	134/136 (98%)	133 (99%)	1 (1%)	0	100	100
24	n	118/127 (93%)	111 (94%)	7 (6%)	0	100	100
25	o	114/117 (97%)	109 (96%)	5 (4%)	0	100	100
26	p	111/115 (96%)	110 (99%)	1 (1%)	0	100	100
27	q	115/118 (98%)	115 (100%)	0	0	100	100
28	r	100/103 (97%)	93 (93%)	5 (5%)	2 (2%)	6	12
29	s	107/110 (97%)	100 (94%)	7 (6%)	0	100	100
30	t	91/100 (91%)	87 (96%)	4 (4%)	0	100	100
31	u	100/104 (96%)	83 (83%)	17 (17%)	0	100	100
32	w	92/94 (98%)	91 (99%)	1 (1%)	0	100	100
33	y	80/85 (94%)	73 (91%)	5 (6%)	2 (2%)	4	8
All	All	3074/3427 (90%)	2917 (95%)	147 (5%)	10 (0%)	37	58

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
28	r	52	PRO
11	8	29	ALA
33	y	5	LYS
3	z	130	GLU
11	8	19	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	z	19/121 (16%)	13 (68%)	6 (32%)	0	0
4	0	67/68 (98%)	67 (100%)	0	100	100
5	1	54/55 (98%)	54 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
6	2	47/49 (96%)	47 (100%)	0	100	100
7	3	46/48 (96%)	44 (96%)	2 (4%)	26	51
8	4	45/49 (92%)	44 (98%)	1 (2%)	45	72
9	6	38/38 (100%)	38 (100%)	0	100	100
10	7	51/52 (98%)	51 (100%)	0	100	100
11	8	34/44 (77%)	28 (82%)	6 (18%)	2	3
14	c	216/218 (99%)	216 (100%)	0	100	100
15	d	164/164 (100%)	164 (100%)	0	100	100
16	e	165/165 (100%)	165 (100%)	0	100	100
17	f	148/150 (99%)	144 (97%)	4 (3%)	39	67
18	g	137/138 (99%)	136 (99%)	1 (1%)	76	89
19	h	30/114 (26%)	30 (100%)	0	100	100
20	j	116/116 (100%)	116 (100%)	0	100	100
21	k	103/104 (99%)	103 (100%)	0	100	100
22	l	102/103 (99%)	94 (92%)	8 (8%)	11	26
23	m	109/109 (100%)	109 (100%)	0	100	100
24	n	100/103 (97%)	100 (100%)	0	100	100
25	o	86/87 (99%)	82 (95%)	4 (5%)	23	48
26	p	98/100 (98%)	98 (100%)	0	100	100
27	q	89/90 (99%)	89 (100%)	0	100	100
28	r	84/84 (100%)	79 (94%)	5 (6%)	17	37
29	s	92/93 (99%)	91 (99%)	1 (1%)	65	84
30	t	80/84 (95%)	78 (98%)	2 (2%)	42	69
31	u	83/85 (98%)	82 (99%)	1 (1%)	63	83
32	w	78/78 (100%)	78 (100%)	0	100	100
33	y	61/63 (97%)	57 (93%)	4 (7%)	15	34
All	All	2542/2772 (92%)	2497 (98%)	45 (2%)	51	76

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
25	o	15	ARG
28	r	51	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
25	o	67	ASN
28	r	40	MET
29	s	85	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 47 such sidechains are listed below:

Mol	Chain	Res	Type
24	n	9	GLN
26	p	66	ASN
25	o	29	HIS
25	o	116	GLN
27	q	72	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	75/77 (97%)	20 (26%)	3 (4%)
1	v	75/77 (97%)	20 (26%)	0
12	a	117/120 (97%)	21 (17%)	0
13	b	2900/2904 (99%)	1199 (41%)	0
2	X	8/9 (88%)	5 (62%)	2 (25%)
All	All	3175/3187 (99%)	1265 (39%)	5 (0%)

5 of 1265 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	v	3	G
1	v	8	U
1	v	9	G
1	v	10	G
1	v	13	C

All (5) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	9	G
1	A	14	A
1	A	20	G
2	X	14	U
2	X	17	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	z	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	z	131:LEU	C	132:PHE	N	1.11


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-52500. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 240




Y Index: 240




Z Index: 240


6.2.2 Raw map



X Index: 240



Y Index: 240




Z Index: 240


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 271




Y Index: 248




Z Index: 255


6.3.2 Raw map



X Index: 266



Y Index: 235




Z Index: 246


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.006. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_52500_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1757 nm^3 ; this corresponds to an approximate mass of 1587 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.385 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.385 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.60	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.99	8.22	4.22

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.99 differs from the reported value 2.6 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-52500 and PDB model 9HY6. Per-residue inclusion information can be found in [section 3](#) on [page 11](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.006 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.006).







































































9.4 Atom inclusion [i](#)



At the recommended contour level, 92% of all backbone atoms, 89% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.006) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8860	 0.3520
0	 0.8870	 0.4580
1	 0.6460	 0.2640
2	 0.7890	 0.3130
3	 0.8090	 0.3580
4	 0.7950	 0.3370
6	 0.9410	 0.4580
7	 0.9410	 0.4540
8	 0.8560	 0.2960
A	 0.9470	 0.3570
X	 0.9520	 0.4080
a	 0.9160	 0.2960
b	 0.9210	 0.3620
c	 0.9050	 0.4420
d	 0.8220	 0.3530
e	 0.7430	 0.3520
f	 0.7040	 0.2640
g	 0.6000	 0.2010
h	 0.4960	 0.3320
j	 0.7930	 0.2940
k	 0.8450	 0.4040
l	 0.8380	 0.4150
m	 0.8220	 0.3240
n	 0.8340	 0.3500
o	 0.7270	 0.2610
p	 0.8030	 0.3390
q	 0.8080	 0.3360
r	 0.7240	 0.3140
s	 0.8290	 0.3690
t	 0.7600	 0.2970
u	 0.6510	 0.2670
v	 0.9010	 0.2860
w	 0.6970	 0.2540
y	 0.8660	 0.3510
z	 0.7690	 0.2600

