wwPDB X-ray Structure Validation Summary Report (i)

Aug 20, 2023 – 08:49 AM EDT

PDB ID	:	2HZV
Title	:	NikR-operator DNA complex
Authors	:	Schreiter, E.R.; Drennan, C.L.
Deposited on	:	2006-08-09
Resolution	:	3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	I	30	10% 90%
1	K	30	23% 77%
2	J	30	17% 83%
2	L	30	20% 77% •
3	A	133	2% 41% 50% 6% • •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	B	133	43% 51% 5%
3	C	133	2% 48% 45% 5%
3	D	133	3% 47% 45% 6%
3	E	133	47% 41% 10%
3	F	133	2% 50% 44%
3	G	133	43% 49% 6%
3	H	133	% 45% 47% 7%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10501 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*AP*GP*TP*AP*TP*GP*AP*CP*GP*AP*AP*TP*AP*CP*TP*TP*AP*AP*AP*AP*TP*CP*GP*TP*CP*AP*TP*AP*CP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	I	30	612	295	113	175	29	0	0	0
1	K	30	612	295	113	175	29	0	0	0

- Molecule 2 is a DNA chain called 5'-D(*AP*GP*TP*AP*TP*GP*AP*CP*GP*AP*TP*TP*TP*TP*AP*AP*GP*TP*AP*TP*TP*CP*GP*TP*CP*AP*TP*AP*CP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	J	30	612	296	106	181	29	0	0	0
2	L	30	612	296	106	181	29	0	0	0

- Molecule 3 is a protein called Nickel-responsive regulator.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
3	A	131	981	603	184	190	2	2	0	0	0
3	B	131	1031	631	198	198	2	2	0	0	0
3	C	131	971	598	183	186	2	2	0	0	0
3	D	131	1027	629	197	197	2	2	0	0	0
3	E	131	981	603	184	190	2	2	0	0	0
3	F	131	1031	631	198	198	2	2	0	0	0
3	G	131	985	605	185	191	2	2	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
3	H	131	1027	629	197	197	2	2	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP P0A6Z6
A	105	MSE	MET	modified residue	UNP P0A6Z6
B	1	MSE	MET	modified residue	UNP P0A6Z6
B	105	MSE	MET	modified residue	UNP P0A6Z6
C	1	MSE	MET	modified residue	UNP P0A6Z6
C	105	MSE	MET	modified residue	UNP P0A6Z6
D	1	MSE	MET	modified residue	UNP P0A6Z6
D	105	MSE	MET	modified residue	UNP P0A6Z6
E	1	MSE	MET	modified residue	UNP P0A6Z6
E	105	MSE	MET	modified residue	UNP P0A6Z6
F	1	MSE	MET	modified residue	UNP P0A6Z6
F	105	MSE	MET	modified residue	UNP P0A6Z6
G	1	MSE	MET	modified residue	UNP P0A6Z6
G	105	MSE	MET	modified residue	UNP P0A6Z6
H	1	MSE	MET	modified residue	UNP P0A6Z6
H	105	MSE	MET	modified residue	UNP P0A6Z6

- Molecule 4 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ni		
4	A	1	1	1	0	0
4	C	1	1	1	0	0
4	D	2	2	2	0	0
4	E	1	1	1	0	0
4	F	1	1	1	0	0
4	G	1	1	1	0	0
4	H	1	1	1	0	0

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	K		
5	A	2	2	2	0	0
			Total	K		
5	B	1	1	1	0	0
			Total	K		
5	C	2	2	2	0	0
			Total	K		
5	E	2	2	2	0	0
			Total	K		
5	F	1	1	1	0	0
			Total	K		
5	G	2	2	2	0	0
			Total	K		
5	H	1	1	1	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 5'-D(*AP*GP*TP*AP*TP*GP*AP*CP*GP*AP*AP*TP*AP*CP*TP*TP*AP*AP*AP*AP*TP*CP*GP*TP*CP*AP*TP*AP*CP*T)-3'

Chain I:





• Molecule 1: 5'-D(*AP*GP*TP*AP*TP*GP*AP*CP*GP*AP*AP*TP*AP*CP*TP*TP*AP*AP*AP*AP*TP*CP*GP*TP*CP*AP*TP*AP*CP*T)-3'

Chain K:





• Molecule 2: 5'-D(*AP*GP*TP*AP*TP*GP*AP*CP*GP*AP*TP*TP*TP*TP*AP*AP*GP*TP*AP*TP*TP*CP*GP*TP*CP*AP*TP*AP*CP*T)-3'

Chain J:



• Molecule 2: 5'-D(*AP*GP*TP*AP*TP*GP*AP*CP*GP*AP*TP*TP*TP*TP*AP*AP*GP*TP*AP*TP*TP*CP*GP*TP*CP*AP*TP*AP*CP*T)-3'

Chain L:





• Molecule 3: Nickel-responsive regulator

Chain A:



• Molecule 3: Nickel-responsive regulator

Chain B:





• Molecule 3: Nickel-responsive regulator

Chain C:



• Molecule 3: Nickel-responsive regulator

Chain D:



• Molecule 3: Nickel-responsive regulator

Chain E:





• Molecule 3: Nickel-responsive regulator

Chain F:



• Molecule 3: Nickel-responsive regulator

Chain G:





• Molecule 3: Nickel-responsive regulator

Chain H:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	197.20Å 76.13Å 132.09Å 90.00° 110.27° 90.00°	Depositor
Resolution (Å)	49.28 – 3.10 49.28 – 3.10	Depositor EDS
% Data completeness (in resolution range)	94.1 (49.28-3.10) 89.8 (49.28-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	5.16 (at 3.12Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.261 , 0.303 0.247 , 0.293	Depositor DCC
R_{free} test set	2119 reflections (6.67%)	wwPDB-VP
Wilson B-factor (Å^2)	72.9	Xtriage
Anisotropy	0.965	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.26 , 56.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	10501	wwPDB-VP
Average B, all atoms (Å^2)	99.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.52% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	I	0.58	0/687	0.87	0/1058
1	K	0.62	0/687	0.87	0/1058
2	J	0.54	0/685	0.81	0/1056
2	L	0.67	0/685	0.85	0/1056
3	A	0.39	0/995	0.66	0/1352
3	B	0.45	0/1047	0.72	1/1416 (0.1%)
3	C	0.40	0/985	0.67	0/1339
3	D	0.42	0/1043	0.66	0/1411
3	E	0.42	0/995	0.67	0/1352
3	F	0.45	0/1047	0.69	0/1416
3	G	0.46	0/999	0.77	3/1357 (0.2%)
3	H	0.49	0/1043	0.70	0/1411
All	All	0.49	0/10898	0.74	4/15282 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	L	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	G	66	ASP	CB-CA-C	7.33	125.06	110.40
3	B	119	ARG	NE-CZ-NH2	6.40	123.50	120.30
3	G	67	LEU	N-CA-C	5.45	125.72	111.00
3	G	23	ARG	NE-CZ-NH1	-5.18	117.71	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	L	29	DC	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	I	612	0	341	36	0
1	K	612	0	341	29	0
2	J	612	0	344	45	0
2	L	612	0	344	36	0
3	A	981	0	912	73	0
3	B	1031	0	989	76	0
3	C	971	0	897	62	0
3	D	1027	0	983	78	0
3	E	981	0	912	61	0
3	F	1031	0	989	70	0
3	G	985	0	918	63	0
3	H	1027	0	983	80	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
4	D	2	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
5	A	2	0	0	0	0
5	B	1	0	0	0	0
5	C	2	0	0	0	0
5	E	2	0	0	0	0
5	F	1	0	0	0	0
5	G	2	0	0	0	0
5	H	1	0	0	0	0
All	All	10501	0	8953	632	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 632 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:4:DA:H2"	1:K:5:DT:H5"	1.21	1.16
1:K:4:DA:H2"	1:K:5:DT:C5'	1.83	1.06
3:E:30:GLU:HG2	3:E:33:ARG:HH22	1.24	1.02
3:G:51:GLN:HG2	3:G:104:ASP:HA	1.45	0.98
1:K:4:DA:C2'	1:K:5:DT:H5"	1.95	0.96

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	A	129/133 (97%)	97 (75%)	23 (18%)	9 (7%)	1 7
3	B	129/133 (97%)	102 (79%)	21 (16%)	6 (5%)	2 14
3	C	129/133 (97%)	99 (77%)	26 (20%)	4 (3%)	4 23
3	D	129/133 (97%)	99 (77%)	24 (19%)	6 (5%)	2 14
3	E	129/133 (97%)	100 (78%)	18 (14%)	11 (8%)	1 5
3	F	129/133 (97%)	102 (79%)	20 (16%)	7 (5%)	2 12
3	G	129/133 (97%)	99 (77%)	23 (18%)	7 (5%)	2 12
3	H	129/133 (97%)	101 (78%)	21 (16%)	7 (5%)	2 12
All	All	1032/1064 (97%)	799 (77%)	176 (17%)	57 (6%)	2 11

5 of 57 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	118	GLN
3	B	38	SER
3	C	130	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	E	44	ALA
3	E	48	HIS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	A	100/114 (88%)	90 (90%)	10 (10%)	7 28
3	B	111/114 (97%)	106 (96%)	5 (4%)	27 60
3	C	97/114 (85%)	85 (88%)	12 (12%)	4 19
3	D	110/114 (96%)	102 (93%)	8 (7%)	14 43
3	E	100/114 (88%)	86 (86%)	14 (14%)	3 15
3	F	111/114 (97%)	108 (97%)	3 (3%)	44 74
3	G	101/114 (89%)	89 (88%)	12 (12%)	5 20
3	H	110/114 (96%)	104 (94%)	6 (6%)	21 53
All	All	840/912 (92%)	770 (92%)	70 (8%)	11 38

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G	42	GLN
3	G	78	HIS
3	H	7	THR
3	C	107	ASP
3	C	78	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	123	HIS
3	E	125	HIS
3	E	46	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	E	127	GLN
3	B	21	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 19 ligands modelled in this entry, 19 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	I	30/30 (100%)	-0.77	0	100	100	69, 95, 115, 131	0
1	K	30/30 (100%)	-0.72	0	100	100	65, 89, 110, 120	0
2	J	30/30 (100%)	-0.74	0	100	100	70, 91, 118, 127	0
2	L	30/30 (100%)	-0.56	0	100	100	63, 86, 133, 150	0
3	A	129/133 (96%)	-0.06	2 (1%)	72	51	58, 105, 169, 192	0
3	B	129/133 (96%)	-0.15	0	100	100	59, 95, 156, 180	0
3	C	129/133 (96%)	-0.06	2 (1%)	72	51	60, 99, 163, 186	0
3	D	129/133 (96%)	0.07	4 (3%)	49	26	63, 96, 167, 199	0
3	E	129/133 (96%)	-0.20	0	100	100	59, 99, 164, 188	0
3	F	129/133 (96%)	-0.07	2 (1%)	72	51	49, 92, 159, 182	0
3	G	129/133 (96%)	-0.04	0	100	100	53, 92, 159, 186	0
3	H	129/133 (96%)	-0.08	1 (0%)	86	72	49, 85, 136, 148	0
All	All	1152/1184 (97%)	-0.14	11 (0%)	82	67	49, 95, 161, 199	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	25	TYR	3.6
3	D	102	LYS	3.1
3	H	2	GLN	3.1
3	A	60	TYR	2.9
3	D	101	LEU	2.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	K	A	409	1/1	0.66	0.13	129,129,129,129	0
5	K	E	403	1/1	0.78	0.12	111,111,111,111	0
5	K	G	404	1/1	0.78	0.18	94,94,94,94	0
5	K	E	406	1/1	0.79	0.17	121,121,121,121	0
5	K	C	411	1/1	0.79	0.12	123,123,123,123	0
5	K	B	410	1/1	0.81	0.20	122,122,122,122	0
5	K	C	402	1/1	0.82	0.12	102,102,102,102	0
5	K	F	408	1/1	0.84	0.17	140,140,140,140	0
5	K	G	407	1/1	0.92	0.18	135,135,135,135	0
5	K	H	405	1/1	0.92	0.21	90,90,90,90	0
4	NI	C	303	1/1	0.93	0.06	109,109,109,109	0
4	NI	A	302	1/1	0.94	0.09	104,104,104,104	0
4	NI	E	306	1/1	0.95	0.12	106,106,106,106	0
5	K	A	401	1/1	0.96	0.17	97,97,97,97	0
4	NI	D	301	1/1	0.97	0.13	84,84,84,84	0
4	NI	H	308	1/1	0.98	0.18	73,73,73,73	0
4	NI	G	307	1/1	0.98	0.10	78,78,78,78	0
4	NI	F	305	1/1	0.99	0.21	72,72,72,72	0
4	NI	D	304	1/1	0.99	0.18	86,86,86,86	0

6.5 Other polymers (i)

There are no such residues in this entry.