



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 10, 2023 – 06:44 AM EDT

PDB ID : 4I9N  
Title : Crystal structure of rabbit LDHA in complex with AP28161 and AP28122  
Authors : Zhou, T.; Kohlmann, A.; Stephan, Z.G.; Li, F.; Commodore, L.; Greenfield, M.T.; Shakespeare, W.C.; Zhu, X.; Dalgarno, D.C.  
Deposited on : 2012-12-05  
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	 2% 75% 21% ..
1	B	331	 4% 70% 27% ..
1	C	331	 3% 75% 23% ..
1	D	331	 3% 69% 30% .
1	E	331	 6% 69% 30% .

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	331	
1	G	331	
1	H	331	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 20952 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-lactate dehydrogenase A chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	324	2502	1602	429	457	14	0	0	0
1	B	324	2502	1602	429	457	14	0	0	0
1	C	328	2532	1618	436	464	14	0	0	0
1	D	331	2559	1633	441	471	14	0	0	0
1	E	331	2559	1633	441	471	14	0	0	0
1	F	329	2540	1622	437	467	14	0	0	0
1	G	328	2537	1621	437	465	14	0	0	0
1	H	325	2508	1605	430	459	14	0	0	0

- Molecule 2 is 6-({2-[(5-chloro-4-[(2S)-2,3-dihydroxypropyl]oxy})-2-methoxyphenyl]amino}-2-oxoethyl)sulfanylpyridine-3-carboxylic acid (three-letter code: 1E6) (formula: C<sub>18</sub>H<sub>19</sub>ClN<sub>2</sub>O<sub>7</sub>S).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
3	A	1	Total 21	C 14	F 1	N 1	O 5	0	0
3	B	1	Total 21	C 14	F 1	N 1	O 5	0	0
3	C	1	Total 21	C 14	F 1	N 1	O 5	0	0
3	D	1	Total 21	C 14	F 1	N 1	O 5	0	0
3	E	1	Total 21	C 14	F 1	N 1	O 5	0	0
3	F	1	Total 21	C 14	F 1	N 1	O 5	0	0
3	G	1	Total 21	C 14	F 1	N 1	O 5	0	0
3	H	1	Total 21	C 14	F 1	N 1	O 5	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	54	Total 54	O 54	0	0
4	B	41	Total 41	O 41	0	0
4	C	36	Total 36	O 36	0	0
4	D	43	Total 43	O 43	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
4	E	26	Total O 26 26	0	0
4	F	32	Total O 32 32	0	0
4	G	42	Total O 42 42	0	0
4	H	39	Total O 39 39	0	0







- Molecule 1: L-lactate dehydrogenase A chain



- Molecule 1: L-lactate dehydrogenase A chain




- Molecule 1: L-lactate dehydrogenase A chain




- Molecule 1: L-lactate dehydrogenase A chain





• Molecule 1: L-lactate dehydrogenase A chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.04Å 139.81Å 138.93Å 90.00° 94.57° 90.00°	Depositor
Resolution (Å)	50.00 – 2.35 49.21 – 2.25	Depositor EDS
% Data completeness (in resolution range)	94.7 (50.00-2.35) 93.7 (49.21-2.25)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.49 (at 2.24Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.220 , 0.264 0.212 , 0.255	Depositor DCC
$R_{free}$ test set	7246 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.4	Xtrriage
Anisotropy	0.322	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	20952	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1E6, 1E5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/2547	0.52	0/3445
1	B	0.36	0/2547	0.54	0/3445
1	C	0.32	0/2577	0.51	0/3484
1	D	0.33	0/2605	0.50	0/3523
1	E	0.31	0/2605	0.48	0/3523
1	F	0.32	0/2584	0.50	0/3493
1	G	0.33	0/2582	0.51	0/3491
1	H	0.34	0/2553	0.52	0/3453
All	All	0.33	0/20600	0.51	0/27857

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	H	0	1
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	107	ASN	Peptide
1	B	16	HIS	Peptide

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	H	15	GLU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2502	0	2589	60	0
1	B	2502	0	2589	74	0
1	C	2532	0	2616	64	0
1	D	2559	0	2639	88	0
1	E	2559	0	2639	89	0
1	F	2540	0	2625	84	0
1	G	2537	0	2621	83	0
1	H	2508	0	2594	91	0
2	A	29	0	18	0	0
2	B	29	0	18	0	0
2	C	29	0	18	0	0
2	D	29	0	18	0	0
2	E	29	0	18	1	0
2	F	29	0	18	0	0
2	G	29	0	18	0	0
2	H	29	0	18	0	0
3	A	21	0	8	1	0
3	B	21	0	9	1	0
3	C	21	0	8	1	0
3	D	21	0	8	1	0
3	E	21	0	8	1	0
3	F	21	0	8	1	0
3	G	21	0	9	1	0
3	H	21	0	9	1	0
4	A	54	0	0	1	0
4	B	41	0	0	0	0
4	C	36	0	0	0	0
4	D	43	0	0	2	0
4	E	26	0	0	0	0
4	F	32	0	0	0	0
4	G	42	0	0	0	0
4	H	39	0	0	3	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	20952	0	21123	580	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 580 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:304:LYS:HD2	1:D:9:HIS:HB2	1.45	0.95
1:D:97:ALA:H	1:D:112:ASN:HD21	1.11	0.94
1:F:276:LEU:HD21	1:F:288:LEU:HD12	1.48	0.94
1:H:110:GLN:HE22	1:H:330:GLN:H	1.10	0.93
1:E:98:ARG:H	1:E:98:ARG:HE	0.94	0.91

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	320/331 (97%)	305 (95%)	12 (4%)	3 (1%)	17 17
1	B	320/331 (97%)	307 (96%)	12 (4%)	1 (0%)	41 47
1	C	324/331 (98%)	310 (96%)	13 (4%)	1 (0%)	41 47
1	D	329/331 (99%)	314 (95%)	15 (5%)	0	100 100
1	E	329/331 (99%)	317 (96%)	10 (3%)	2 (1%)	25 27
1	F	325/331 (98%)	311 (96%)	11 (3%)	3 (1%)	17 17
1	G	324/331 (98%)	312 (96%)	11 (3%)	1 (0%)	41 47
1	H	321/331 (97%)	309 (96%)	10 (3%)	2 (1%)	25 27
All	All	2592/2648 (98%)	2485 (96%)	94 (4%)	13 (0%)	29 32

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	107	ASN
1	A	329	LEU
1	E	14	GLU
1	E	16	HIS
1	F	101	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/284 (98%)	270 (97%)	8 (3%)	42	52
1	B	278/284 (98%)	269 (97%)	9 (3%)	39	47
1	C	281/284 (99%)	274 (98%)	7 (2%)	47	58
1	D	284/284 (100%)	278 (98%)	6 (2%)	53	65
1	E	284/284 (100%)	274 (96%)	10 (4%)	36	44
1	F	282/284 (99%)	272 (96%)	10 (4%)	36	44
1	G	282/284 (99%)	274 (97%)	8 (3%)	43	53
1	H	279/284 (98%)	270 (97%)	9 (3%)	39	47
All	All	2248/2272 (99%)	2181 (97%)	67 (3%)	41	50

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	330	GLN
1	H	14	GLU
1	H	310	GLU
1	D	12	LEU
1	C	285	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	297	ASN
1	H	9	HIS
1	E	204	ASN
1	G	228	GLN
1	H	114	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	1E5	C	402	-	22,22,22	3.08	11 (50%)	30,30,30	1.47	6 (20%)
2	1E6	D	401	-	30,30,30	4.04	13 (43%)	39,40,40	2.46	8 (20%)
3	1E5	D	402	-	22,22,22	2.95	7 (31%)	30,30,30	1.49	4 (13%)
3	1E5	G	402	-	22,22,22	2.74	9 (40%)	30,30,30	1.31	4 (13%)
2	1E6	C	401	-	30,30,30	3.95	11 (36%)	39,40,40	2.41	7 (17%)
2	1E6	F	401	-	30,30,30	3.92	11 (36%)	39,40,40	2.43	7 (17%)
2	1E6	B	401	-	30,30,30	3.84	11 (36%)	39,40,40	2.50	7 (17%)
3	1E5	B	402	-	22,22,22	2.72	10 (45%)	30,30,30	1.35	3 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	1E6	G	401	-	30,30,30	3.86	12 (40%)	39,40,40	2.39	7 (17%)
3	1E5	F	402	-	22,22,22	3.02	10 (45%)	30,30,30	1.46	7 (23%)
2	1E6	A	401	-	30,30,30	3.84	12 (40%)	39,40,40	2.49	7 (17%)
2	1E6	H	401	-	30,30,30	3.85	13 (43%)	39,40,40	2.46	7 (17%)
3	1E5	H	402	-	22,22,22	2.73	10 (45%)	30,30,30	1.36	4 (13%)
3	1E5	A	402	-	22,22,22	2.83	8 (36%)	30,30,30	1.30	4 (13%)
3	1E5	E	402	-	22,22,22	3.12	13 (59%)	30,30,30	1.44	5 (16%)
2	1E6	E	401	-	30,30,30	4.07	15 (50%)	39,40,40	2.54	8 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1E5	C	402	-	-	2/13/13/13	0/2/2/2
2	1E6	D	401	-	-	3/22/22/22	0/2/2/2
3	1E5	D	402	-	-	3/13/13/13	0/2/2/2
3	1E5	G	402	-	-	2/13/13/13	0/2/2/2
2	1E6	C	401	-	-	5/22/22/22	0/2/2/2
2	1E6	F	401	-	-	5/22/22/22	0/2/2/2
2	1E6	B	401	-	-	2/22/22/22	0/2/2/2
3	1E5	B	402	-	-	2/13/13/13	0/2/2/2
2	1E6	G	401	-	-	1/22/22/22	0/2/2/2
3	1E5	F	402	-	-	2/13/13/13	0/2/2/2
2	1E6	A	401	-	-	4/22/22/22	0/2/2/2
2	1E6	H	401	-	-	5/22/22/22	0/2/2/2
3	1E5	H	402	-	-	4/13/13/13	0/2/2/2
3	1E5	A	402	-	-	2/13/13/13	0/2/2/2
3	1E5	E	402	-	-	3/13/13/13	0/2/2/2
2	1E6	E	401	-	-	6/22/22/22	0/2/2/2

The worst 5 of 176 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	401	1E6	C11-C12	15.45	1.69	1.51
2	C	401	1E6	C11-C12	15.17	1.68	1.51

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	401	1E6	C11-C12	14.70	1.68	1.51
2	G	401	1E6	C11-C12	14.63	1.68	1.51
2	F	401	1E6	C11-C12	14.56	1.68	1.51

The worst 5 of 95 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	401	1E6	O43-C12-C11	-10.86	102.82	121.58
2	H	401	1E6	O43-C12-C11	-10.83	102.88	121.58
2	B	401	1E6	O43-C12-C11	-10.79	102.95	121.58
2	F	401	1E6	O43-C12-C11	-10.63	103.21	121.58
2	D	401	1E6	O43-C12-C11	-10.63	103.22	121.58

There are no chirality outliers.

5 of 51 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	401	1E6	C23-C24-C27-O28
2	H	401	1E6	O44-C24-C27-O28
3	H	402	1E5	C19-C20-C23-O24
3	H	402	1E5	C21-C20-C23-O24
2	C	401	1E6	O3-C1-C2-C5


There are no ring outliers.



9 monomers are involved in 9 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	402	1E5	1	0
3	D	402	1E5	1	0
3	G	402	1E5	1	0
3	B	402	1E5	1	0
3	F	402	1E5	1	0
3	H	402	1E5	1	0
3	A	402	1E5	1	0
3	E	402	1E5	1	0
2	E	401	1E6	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is



within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	324/331 (97%)	0.01	8 (2%) 57 67	14, 25, 44, 65	1 (0%)
1	B	324/331 (97%)	-0.06	13 (4%) 38 51	14, 24, 45, 63	1 (0%)
1	C	328/331 (99%)	0.17	10 (3%) 50 61	17, 30, 50, 85	1 (0%)
1	D	331/331 (100%)	0.00	11 (3%) 46 59	15, 26, 48, 72	1 (0%)
1	E	331/331 (100%)	0.29	21 (6%) 20 29	19, 33, 61, 77	1 (0%)
1	F	329/331 (99%)	0.18	19 (5%) 23 33	13, 30, 59, 79	1 (0%)
1	G	328/331 (99%)	0.09	12 (3%) 41 54	15, 27, 55, 86	1 (0%)
1	H	325/331 (98%)	0.04	10 (3%) 49 61	13, 26, 47, 94	1 (0%)
All	All	2620/2648 (98%)	0.09	104 (3%) 38 51	13, 27, 53, 94	8 (0%)

The worst 5 of 104 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	102	GLY	13.0
1	H	105	ARG	8.4
1	G	98	ARG	8.2
1	G	106	LEU	7.0
1	H	12	LEU	6.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	1E6	E	401	29/29	0.92	0.16	22,37,40,44	0
3	1E5	E	402	21/21	0.92	0.12	29,32,35,36	0
3	1E5	F	402	21/21	0.92	0.15	23,27,29,31	0
2	1E6	F	401	29/29	0.93	0.16	14,33,40,43	0
2	1E6	C	401	29/29	0.93	0.18	18,36,40,41	0
2	1E6	B	401	29/29	0.93	0.17	3,28,33,34	0
3	1E5	C	402	21/21	0.94	0.13	22,23,28,31	0
2	1E6	A	401	29/29	0.94	0.18	17,31,35,36	0
2	1E6	D	401	29/29	0.94	0.15	17,34,35,37	0
2	1E6	G	401	29/29	0.95	0.17	13,28,35,39	0
3	1E5	D	402	21/21	0.95	0.13	18,23,24,28	0
2	1E6	H	401	29/29	0.95	0.19	11,31,39,42	0
3	1E5	B	402	21/21	0.95	0.11	19,22,24,26	0
3	1E5	G	402	21/21	0.95	0.12	24,27,28,30	0
3	1E5	H	402	21/21	0.95	0.13	22,25,27,28	0
3	1E5	A	402	21/21	0.96	0.12	22,25,31,33	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 1E6 E 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 1E5 E 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 1E5 F 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 1E6 F 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 1E6 C 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 1E6 B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 1E5 C 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 1E6 A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 1E6 D 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 1E6 G 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 1E5 D 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 1E6 H 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 1E5 B 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 1E5 G 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 1E5 H 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 1E5 A 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers

There are no such residues in this entry.