



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 03:07 PM UTC

PDB ID : 9IDZ / pdb_00009idz
Title : Human Deoxyhypusine Synthase Fragment Screening Campaign - ligand VT00221
Authors : Wilk, P.; Water-Wilk, E.; Krojer, T.; Grudnik, P.
Deposited on : 2025-02-15
Resolution : 1.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3234 (1.46-1.42)
Clashscore	190562	3289 (1.46-1.42)
Ramachandran outliers	187476	3248 (1.46-1.42)
Sidechain outliers	187428	3248 (1.46-1.42)
RSRZ outliers	180081	3234 (1.46-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	
1	B	369	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6034 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deoxyhypusine synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	335	Total	C	N	O	S	0	6	0
			2663	1700	450	497	16			
1	B	352	Total	C	N	O	S	0	6	0
			2781	1770	471	523	17			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is (4R)-2-METHYLPENTANE-2,4-DIOL (CCD ID: MRD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 4 is QUINOLIN-2-AMINE (CCD ID: 2AQ) (formula: $C_9H_8N_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	N	0	0
			11	9	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	247	Total 247	O 247	0	0
5	B	236	Total 236	O 236	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Deoxyhypusine synthase



• Molecule 1: Deoxyhypusine synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.78Å 104.78Å 160.35Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.05 – 1.44 46.05 – 1.44	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.05-1.44) 99.9 (46.05-1.44)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.01 (at 1.44Å)	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.200 , 0.214 (Not available) , 0.221	Depositor DCC
R_{free} test set	9107 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtriage
Anisotropy	0.090	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.012 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6034	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2AQ, NAD, MRD, CSS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	1/2722 (0.0%)	1.31	9/3690 (0.2%)
1	B	1.16	5/2845 (0.2%)	1.26	9/3857 (0.2%)
All	All	1.15	6/5567 (0.1%)	1.28	18/7547 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	29	THR	C-N	-17.69	1.10	1.33
1	B	46[A]	LEU	N-CA	7.19	1.55	1.46
1	B	46[B]	LEU	N-CA	7.19	1.55	1.46
1	B	271	ILE	C-O	-6.86	1.16	1.24
1	B	65	GLN	CD-OE1	6.45	1.35	1.23
1	A	172	PRO	C-O	-5.60	1.17	1.23

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	24	LEU	CA-C-N	10.13	130.81	120.38
1	A	24	LEU	C-N-CA	10.13	130.81	120.38
1	B	29	THR	O-C-N	-9.35	110.92	122.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	65	GLN	CG-CD-NE2	-8.50	103.65	116.40
1	B	46[A]	LEU	CA-C-O	-8.42	109.02	119.38
1	B	46[B]	LEU	CA-C-O	-8.42	109.02	119.38
1	A	285[A]	VAL	CA-C-O	8.25	129.40	120.57
1	A	285[B]	VAL	CA-C-O	8.25	129.40	120.57
1	A	53	GLY	CA-C-O	-6.73	117.20	122.52
1	A	229	ILE	O-C-N	6.45	125.44	121.37
1	B	46[A]	LEU	N-CA-C	5.83	119.65	112.54
1	B	46[B]	LEU	N-CA-C	5.83	119.65	112.54
1	B	65	GLN	OE1-CD-NE2	5.50	128.10	122.60
1	B	305	TYR	CA-CB-CG	5.49	123.78	113.90
1	A	305	TYR	CA-CB-CG	5.35	123.53	113.90
1	A	333	ASP	CA-CB-CG	5.30	117.90	112.60
1	B	47	GLU	CB-CG-CD	5.26	121.55	112.60
1	A	274	LYS	N-CA-C	-5.16	105.74	111.36

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	286[A]	VAL	Mainchain
1	B	46[A]	LEU	Mainchain
1	B	46[B]	LEU	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2663	0	2652	15	0
1	B	2781	0	2768	11	0
2	A	44	0	26	0	0
2	B	44	0	26	0	0
3	B	8	0	14	0	0
4	B	11	0	8	0	0
5	A	247	0	0	5	1
5	B	236	0	0	1	1
All	All	6034	0	5494	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:27:GLU:O	5:B:502:HOH:O	1.97	0.82
1:A:77:LEU:CD1	1:A:274:LYS:HE3	2.14	0.77
1:A:295:LEU:HD12	1:A:295:LEU:O	1.95	0.66
1:A:77:LEU:HD13	1:A:274:LYS:HE3	1.77	0.65
1:A:103:TYR:N	1:A:286[A]:VAL:HG11	2.12	0.65
1:A:137[B]:GLU:OE1	5:A:501:HOH:O	2.15	0.64
1:B:25:PRO:O	1:B:28:SER:HB2	1.97	0.64
1:B:14:LEU:C	1:B:14:LEU:HD13	2.30	0.57
1:A:261:GLU:HG2	5:A:505:HOH:O	2.05	0.56
1:A:359:MET:O	1:A:359:MET:HE3	2.04	0.56
1:B:14:LEU:HD11	1:B:18:LEU:HD11	1.91	0.52
1:A:275:CYS:HB2	5:A:714:HOH:O	2.12	0.49
1:A:275:CYS:SG	1:A:302:TYR:HE2	2.37	0.48
1:B:19:LYS:O	1:B:22:SER:OG	2.31	0.47
1:A:222:TYR:OH	1:A:226:LYS:NZ	2.47	0.46
5:A:541:HOH:O	1:B:296:MET:HE2	2.17	0.44
1:A:359:MET:HE3	1:A:359:MET:HA	1.99	0.43
1:B:103:TYR:N	1:B:286[A]:VAL:HG11	2.34	0.43
1:A:131:THR:CG2	1:A:285[B]:VAL:CG2	2.97	0.42
1:B:27:GLU:H	1:B:27:GLU:CD	2.26	0.42
1:B:122:HIS:CE1	1:B:359[B]:MET:HE1	2.54	0.41
1:A:305:TYR:OH	5:A:502:HOH:O	2.22	0.41
1:A:164:ASN:HB3	1:B:327:TRP:CE2	2.56	0.40
1:B:123:ASN:OD1	1:B:227:ASN:HB3	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:706:HOH:O	5:B:710:HOH:O[4_555]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	336/369 (91%)	332 (99%)	4 (1%)	0	100	100
1	B	355/369 (96%)	350 (99%)	4 (1%)	1 (0%)	36	17
All	All	691/738 (94%)	682 (99%)	8 (1%)	1 (0%)	48	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	26	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/309 (93%)	285 (99%)	3 (1%)	68	39
1	B	301/309 (97%)	300 (100%)	1 (0%)	86	74
All	All	589/618 (95%)	585 (99%)	4 (1%)	76	53

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	LEU
1	A	109	SER
1	A	115	THR
1	B	115	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	GLN
1	B	310	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSS	B	177	1	4,6,7	0.72	0	2,6,8	1.05	0
1	CSS	A	177	1	4,6,7	0.62	0	2,6,8	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	B	177	1	-	0/1/5/7	-
1	CSS	A	177	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	2AQ	B	403	-	12,12,12	1.96	3 (25%)	15,16,16	0.88	1 (6%)
2	NAD	B	402	-	46,48,48	0.54	1 (2%)	64,73,73	0.62	1 (1%)
3	MRD	B	401	-	7,7,7	0.25	0	9,10,10	0.75	0
2	NAD	A	401	-	46,48,48	0.49	0	64,73,73	0.62	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	2AQ	B	403	-	-	-	0/2/2/2
2	NAD	B	402	-	-	6/30/62/62	0/5/5/5
3	MRD	B	401	-	-	0/5/5/5	-
2	NAD	A	401	-	-	6/30/62/62	0/5/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	403	2AQ	C5-C10	-4.73	1.35	1.42
4	B	403	2AQ	C2-N11	-3.05	1.30	1.33
4	B	403	2AQ	C2-N1	2.69	1.42	1.35
2	B	402	NAD	C2N-N1N	2.08	1.37	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	402	NAD	C6N-N1N-C2N	-2.65	119.62	121.88
2	A	401	NAD	C6N-N1N-C2N	-2.40	119.84	121.88
4	B	403	2AQ	C3-C4-C5	-2.10	117.70	120.84

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	NAD	C5D-O5D-PN-O3
2	A	401	NAD	C5D-O5D-PN-O1N
2	B	402	NAD	C5D-O5D-PN-O3
2	B	402	NAD	C5D-O5D-PN-O1N
2	A	401	NAD	PA-O3-PN-O5D
2	B	402	NAD	PA-O3-PN-O5D
2	A	401	NAD	C5D-O5D-PN-O2N
2	B	402	NAD	C5D-O5D-PN-O2N
2	B	402	NAD	PA-O3-PN-O2N
2	A	401	NAD	C4D-C5D-O5D-PN
2	B	402	NAD	C4D-C5D-O5D-PN
2	A	401	NAD	PA-O3-PN-O2N

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 2AQ B 403



Ligand NAD B 402





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	29:THR	C	30:GLN	N	1.10

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	334/369 (90%)	0.70	29 (8%) 16 17	12, 29, 54, 129	7 (2%)
1	B	351/369 (95%)	0.97	39 (11%) 10 12	14, 30, 51, 83	7 (1%)
All	All	685/738 (92%)	0.83	68 (9%) 13 14	12, 29, 52, 129	14 (2%)

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	12	GLY	12.6
1	B	18	LEU	10.8
1	B	24	LEU	10.4
1	B	17	VAL	10.1
1	B	13	ALA	9.9
1	B	15	ALA	9.9
1	B	25	PRO	9.7
1	B	22	SER	8.9
1	B	19	LYS	8.8
1	B	23	THR	8.8
1	B	16	ALA	8.7
1	B	20	HIS	8.4
1	B	14	LEU	8.4
1	A	24	LEU	7.9
1	B	28	SER	7.6
1	B	21	SER	7.3
1	B	27	GLU	7.1
1	B	26	PRO	6.7
1	A	87	LEU	6.7
1	A	26	PRO	6.4
1	A	91	ARG	6.1
1	B	89	GLN	6.1
1	A	77	LEU	5.8
1	A	275	CYS	5.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	29	THR	5.3
1	A	84	HIS	5.3
1	A	23	THR	4.5
1	B	250	TYR	4.4
1	B	85	ALA	4.3
1	A	25	PRO	4.3
1	A	85	ALA	4.1
1	B	251	LYS	4.0
1	A	361	ALA	4.0
1	B	94	LEU	4.0
1	A	363	MET	3.9
1	A	86	ASP	3.5
1	A	274	LYS	3.5
1	B	90	SER	3.4
1	B	192	MET	3.2
1	A	362	PHE	3.1
1	A	27	GLU	3.0
1	B	88	THR	3.0
1	A	305	TYR	2.9
1	A	285[A]	VAL	2.9
1	B	363	MET	2.7
1	A	272	PHE	2.7
1	B	93	PRO	2.6
1	A	195	ASN	2.5
1	A	360	ASP	2.5
1	A	93	PRO	2.5
1	A	74	LEU	2.4
1	B	171	VAL	2.4
1	B	200	LYS	2.4
1	B	305	TYR	2.4
1	B	87	LEU	2.3
1	B	153	LEU	2.3
1	B	174	GLU	2.3
1	B	71	GLU	2.3
1	B	196	THR	2.3
1	B	91	ARG	2.2
1	B	75	GLU	2.2
1	A	271	ILE	2.1
1	A	332	VAL	2.1
1	A	298	ASN	2.1
1	B	256	VAL	2.0
1	A	76	PRO	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	199	VAL	2.0
1	A	295	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSS	B	177	7/8	0.90	0.14	36,40,55,96	0
1	CSS	A	177	7/8	0.93	0.10	28,30,43,56	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	2AQ	B	403	11/11	0.74	0.21	20,20,20,20	11
3	MRD	B	401	8/8	0.89	0.21	46,48,56,58	8
2	NAD	A	401	44/44	0.97	0.07	19,23,32,42	0
2	NAD	B	402	44/44	0.97	0.07	19,24,33,41	44




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around 2AQ B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD A 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.