wwPDB X-ray Structure Validation Summary Report (i)

Aug 20, 2023 – 01:38 PM EDT

PDB ID	:	2ISZ
Title	:	Crystal structure of a two-domain IdeR-DNA complex crystal form I
Authors	:	Wisedchaisri, G.; Chou, C.J.; Wu, M.; Roach, C.; Rice, A.E.; Holmes, R.K.; Beeson, C.; Hol, W.G.
Deposited on	:	2006-10-18
Resolution	:	2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	33	45% 45% 9%
2	F	33	9% 45% 45% 9%
3	A	157	73% 14% • 11%
3	B	157	% 71% 16% • 11%
3	C	157	3% 75% 13% • 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	157	

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5944 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called mbtA/mbtB operator strand 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	E	33	670	321	117	200	32	0	0	0

- Molecule 2 is a DNA chain called mbtA/mbtB operator strand 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	F	33	677	322	131	192	32	0	0	0

- Molecule 3 is a protein called Iron-dependent repressor ideR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	140	1112	692	204	210	6	0	0	0
3	B	140	1112	692	204	210	6	0	0	0
3	C	140	1104	685	203	210	6	0	0	0
3	D	140	1112	692	204	210	6	0	0	0

There are 68 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	141	ALA	-	expression tag	UNP P0A672
A	142	SER	-	expression tag	UNP P0A672
A	143	GLU	-	expression tag	UNP P0A672
A	144	ASN	-	expression tag	UNP P0A672
A	145	LEU	-	expression tag	UNP P0A672
A	146	TYR	-	expression tag	UNP P0A672
A	147	PHE	-	expression tag	UNP P0A672
A	148	GLN	-	expression tag	UNP P0A672

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	149	GLY	-	expression tag	UNP P0A672
A	150	GLY	-	expression tag	UNP P0A672
A	151	GLY	-	expression tag	UNP P0A672
A	152	HIS	-	expression tag	UNP P0A672
A	153	HIS	-	expression tag	UNP P0A672
A	154	HIS	-	expression tag	UNP P0A672
A	155	HIS	-	expression tag	UNP P0A672
A	156	HIS	-	expression tag	UNP P0A672
A	157	HIS	-	expression tag	UNP P0A672
B	141	ALA	-	expression tag	UNP P0A672
B	142	SER	-	expression tag	UNP P0A672
B	143	GLU	-	expression tag	UNP P0A672
B	144	ASN	-	expression tag	UNP P0A672
B	145	LEU	-	expression tag	UNP P0A672
B	146	TYR	-	expression tag	UNP P0A672
B	147	PHE	-	expression tag	UNP P0A672
B	148	GLN	-	expression tag	UNP P0A672
B	149	GLY	-	expression tag	UNP P0A672
B	150	GLY	-	expression tag	UNP P0A672
B	151	GLY	-	expression tag	UNP P0A672
B	152	HIS	-	expression tag	UNP P0A672
B	153	HIS	-	expression tag	UNP P0A672
B	154	HIS	-	expression tag	UNP P0A672
B	155	HIS	-	expression tag	UNP P0A672
B	156	HIS	-	expression tag	UNP P0A672
B	157	HIS	-	expression tag	UNP P0A672
C	141	ALA	-	expression tag	UNP P0A672
C	142	SER	-	expression tag	UNP P0A672
C	143	GLU	-	expression tag	UNP P0A672
C	144	ASN	-	expression tag	UNP P0A672
C	145	LEU	-	expression tag	UNP P0A672
C	146	TYR	-	expression tag	UNP P0A672
C	147	PHE	-	expression tag	UNP P0A672
C	148	GLN	-	expression tag	UNP P0A672
C	149	GLY	-	expression tag	UNP P0A672
C	150	GLY	-	expression tag	UNP P0A672
C	151	GLY	-	expression tag	UNP P0A672
C	152	HIS	-	expression tag	UNP P0A672
C	153	HIS	-	expression tag	UNP P0A672
C	154	HIS	-	expression tag	UNP P0A672
C	155	HIS	-	expression tag	UNP P0A672
C	156	HIS	-	expression tag	UNP P0A672

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	157	HIS	-	expression tag	UNP P0A672
D	141	ALA	-	expression tag	UNP P0A672
D	142	SER	-	expression tag	UNP P0A672
D	143	GLU	-	expression tag	UNP P0A672
D	144	ASN	-	expression tag	UNP P0A672
D	145	LEU	-	expression tag	UNP P0A672
D	146	TYR	-	expression tag	UNP P0A672
D	147	PHE	-	expression tag	UNP P0A672
D	148	GLN	-	expression tag	UNP P0A672
D	149	GLY	-	expression tag	UNP P0A672
D	150	GLY	-	expression tag	UNP P0A672
D	151	GLY	-	expression tag	UNP P0A672
D	152	HIS	-	expression tag	UNP P0A672
D	153	HIS	-	expression tag	UNP P0A672
D	154	HIS	-	expression tag	UNP P0A672
D	155	HIS	-	expression tag	UNP P0A672
D	156	HIS	-	expression tag	UNP P0A672
D	157	HIS	-	expression tag	UNP P0A672

- Molecule 4 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ni		
4	A	3	3	3	0	0
4	B	3	3	3	0	0
4	C	3	3	3	0	0
4	D	3	3	3	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
5	A	1	1	1	0	0
5	C	1	1	1	0	0
5	D	1	1	1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	E	25	25	25	0	0
			Total	O		
6	F	18	18	18	0	0
			Total	O		
6	A	30	30	30	0	0
			Total	O		
6	B	27	27	27	0	0
			Total	O		
6	C	24	24	24	0	0
			Total	O		
6	D	18	18	18	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: mbtA/mbtB operator strand 1

Chain E:



• Molecule 2: mbtA/mbtB operator strand 2

Chain F:



• Molecule 3: Iron-dependent repressor ideR

Chain A:



• Molecule 3: Iron-dependent repressor ideR

Chain B:



• Molecule 3: Iron-dependent repressor ideR

Chain C:



• Molecule 3: Iron-dependent repressor ideR

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	53.83Å 69.74Å 76.46Å 106.57° 104.85° 99.66°	Depositor
Resolution (Å)	50.00 – 2.40 48.84 – 2.40	Depositor EDS
% Data completeness (in resolution range)	90.0 (50.00-2.40) 90.1 (48.84-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.82 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.199 , 0.239 0.200 , 0.236	Depositor DCC
R_{free} test set	1754 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å^2)	35.8	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 44.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	5944	wwPDB-VP
Average B, all atoms (Å^2)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.32% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	E	0.66	0/749	1.45	12/1154 (1.0%)
2	F	0.68	0/761	1.32	9/1173 (0.8%)
3	A	0.37	0/1129	0.53	0/1530
3	B	0.36	0/1129	0.52	0/1530
3	C	0.35	0/1119	0.51	0/1515
3	D	0.37	0/1129	0.52	0/1530
All	All	0.46	0/6016	0.85	21/8432 (0.2%)

There are no bond length outliers.

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	F	12	DG	O4'-C1'-N9	9.20	114.44	108.00
1	E	20	DC	O4'-C4'-C3'	-9.08	100.55	106.00
1	E	1	DC	O4'-C4'-C3'	-9.00	100.60	106.00
2	F	23	DG	O4'-C4'-C3'	-8.63	100.82	106.00
2	F	31	DG	O4'-C1'-N9	7.97	113.58	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	670	0	375	13	0
2	F	677	0	371	9	0
3	A	1112	0	1128	14	0
3	B	1112	0	1127	17	0
3	C	1104	0	1122	14	0
3	D	1112	0	1128	16	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
4	C	3	0	0	0	0
4	D	3	0	0	0	0
5	A	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	A	30	0	0	0	0
6	B	27	0	0	2	0
6	C	24	0	0	2	0
6	D	18	0	0	2	0
6	E	25	0	0	4	0
6	F	18	0	0	0	0
All	All	5944	0	5251	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 79 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:102:CYS:SG	6:B:1029:HOH:O	2.31	0.88
3:C:80:ARG:HG2	3:C:126:SER:HB2	1.63	0.81
3:B:111:ASP:OD1	3:B:115:ARG:NH1	2.18	0.76
3:D:18:LEU:HD22	3:D:23:VAL:CG2	2.20	0.72
1:E:8:DA:H5"	6:E:48:HOH:O	1.93	0.68

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	138/157 (88%)	131 (95%)	7 (5%)	0	100	100
3	B	138/157 (88%)	135 (98%)	3 (2%)	0	100	100
3	C	138/157 (88%)	131 (95%)	7 (5%)	0	100	100
3	D	138/157 (88%)	131 (95%)	7 (5%)	0	100	100
All	All	552/628 (88%)	528 (96%)	24 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	122/135 (90%)	116 (95%)	6 (5%)	25	40
3	B	122/135 (90%)	118 (97%)	4 (3%)	38	57
3	C	121/135 (90%)	113 (93%)	8 (7%)	16	26
3	D	122/135 (90%)	114 (93%)	8 (7%)	16	26
All	All	487/540 (90%)	461 (95%)	26 (5%)	22	37

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	71	LEU
3	C	138	LEU
3	D	130	ASN
3	C	111	ASP
3	D	23	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	B	122	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 15 ligands modelled in this entry, 15 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	E	33/33 (100%)	-0.23	0 100 100	24, 45, 77, 80	0
2	F	33/33 (100%)	0.03	3 (9%) 9 8	23, 37, 95, 102	0
3	A	140/157 (89%)	-0.32	0 100 100	25, 33, 50, 62	0
3	B	140/157 (89%)	-0.02	2 (1%) 75 73	25, 34, 52, 61	0
3	C	140/157 (89%)	-0.08	4 (2%) 51 50	27, 39, 59, 67	0
3	D	140/157 (89%)	-0.18	1 (0%) 87 86	28, 39, 55, 67	0
All	All	626/694 (90%)	-0.14	10 (1%) 72 70	23, 37, 59, 102	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	140	VAL	8.4
3	D	140	VAL	4.1
2	F	1	DC	4.0
3	B	94	TRP	3.1
3	B	1	MET	2.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	NI	C	1001	1/1	0.91	0.09	55,55,55,55	0
4	NI	B	1003	1/1	0.92	0.14	52,52,52,52	0
4	NI	D	1003	1/1	0.93	0.12	58,58,58,58	0
5	NA	D	2001	1/1	0.93	0.07	49,49,49,49	0
5	NA	C	2001	1/1	0.94	0.14	48,48,48,48	0
4	NI	D	1001	1/1	0.94	0.08	56,56,56,56	0
4	NI	A	1003	1/1	0.95	0.22	45,45,45,45	0
4	NI	A	1001	1/1	0.97	0.04	49,49,49,49	0
4	NI	B	1001	1/1	0.97	0.06	49,49,49,49	0
4	NI	C	1003	1/1	0.98	0.25	53,53,53,53	0
5	NA	A	2001	1/1	0.98	0.10	50,50,50,50	0
4	NI	C	1002	1/1	0.99	0.12	36,36,36,36	0
4	NI	A	1002	1/1	0.99	0.16	28,28,28,28	0
4	NI	B	1002	1/1	0.99	0.16	29,29,29,29	0
4	NI	D	1002	1/1	0.99	0.16	34,34,34,34	0

6.5 Other polymers (i)

There are no such residues in this entry.