Full wwPDB X-ray Structure Validation Report (i)

Aug 7, 2023 – 03:18 AM EDT

PDB ID	:	1JFQ
Title	:	ANTIGEN-BINDING FRAGMENT OF THE MURINE ANTI-PHENYLARSONATE ANTIBODY 36-71, "FAB 36-71"
Authors	:	Parhami-Seren, B.; Viswanathan, M.; Strong, R.K.; Margolies, M.N.
Deposited on	:	2001-06-21
Resolution	:	1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	215	78% 19% •
2	H	222	65% 29% 5% •

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3447 atoms, of which 15 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ANTIGEN-BINDING FRAGMENT OF ANTI-PHENYLARSONATE ANTIBODY.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	L	215	1665	1027	5	284	342	7	0	0	0

- Molecule 2 is a protein called ANTIGEN-BINDING FRAGMENT OF ANTI-PHENYLARSONATE ANTIBODY.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	H	220	1670	1046	10	277	330	7	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	L	60	60	60	0	0
3	H	52	52	52	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: ANTIGEN-BINDING FRAGMENT OF ANTI-PHENYLARSONATE ANTIBODY

Chain L:



• Molecule 2: ANTIGEN-BINDING FRAGMENT OF ANTI-PHENYLARSONATE ANTIBODY

Chain H:



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	67.13Å 73.05Å 46.78Å 90.00° 104.48° 90.00°	Depositor
Resolution (Å)	6.00 – 1.90	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-1.90)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.196 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3447	wwPDB-VP
Average B, all atoms ($Å^2$)	19.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#|Z| >5	RMSZ	#|Z| >5
1	L	0.64	0/1693	0.86	0/2295
2	H	0.61	0/1702	0.87	2/2322 (0.1%)
All	All	0.62	0/3395	0.86	2/4617 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	H	485	LEU	CA-CB-CG	7.12	131.67	115.30
2	H	344	GLY	N-CA-C	6.10	128.35	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1660	5	1589	36	0
2	H	1660	10	1616	41	0
3	H	52	0	0	1	0
3	L	60	0	0	1	0
All	All	3432	15	3205	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:148:TRP:HB2	1:L:155:ARG:HB3	1.28	1.15
2:H:492:PRO:O	2:H:495:PRO:HD2	1.74	0.87
2:H:384:ARG:HD3	3:H:657:HOH:O	1.78	0.82
2:H:461:THR:HG22	2:H:504:ASN:HB2	1.64	0.79
2:H:494:SER:HB2	2:H:495:PRO:HD3	1.66	0.78
2:H:350:TYR:HB3	2:H:359:THR:HG22	1.69	0.73
2:H:335:ASN:HD22	2:H:347:TRP:HE1	1.38	0.71
1:L:2:ILE:HD13	1:L:29:ILE:HD11	1.74	0.68
1:L:147:LYS:HA	1:L:155:ARG:HD3	1.76	0.67
1:L:2:ILE:HG21	1:L:29:ILE:HD11	1.78	0.65
2:H:306:GLN:O	2:H:307:SER:HB2	1.95	0.65
1:L:147:LYS:HG3	1:L:154:GLU:HG3	1.81	0.63
1:L:147:LYS:HD3	1:L:149:LYS:HE3	1.81	0.62
2:H:310:GLU:HB3	2:H:416:THR:O	1.99	0.61
1:L:189:HIS:O	1:L:211:ARG:HD3	1.99	0.61
2:H:434:PRO:HB3	2:H:439:GLN:HB3	1.83	0.61
1:L:7:ILE:HB	1:L:8:PRO:HD3	1.85	0.58
1:L:36:TYR:HB3	1:L:44:ILE:HD12	1.85	0.58
1:L:154:GLU:HG2	1:L:155:ARG:N	2.19	0.58
2:H:353:PRO:HB3	2:H:372:VAL:HG21	1.85	0.57
1:L:154:GLU:HG2	1:L:155:ARG:H	1.73	0.54
1:L:147:LYS:HD3	1:L:149:LYS:CE	2.38	0.53
2:H:361:ASN:HB3	2:H:364:PHE:HD1	1.74	0.53
1:L:44:ILE:HG12	2:H:411:TRP:CD2	2.46	0.51
1:L:148:TRP:H	1:L:155:ARG:HG2	1.76	0.50
2:H:313:ARG:HA	2:H:313:ARG:HE	1.76	0.50
2:H:338:LYS:HE2	2:H:340:ARG:HG2	1.93	0.50
2:H:427:PRO:HB3	2:H:453:TYR:HB3	1.95	0.49
2:H:449:LEU:HD11	2:H:451:LYS:HD2	1.95	0.49
1:L:2:ILE:HG21	1:L:29:ILE:CD1	2.41	0.48
2:H:340:ARG:HD2	2:H:391:SER:O	2.13	0.48
2:H:312:VAL:HG11	2:H:318:VAL:CG1	2.44	0.48
1:L:44:ILE:HD11	2:H:411:TRP:CZ2	2.49	0.48
1:L:135:PHE:CE1	2:H:488:SER:HB3	2.49	0.47
1:L:149:LYS:HA	1:L:153:SER:O	2.14	0.47
1:L:155:ARG:CG	1:L:156:GLN:H	2.27	0.47
2:H:313:ARG:CZ	2:H:314:ALA:H	2.28	0.47
1:L:148:TRP:H	1:L:155:ARG:CG	2.27	0.47
2:H:443:MET:HB3	2:H:490:THR:CG2	2.45	0.47
2:H:444:VAL:O	2:H:490:THR:HA	2.15	0.47
1:L:184:ASP:O	1:L:188:ARG:HG3	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:21:ILE:O	1:L:72:SER:HA	2.16	0.46
2:H:313:ARG:NE	2:H:314:ALA:H	2.13	0.46
1:L:150:ILE:HD11	1:L:179:LEU:HD11	1.97	0.46
2:H:350:TYR:HB3	2:H:359:THR:CG2	2.40	0.46
2:H:453:TYR:CE2	2:H:458:VAL:HG13	2.50	0.46
1:L:167:ASP:OD1	1:L:169:LYS:HG2	2.16	0.46
1:L:184:ASP:OD1	1:L:188:ARG:NH1	2.48	0.46
1:L:210:ASN:ND2	1:L:212:ASN:OD1	2.49	0.46
2:H:308:GLY:HA3	2:H:320:MET:HA	1.98	0.45
2:H:468:SER:O	2:H:471:VAL:HG22	2.17	0.45
2:H:478:LEU:HD13	2:H:483:TYR:CE1	2.50	0.45
1:L:118:PHE:HA	1:L:119:PRO:HD3	1.80	0.44
2:H:480:SER:O	2:H:481:ASP:HB2	2.16	0.44
2:H:441:ASN:O	2:H:442:SER:HB2	2.19	0.43
2:H:318:VAL:HG22	2:H:386:LEU:HD11	2.00	0.43
1:L:50:PHE:HB2	2:H:402:TYR:CE2	2.53	0.43
2:H:338:LYS:HB2	2:H:348:ILE:HD11	2.00	0.43
1:L:212:ASN:O	1:L:214:CYS:N	2.51	0.43
1:L:163:TRP:HH2	3:L:623:HOH:O	2.01	0.43
2:H:376:SER:O	2:H:377:ASN:HB2	2.19	0.42
1:L:142:LYS:HE3	1:L:163:TRP:CD1	2.54	0.42
1:L:169:LYS:HB3	1:L:169:LYS:HE3	1.87	0.42
1:L:85:THR:HA	1:L:102:THR:O	2.20	0.42
2:H:367:LYS:HA	2:H:384:ARG:HH21	1.83	0.42
2:H:384:ARG:CG	2:H:384:ARG:HH21	2.32	0.42
1:L:83:ILE:HD11	1:L:106:ILE:HG13	2.01	0.41
2:H:400:GLU:HG3	2:H:409:ASP:OD2	2.21	0.41
2:H:340:ARG:HD2	2:H:392:ALA:HB2	2.03	0.41
1:L:6:GLN:HA	1:L:22:SER:O	2.20	0.41
2:H:339:GLN:HB2	2:H:345:LEU:HD12	2.03	0.40
2:H:352:ASN:HB3	2:H:355:ASN:OD1	2.21	0.40
1:L:12:SER:HB3	1:L:107:LYS:HE2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	L	213/215 (99%)	205 (96%)	6 (3%)	2 (1%)	17 7
2	H	218/222 (98%)	200 (92%)	12 (6%)	6 (3%)	5 1
All	All	431/437 (99%)	405 (94%)	18 (4%)	8 (2%)	8 1

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	155	ARG
2	H	307	SER
2	H	343	GLN
2	H	403	GLY
2	H	436	SER
2	H	495	PRO
1	L	156	GLN
2	H	438	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	L	190/191 (100%)	178 (94%)	12 (6%)	18 8
2	H	188/190 (99%)	169 (90%)	19 (10%)	7 2
All	All	378/381 (99%)	347 (92%)	31 (8%)	11 4

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	29	ILE
1	L	44	ILE
1	L	46	LEU
1	L	47	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	L	53	ARG
1	L	72	SER
1	L	74	THR
1	L	147	LYS
1	L	160	LEU
1	L	168	SER
1	L	179	LEU
1	L	190	ASN
2	H	303	GLN
2	H	307	SER
2	H	310	GLU
2	H	313	ARG
2	H	345	LEU
2	H	361	ASN
2	H	375	SER
2	H	384	ARG
2	H	388	SER
2	H	421	SER
2	H	440	THR
2	H	441	ASN
2	H	449	LEU
2	H	457	PRO
2	H	461	THR
2	H	479	GLN
2	H	485	LEU
2	H	496	ARG
2	H	511	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	34	ASN
1	L	55	GLN
1	L	77	ASN
1	L	161	ASN
1	L	189	HIS
2	H	335	ASN
2	H	361	ASN
2	H	439	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.