



Full wwPDB EM Validation Report ⓘ

Nov 24, 2025 – 04:43 PM JST

PDB ID : 9JG1 / pdb_00009jg1
EMDB ID : EMD-61447
Title : Cryo-EM structure of Adriformant-bound Histamine receptor 4 H4R at inactive state
Authors : Jin, S.S.; Zhang, H.; Jiang, Y.
Deposited on : 2024-09-05
Resolution : 3.62 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev129
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	210492	15764	-
Ramachandran outliers	207382	16835	-
Sidechain outliers	206894	16415	-
Q-score	-	25397	11773 (3.12 - 4.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	R	427	
2	H	222	
3	L	226	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	A1EBW	R	701	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4730 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histamine H4 receptor,Soluble cytochrome b562.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	386	Total	C	N	O	S	0	0
			3026	1996	494	525	11		

There are 37 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	378	ALA	-	linker	UNP Q9H3N8
R	379	ARG	-	linker	UNP Q9H3N8
R	380	ARG	-	linker	UNP Q9H3N8
R	381	GLN	-	linker	UNP Q9H3N8
R	382	LEU	-	linker	UNP Q9H3N8
R	389	TRP	MET	conflict	UNP P0ABE7
R	484	ILE	HIS	conflict	UNP P0ABE7
R	488	LEU	-	linker	UNP P0ABE7
R	489	GLU	-	linker	UNP P0ABE7
R	490	ARG	-	linker	UNP P0ABE7
R	491	ALA	-	linker	UNP P0ABE7
R	492	ARG	-	linker	UNP P0ABE7
R	493	SER	-	linker	UNP P0ABE7
R	494	THR	-	linker	UNP P0ABE7
R	495	LEU	-	linker	UNP P0ABE7
R	576	PHE	-	expression tag	UNP Q9H3N8
R	577	LEU	-	expression tag	UNP Q9H3N8
R	578	LYS	-	expression tag	UNP Q9H3N8
R	579	ILE	-	expression tag	UNP Q9H3N8
R	580	PHE	-	expression tag	UNP Q9H3N8
R	581	CYS	-	expression tag	UNP Q9H3N8
R	582	ILE	-	expression tag	UNP Q9H3N8
R	583	LYS	-	expression tag	UNP Q9H3N8
R	584	LYS	-	expression tag	UNP Q9H3N8
R	585	GLN	-	expression tag	UNP Q9H3N8
R	586	PRO	-	expression tag	UNP Q9H3N8
R	587	LEU	-	expression tag	UNP Q9H3N8
R	588	PRO	-	expression tag	UNP Q9H3N8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
R	589	SER	-	expression tag	UNP Q9H3N8
R	590	GLN	-	expression tag	UNP Q9H3N8
R	591	HIS	-	expression tag	UNP Q9H3N8
R	592	SER	-	expression tag	UNP Q9H3N8
R	593	ARG	-	expression tag	UNP Q9H3N8
R	594	SER	-	expression tag	UNP Q9H3N8
R	595	VAL	-	expression tag	UNP Q9H3N8
R	596	SER	-	expression tag	UNP Q9H3N8
R	597	SER	-	expression tag	UNP Q9H3N8


- Molecule 2 is a protein called anti-BRIL Fab Heavy chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	H	106	Total	C	N	O	S	0	0
			787	494	131	159	3		

- Molecule 3 is a protein called anti-BRIL Fab Light chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	L	121	Total	C	N	O	S	0	0
			898	573	152	170	3		

- Molecule 4 is Adriforant (CCD ID: A1EBW) (formula: $C_{13}H_{22}N_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
			Total	C	N	
4	R	1	19	13	6	0

THR	THR
ALA	ALA
ASN	ASN
VAL	VAL
LEU	LEU
GLY	GLY
ASN	ASN
CYS	CYS
LEU	LEU
VAL	VAL
LYS	LYS
ASP	ASP
THR	THR
TYR	TYR
PHE	PHE
PRO	PRO
GLU	GLU
PRO	PRO
VAL	VAL
THR	THR
VAL	VAL
SER	SER
TRP	TRP
ASN	ASN
SER	SER
GLY	GLY
ALA	ALA
LEU	LEU
THR	THR
SER	SER
GLY	GLY
VAL	VAL
HIS	HIS
THR	THR
PHE	PHE
PRO	PRO
ALA	ALA
VAL	VAL
LEU	LEU
GLN	GLN
SER	SER
SER	SER
GLY	GLY
LEU	LEU
TYR	TYR
SER	SER
LEU	LEU
SER	SER
VAL	VAL
VAL	VAL
THR	THR
VAL	VAL
PRO	PRO
SER	SER
SER	SER
SER	SER
LEU	LEU
GLY	GLY
THR	THR
GLN	GLN
THR	THR
TYR	TYR

ILE
CYS
ASN
VAL
HIS
LYS
PRO
SER
ASN
THR
LYS
VAL
ASP
LYS
LYS
VAL
GLU
PRO

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	132359	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TECNAI ARCTICA	Depositor
Voltage (kV)	50	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	1.459	Depositor
Minimum map value	-0.972	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.033	Depositor
Recommended contour level	0.19	Depositor
Map size (\AA)	233.6, 233.6, 233.6	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.73, 0.73, 0.73	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1EBW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	R	0.28	1/3107 (0.0%)	0.53	5/4244 (0.1%)
2	H	0.26	0/804	0.61	1/1094 (0.1%)
3	L	0.25	0/924	0.67	2/1261 (0.2%)
All	All	0.27	1/4835 (0.0%)	0.57	8/6599 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	R	541	GLY	C-N	5.10	1.38	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	R	541	GLY	CA-C-N	-6.69	114.01	120.83
1	R	541	GLY	C-N-CA	-6.69	114.01	120.83
1	R	464	GLY	N-CA-C	-6.08	106.85	115.43
1	R	413	THR	N-CA-C	-5.33	105.37	111.07
3	L	105	TRP	CA-C-N	-5.25	114.17	119.83
3	L	105	TRP	C-N-CA	-5.25	114.17	119.83
1	R	473	ALA	N-CA-C	-5.10	104.53	111.56
2	H	54	SER	N-CA-C	5.01	117.73	109.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	3026	0	2965	45	0
2	H	787	0	751	13	0
3	L	898	0	825	9	0
4	R	19	0	0	12	0
All	All	4730	0	4541	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:95:TYR:HB3	4:R:701:A1EBW:N13	1.61	1.14
1:R:95:TYR:HB3	4:R:701:A1EBW:C14	1.95	0.96
2:H:30:VAL:CG1	2:H:33:ALA:HB3	1.98	0.93
1:R:554:GLN:HG2	4:R:701:A1EBW:C04	2.13	0.78
1:R:68:SER:HA	1:R:90:TRP:HE1	1.55	0.72
2:H:30:VAL:HG11	2:H:33:ALA:HB3	1.75	0.68
2:H:17:GLY:H	2:H:79:LEU:HB2	1.60	0.67
1:R:182:GLU:HG3	4:R:701:A1EBW:N10	2.10	0.67
1:R:116:VAL:HG11	1:R:197:ILE:HG23	1.77	0.67
1:R:554:GLN:CB	4:R:701:A1EBW:C04	2.73	0.67
1:R:554:GLN:CG	4:R:701:A1EBW:C04	2.72	0.67
1:R:46:LEU:HD22	1:R:571:ARG:HH21	1.60	0.66
1:R:59:ILE:O	1:R:62:PHE:HB3	1.97	0.65
1:R:554:GLN:HG2	4:R:701:A1EBW:C05	2.27	0.64
3:L:42:GLN:HB2	3:L:48:LEU:HD23	1.79	0.64
1:R:61:ASP:O	1:R:64:VAL:HB	1.98	0.63
1:R:100:ALA:O	1:R:104:ASN:ND2	2.31	0.61
2:H:86:THR:HG22	2:H:105:LYS:HG2	1.83	0.61
1:R:551:PHE:CE1	4:R:701:A1EBW:N02	2.69	0.60
2:H:90:GLN:HE22	2:H:92:TYR:HB3	1.68	0.58
1:R:551:PHE:HE1	4:R:701:A1EBW:N02	2.01	0.58
3:L:76:ASP:OD2	3:L:78:SER:OG	2.20	0.58
3:L:89:LEU:HD13	3:L:125:VAL:HG22	1.87	0.56
2:H:3:ILE:HD12	2:H:91:GLN:HE22	1.69	0.56
1:R:95:TYR:CB	4:R:701:A1EBW:N13	2.52	0.56
2:H:30:VAL:HG13	2:H:33:ALA:HB3	1.87	0.54
3:L:64:ALA:HB3	3:L:67:VAL:HG22	1.89	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:530:THR:O	1:R:534:SER:OG	2.25	0.54
1:R:381:GLN:HE22	1:R:427:PRO:HA	1.73	0.53
1:R:169:PHE:HA	1:R:175:LEU:HD11	1.89	0.52
2:H:65:GLY:HA2	2:H:73:THR:O	2.10	0.52
3:L:94:THR:HG23	3:L:124:THR:HA	1.92	0.51
1:R:74:PRO:HA	1:R:78:PHE:HB2	1.91	0.51
1:R:166:PRO:HB2	1:R:168:PHE:HD1	1.76	0.51
1:R:139:VAL:O	1:R:142:LEU:HB2	2.11	0.50
1:R:551:PHE:HE1	4:R:701:A1EBW:C01	2.24	0.50
3:L:43:ALA:HB3	3:L:46:LYS:HB2	1.94	0.50
2:H:90:GLN:HE21	2:H:98:VAL:HG13	1.77	0.50
2:H:22:ILE:HD12	2:H:74:LEU:HD23	1.92	0.50
1:R:30:MET:HG2	1:R:62:PHE:HE1	1.77	0.48
1:R:166:PRO:HB2	1:R:168:PHE:CD1	2.49	0.48
1:R:113:TYR:O	1:R:117:SER:OG	2.31	0.47
1:R:80:TRP:NE1	1:R:82:PHE:O	2.47	0.47
1:R:439:GLU:O	1:R:487:TYR:OH	2.28	0.47
1:R:545:VAL:HG22	1:R:548:ARG:HH21	1.80	0.46
1:R:136:MET:O	1:R:139:VAL:HB	2.15	0.46
1:R:103:TYR:HB3	1:R:139:VAL:HG13	1.97	0.46
1:R:162:SER:HB3	1:R:165:GLU:HB2	1.98	0.45
3:L:105:TRP:HA	3:L:110:TRP:HB2	1.98	0.45
1:R:554:GLN:HB3	4:R:701:A1EBW:C04	2.47	0.45
1:R:389:TRP:HE1	1:R:480:ARG:NH1	2.16	0.44
1:R:92:THR:HA	1:R:95:TYR:CE2	2.53	0.44
2:H:48:LEU:HA	2:H:59:VAL:HG21	2.01	0.43
1:R:416:ARG:HB3	1:R:416:ARG:HH11	1.83	0.43
1:R:178:THR:O	1:R:179:SER:C	2.62	0.42
1:R:92:THR:HA	1:R:95:TYR:CZ	2.54	0.42
1:R:103:TYR:OH	1:R:146:VAL:HG21	2.20	0.42
1:R:187:VAL:HG22	1:R:516:LEU:HD21	2.02	0.42
1:R:443:PHE:HB2	1:R:487:TYR:CE1	2.55	0.42
3:L:35:PHE:HB3	3:L:101:ARG:HG3	2.01	0.41
1:R:107:LEU:HD11	1:R:135:LEU:HD13	2.01	0.41
1:R:570:LYS:HD3	1:R:570:LYS:HA	1.91	0.41
1:R:473:ALA:HA	1:R:475:GLN:HE22	1.86	0.41
1:R:526:TYR:HB2	1:R:554:GLN:NE2	2.36	0.41
2:H:90:GLN:NE2	2:H:92:TYR:HB3	2.35	0.40
3:L:35:PHE:CD2	3:L:101:ARG:HD2	2.56	0.40
2:H:89:CYS:O	2:H:101:GLY:N	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	R	382/427 (90%)	370 (97%)	12 (3%)	0	100	100
2	H	104/222 (47%)	101 (97%)	3 (3%)	0	100	100
3	L	119/226 (53%)	116 (98%)	3 (2%)	0	100	100
All	All	605/875 (69%)	587 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	R	309/376 (82%)	308 (100%)	1 (0%)	91	96
2	H	86/196 (44%)	85 (99%)	1 (1%)	67	82
3	L	88/187 (47%)	88 (100%)	0	100	100
All	All	483/759 (64%)	481 (100%)	2 (0%)	88	94

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	R	92	THR
2	H	46	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	R	48	HIS
1	R	75	HIS
1	R	147	ASN
1	R	206	HIS
1	R	381	GLN
1	R	485	GLN
1	R	569	HIS
1	R	573	GLN
2	H	4	GLN
2	H	39	GLN
2	H	90	GLN
2	H	102	GLN
3	L	42	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	A1EBW	R	701	-	21,21,21	4.58	8 (38%)	22,29,29	4.09	11 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	A1EBW	R	701	-	-	4/9/22/22	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	R	701	A1EBW	C04-C03	-10.51	1.25	1.52
4	R	701	A1EBW	C19-N06	-8.82	1.35	1.46
4	R	701	A1EBW	C09-N10	8.40	1.50	1.33
4	R	701	A1EBW	C19-C03	6.90	1.66	1.53
4	R	701	A1EBW	C12-N13	6.86	1.46	1.36
4	R	701	A1EBW	C07-N06	5.72	1.49	1.37
4	R	701	A1EBW	C05-N06	5.02	1.54	1.47
4	R	701	A1EBW	C04-C05	3.69	1.59	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	R	701	A1EBW	N08-C07-N06	11.31	129.24	116.55
4	R	701	A1EBW	C18-C07-N06	-8.38	112.33	122.29
4	R	701	A1EBW	C09-N08-C07	6.85	121.58	116.73
4	R	701	A1EBW	N10-C09-N08	4.94	124.94	117.25
4	R	701	A1EBW	N13-C12-N11	4.55	124.09	116.43
4	R	701	A1EBW	N11-C09-N08	-4.25	118.75	125.42
4	R	701	A1EBW	C14-N13-C12	-3.82	116.37	123.36
4	R	701	A1EBW	C09-N11-C12	3.17	121.88	114.68
4	R	701	A1EBW	C18-C12-N13	-3.12	115.76	120.86
4	R	701	A1EBW	C05-N06-C19	-2.76	104.42	111.55
4	R	701	A1EBW	C18-C07-N08	-2.42	118.43	123.15

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	R	701	A1EBW	N13-C14-C15-C16
4	R	701	A1EBW	C18-C07-N06-C05
4	R	701	A1EBW	N08-C07-N06-C05
4	R	701	A1EBW	N13-C14-C15-C17

There are no ring outliers.

1 monomer is involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	R	701	A1EBW	12	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-61447. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 153




Y Index: 167




Z Index: 214

6.3.2 Raw map



X Index: 152



Y Index: 186




Z Index: 214


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.19. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 58 nm^3 ; this corresponds to an approximate mass of 52 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.276 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.276 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.62	-	-
Author-provided FSC curve	3.62	4.24	3.85
Unmasked-calculated*	4.39	7.28	4.57

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.39 differs from the reported value 3.62 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-61447 and PDB model 9JG1. Per-residue inclusion information can be found in section [3](#) on page [7](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.19 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.19).

9.4 Atom inclusion [i](#)



At the recommended contour level, 88% of all backbone atoms, 71% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.19) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.7130	<div></div> 0.3430
H	<div></div> 0.8250	<div></div> 0.4470
L	<div></div> 0.7730	<div></div> 0.4060
R	<div></div> 0.6670	<div></div> 0.2970

