Full wwPDB X-ray Structure Validation Report (i)

Nov 1, 2023 – 03:22 PM JST

PDB ID	:	5JHM
Title	:	Crystal structure of Zika virus Envelope protein
Authors	:	Dai, L.; Shi, Y.; Qi, J.; Gao, G.F.
Deposited on	:	2016-04-21
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	416	6% 74% 17% • 6%
1	B	416	3% 79% 14% • 6%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 6236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	391	2988	1867	521	575	25	0	1	0
1	B	390	2975	1860	519	571	25	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	128	128	128	0	0
2	B	145	145	145	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Envelope protein

Chain A:



- Molecule 1: Envelope protein

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	76.45Å 59.06Å 103.67Å 90.00° 103.72° 90.00°	Depositor
Resolution (Å)	33.98 – 2.00 33.98 – 1.99	Depositor EDS
% Data completeness (in resolution range)	92.7 (33.98-2.00) 85.3 (33.98-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.94 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.208 , 0.237 0.213 , 0.240	Depositor DCC
R_{free} test set	2901 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å^2)	24.4	Xtriage
Anisotropy	0.571	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 41.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	6236	wwPDB-VP
Average B, all atoms (Å^2)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.34	0/3050	0.59	2/4130 (0.0%)
1	B	0.33	0/3037	0.55	0/4112
All	All	0.33	0/6087	0.57	2/8242 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	232	GLY	N-CA-C	-5.29	99.87	113.10
1	A	228	GLY	C-N-CA	5.20	134.70	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2988	0	2926	53	0
1	B	2975	0	2915	35	0
2	A	128	0	0	4	0
2	B	145	0	0	7	0
All	All	6236	0	5841	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:27:HIS:ND1	2:B:502:HOH:O	1.99	0.92
1:A:67:ASP:OD2	1:A:90:TYR:OH	1.93	0.85
1:A:274:GLU:HB3	1:A:275:ALA:HB2	1.58	0.83
1:B:64:SER:OG	2:B:501:HOH:O	1.96	0.82
1:A:37:ASP:OD2	2:A:501:HOH:O	1.97	0.81
1:B:244:GLU:OE2	1:B:246:LYS:HE2	1.85	0.76
1:A:253:GLN:OE1	2:A:502:HOH:O	2.03	0.76
1:B:277:MET:SD	2:B:595:HOH:O	2.45	0.75
1:A:74:CYS:HB2	1:A:77:GLN:HG3	1.73	0.70
1:A:205:THR:HG23	2:A:544:HOH:O	1.92	0.69
1:A:228:GLY:HA2	1:A:229:ALA:CB	2.24	0.68
1:B:246:LYS:O	2:B:505:HOH:O	2.14	0.66
1:A:276:GLU:OE2	1:A:285:SER:HB3	1.97	0.64
1:A:170:THR:HG22	1:A:172:ASN:H	1.64	0.62
1:A:2:ARG:NE	1:A:44:GLU:OE2	2.23	0.62
1:A:194:THR:HG21	1:A:290:LYS:H	1.66	0.60
1:B:254:THR:N	2:B:505:HOH:O	2.16	0.60
1:B:14:GLY:O	1:B:15:MET:HB2	2.01	0.60
1:B:343:ALA:O	2:B:506:HOH:O	2.16	0.60
1:A:138:ARG:NH2	1:A:168:GLU:OE2	2.35	0.59
1:A:14:GLY:N	1:A:34:MET:O	2.31	0.58
1:A:166:LYS:NZ	2:A:504:HOH:O	2.37	0.57
1:A:228:GLY:HA2	1:A:229:ALA:HB3	1.86	0.56
1:B:308:CYS:HB3	1:B:332:TYR:CZ	2.40	0.56
1:A:135:LEU:HD13	1:A:198:PHE:HZ	1.71	0.56
1:A:274:GLU:HB3	1:A:275:ALA:CB	2.34	0.55
1:B:359:ILE:HD11	1:B:379:ASP:HB2	1.88	0.55
1:A:198:PHE:O	1:A:200:ASP:N	2.31	0.55
1:A:13:GLU:OE2	1:A:357:ARG:NH1	2.40	0.54
1:A:22:ASP:OD1	1:A:292:ARG:NH1	2.41	0.54
1:A:348:ASP:OD2	1:A:351:THR:HG22	2.09	0.53
1:B:139:ILE:HD12	1:B:169:ILE:HD12	1.90	0.52
1:A:194:THR:CG2	1:A:290:LYS:H	2.22	0.52
1:A:274:GLU:OE2	1:B:246:LYS:HE3	2.10	0.51
1:B:74:CYS:HB2	1:B:77:GLN:HG3	1.93	0.51
1:B:403:SER:OG	1:B:404:GLY:N	2.44	0.50
1:A:142:SER:HB3	1:A:164:ARG:HG2	1.92	0.50
1:A:247:ASP:OD2	1:A:249:HIS:CE1	2.65	0.50
1:A:274:GLU:CB	1:A:275:ALA:HB2	2.37	0.50
1:B:301:LYS:HB2	2:B:518:HOH:O	2.13	0.49
1:A:308:CYS:HB3	1:A:332:TYR:CZ	2.47	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:GLU:CD	1:A:357:ARG:HH12	2.17	0.48
1:B:20:TRP:HA	1:B:293:LEU:O	2.13	0.48
1:B:263:GLY:O	1:B:267:THR:HG23	2.13	0.48
1:B:142:SER:HB3	1:B:164:ARG:HG2	1.95	0.48
1:B:341:VAL:HB	1:B:363:PRO:HG2	1.95	0.48
1:A:229:ALA:HA	1:A:230:ASP:HB3	1.95	0.47
1:B:338:PRO:HA	1:B:365:ILE:O	2.13	0.47
1:A:27:HIS:HD2	1:A:287:GLY:HA3	1.79	0.46
1:B:138:ARG:NH1	1:B:168:GLU:OE2	2.48	0.45
1:A:130:ILE:HD11	1:A:203:TYR:HB2	1.98	0.45
1:A:228:GLY:HA2	1:A:229:ALA:HB2	1.98	0.45
1:A:15:MET:HG3	1:A:18:GLY:HA3	1.98	0.45
1:B:115:THR:OG1	1:B:253:GLN:HB2	2.16	0.45
1:A:217:TRP:CH2	1:A:269:LEU:HD21	2.51	0.45
1:B:223:LEU:HD21	1:B:261:GLN:HG3	1.99	0.44
1:B:164:ARG:HG2	1:B:164:ARG:HH11	1.83	0.44
1:A:71:ASP:OD2	1:A:73:ARG:NH1	2.51	0.44
1:A:247:ASP:O	1:A:248:ALA:CB	2.66	0.43
1:B:175:ARG:HA	1:B:188:LEU:O	2.17	0.43
1:A:359:ILE:HD11	1:A:379:ASP:HB2	2.01	0.43
1:B:14:GLY:HA2	1:B:21:VAL:CG1	2.48	0.43
1:B:136:GLU:OE2	1:B:168:GLU:HG2	2.18	0.43
1:B:138:ARG:HH12	1:B:168:GLU:CD	2.21	0.43
1:A:279:GLY:C	1:A:281:LYS:H	2.21	0.43
1:A:341:VAL:HG22	1:A:365:ILE:HD12	1.99	0.43
1:A:132:PRO:O	1:A:171:PRO:HG2	2.19	0.42
1:A:351:THR:HG23	1:A:353:THR:HG23	2.01	0.42
1:B:57:ARG:NH1	1:B:59:TYR:OH	2.52	0.42
1:B:14:GLY:HA2	1:B:21:VAL:HG11	2.00	0.42
1:A:175:ARG:HA	1:A:175:ARG:HD2	1.61	0.42
1:A:14:GLY:HA3	1:A:35:ALA:HB2	2.01	0.42
1:A:194:THR:HG21	1:A:290:LYS:HB2	2.02	0.42
1:A:20:TRP:HA	1:A:293:LEU:O	2.20	0.41
1:B:13:GLU:HG2	1:B:357:ARG:HH12	1.85	0.41
1:B:314:PHE:CZ	1:B:396:ILE:HG21	2.55	0.41
1:A:202:TYR:CE1	1:A:215:LYS:HE2	2.55	0.41
1:A:205:THR:HG22	1:A:210:HIS:ND1	2.36	0.41
1:A:276:GLU:HG3	1:A:283:ARG:O	2.19	0.41
1:A:278:ASP:HA	1:A:279:GLY:HA2	1.70	0.41
1:B:132:PRO:O	1:B:171:PRO:HG2	2.20	0.41
1:B:1:ILE:HD13	1:B:1:ILE:HG21	1.88	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:PRO:HD3	1:A:300:LEU:HD13	2.03	0.40
1:A:196:LEU:HD23	1:A:198:PHE:HE1	1.86	0.40
1:A:191:GLU:O	1:A:194:THR:HG22	2.22	0.40
1:A:210:HIS:O	1:A:275:ALA:HA	2.21	0.40
1:B:278:ASP:HA	1:B:279:GLY:HA2	1.81	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	388/416 (93%)	366 (94%)	8 (2%)	14 (4%)	3 1
1	B	386/416 (93%)	366 (95%)	13 (3%)	7 (2%)	8 3
All	All	774/832 (93%)	732 (95%)	21 (3%)	21 (3%)	5 1

All (21) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	229	ALA
1	A	231	THR
1	A	248	ALA
1	A	272	ALA
1	A	276	GLU
1	B	15	MET
1	B	231	THR
1	A	14	GLY
1	A	15	MET
1	A	16	SER
1	A	275	ALA
1	B	14	GLY
1	B	232	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	199	SER
1	A	230	ASP
1	A	247	ASP
1	B	67	ASP
1	B	403	SER
1	A	236	TRP
1	A	280	ALA
1	B	248	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	327/348 (94%)	319 (98%)	8 (2%)	49	51
1	B	325/348 (93%)	319 (98%)	6 (2%)	59	63
All	All	652/696 (94%)	638 (98%)	14 (2%)	55	57

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	THR
1	A	185[A]	SER
1	A	185[B]	SER
1	A	191	GLU
1	A	276	GLU
1	A	293	LEU
1	A	300	LEU
1	A	365	ILE
1	B	49	THR
1	B	84	LYS
1	B	174	PRO
1	B	292	ARG
1	B	300	LEU
1	B	328	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	253	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	391/416 (93%)	0.39	23 (5%)	22	21	16, 37, 74, 124	0
1	B	390/416 (93%)	0.29	14 (3%)	42	42	19, 31, 67, 133	0
All	All	781/832 (93%)	0.34	37 (4%)	31	30	16, 34, 70, 133	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	405	SER	10.6
1	B	231	THR	6.7
1	A	196	LEU	6.6
1	B	278	ASP	5.8
1	B	229	ALA	5.6
1	A	278	ASP	5.4
1	B	404	GLY	4.7
1	B	247	ASP	4.4
1	B	230	ASP	4.1
1	B	232	GLY	4.0
1	A	231	THR	3.9
1	A	198	PHE	3.9
1	B	20	TRP	3.8
1	A	15	MET	3.6
1	A	194	THR	3.4
1	A	193	ARG	3.3
1	B	233	THR	3.2
1	A	230	ASP	3.1
1	B	16	SER	3.0
1	A	248	ALA	2.9
1	A	235	HIS	2.8
1	A	246	LYS	2.8
1	B	370	GLU	2.7
1	A	247	ASP	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	14	GLY	2.6
1	A	162	GLU	2.5
1	A	350	GLN	2.5
1	A	197	ASP	2.5
1	B	18	GLY	2.4
1	A	276	GLU	2.2
1	A	406	THR	2.2
1	A	287	GLY	2.2
1	A	199	SER	2.2
1	A	175	ARG	2.2
1	B	162	GLU	2.1
1	A	16	SER	2.0
1	A	232	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.