Full wwPDB EM Validation Report (i)

Nov 14, 2022 – 05:08 PM JST

PDB ID	:	6JHS
EMDB ID	:	EMD-9829
Title	:	The cryo-EM structure of HAV bound to a neutralizing antibody-F7
Authors	:	Cao, L.; Liu, P.; Yang, P.; Gao, Q.; Li, H.; Sun, Y.; Zhu, L.; Lin, J.; Su, D.; Rao, Z.; Wang, X.
Deposited on	:	2019-02-19
Resolution	:	3.05 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev43
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	278	70% 85% 9% 6%
2	B	222	68% 94% · ·
3	C	246	67% 90% 10%
4	D	213	77% 86% 14%
5	E	221	90% 84% 14% ·

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8978 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VP1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	260	2032	1298	332	395	7	0	0

- Molecule 2 is a protein called VP2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	216	1708	1098	291	315	4	0	0

- Molecule 3 is a protein called VP3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	246	1956	1248	331	363	14	0	0

- Molecule 4 is a protein called FAB Light Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	D	213	1641	1022	274	336	9	0	0

- Molecule 5 is a protein called FAB Heavy Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	E	217	1641	1042	268	322	9	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: VP1

Chain A: 70% 85% 9% 6%



- Molecule 2: VP2

Chain B: 68% 94%





- Molecule 3: VP3

Chain C:





- Molecule 4: FAB Light Chain

Chain D:

- Molecule 5: FAB Heavy Chain

Chain E:




D62 S63 V64 K65 G66 R67 F68 T69 I70 S71 R72 D73 N74 A75 K76 N77 T78 L79 Y80 L81 Q82 M83 S84 S85 L86 R87 S88 G89 D90 T91 A92 M93 Y94 Y95 C96 A97 R98 R99 D100 N101 T102 T103 A104 Q105 Y106 Y107 F108 D109 Y110 W111 G112 Q113 G114 T115 T116 L117 T118 V119 S120 S121
P122 K123 T124 T125 P126 P127 S128 V129 Y130 P131 L132 A133 P134 A135 SER ALA SER THR A140 A141 S142 M143 V144 T145 L146 G147 C148 L149 V150 K151 G152 Y153 F154 P155 E156 P157 V158 T159 V160 T161 W162 N163 S164 G165 S166 L167 S168 S169 G170 V171 H172 T173 F174 P175 A176 V177 L178 Q179 S180 D181
L182 Y183 T184 L185 S186 S187 S188 V189 T190 V191 P192 S193 S194 T195 W196 P197 S198 E199 T200 V201 T202 C203 N204 V205 A206 H207 P208 A209 S210 S211 T212 K213 V214 D215 K216 K217 I218 V219 P220 R221

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	16743	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	1.3	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 BASE (4k x 4k)	Depositor
Maximum map value	0.482	Depositor
Minimum map value	-0.291	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.020	Depositor
Recommended contour level	0.035	Depositor
Map size (Å)	648.0, 648.0, 648.0	wwPDB
Map dimensions	480, 480, 480	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.35, 1.35, 1.35	Depositor

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.27	0/2087	0.53	0/2848
2	B	0.28	0/1752	0.53	0/2387
3	C	0.29	0/2007	0.55	1/2733 (0.0%)
4	D	0.28	0/1681	0.54	0/2285
5	E	0.28	0/1685	0.55	0/2305
All	All	0.28	0/9212	0.54	1/12558 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	D	0	1
5	E	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	C	141	LEU	CA-CB-CG	5.53	128.01	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	D	48	TYR	Peptide
5	E	121	SER	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2032	0	1955	17	0
2	B	1708	0	1692	7	0
3	C	1956	0	1911	17	0
4	D	1641	0	1559	15	0
5	E	1641	0	1601	16	0
All	All	8978	0	8718	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:118:ASP:HB2	3:C:225:TYR:HB2	1.74	0.69
5:E:39:GLN:HB3	5:E:93:MET:HB3	1.76	0.66
1:A:65:PRO:HB3	3:C:170:ARG:HE	1.60	0.66
5:E:91:THR:HG23	5:E:118:THR:HA	1.79	0.64
4:D:34:TRP:HB2	4:D:47:ILE:HB	1.81	0.62
4:D:89:GLN:HE21	4:D:96:THR:H	1.48	0.61
4:D:161:SER:HB3	4:D:175:SER:HB3	1.84	0.57
1:A:56:GLU:OE2	3:C:120:GLN:NE2	2.39	0.56
2:B:69:LEU:H	2:B:72:HIS:HD2	1.53	0.55
5:E:29:PHE:O	5:E:72:ARG:NH1	2.40	0.55
4:D:185:TYR:O	4:D:191:TYR:OH	2.26	0.54
4:D:19:VAL:HB	4:D:74:ILE:HB	1.90	0.54
3:C:121:VAL:HG12	3:C:222:VAL:HG22	1.90	0.54
2:B:191:ASN:OD1	3:C:221:ARG:NH2	2.40	0.53
1:A:195:THR:O	3:C:13:ASN:ND2	2.40	0.53
1:A:110:PRO:HD3	1:A:169:PRO:HB3	1.91	0.53
4:D:2:ILE:HG23	4:D:89:GLN:HE22	1.73	0.53
1:A:112:THR:HG22	1:A:114:SER:H	1.74	0.52
5:E:24:ALA:HB3	5:E:77:ASN:HB3	1.92	0.52
4:D:119:PRO:HG2	4:D:129:ALA:HB1	1.93	0.51
5:E:69:THR:HB	5:E:82:GLN:HB3	1.93	0.51
3:C:63:GLN:NE2	3:C:223:ASN:OD1	2.45	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:120:GLN:HG2	2:B:192:ILE:HG22	1.95	0.48
5:E:38:ARG:NH1	5:E:46:GLU:OE1	2.47	0.47
4:D:116:ILE:HG12	4:D:208:PHE:HE1	1.79	0.47
3:C:143:ASP:OD2	5:E:56:SER:OG	2.32	0.47
5:E:52:ASP:O	5:E:72:ARG:NH2	2.48	0.47
4:D:159:LEU:HB2	4:D:177:THR:HB	1.96	0.47
3:C:87:GLN:HA	3:C:243:VAL:HG13	1.96	0.46
5:E:40:THR:OG1	5:E:43:LYS:O	2.25	0.46
1:A:197:ASN:HD22	3:C:15:VAL:HG22	1.80	0.46
4:D:135:LEU:HB2	4:D:174:MET:HB3	1.98	0.45
4:D:29:VAL:O	4:D:70:TYR:OH	2.31	0.45
2:B:137:SER:HB2	3:C:98:ILE:HG22	1.99	0.45
5:E:44:GLY:HA2	5:E:45:LEU:HA	1.78	0.44
4:D:209:ASN:HB2	4:D:212:GLU:HB2	1.98	0.44
1:A:211:VAL:HG11	1:A:226:PHE:HB3	2.00	0.44
3:C:132:LEU:HB3	3:C:205:TYR:HB2	2.00	0.43
1:A:128:ARG:HH12	1:A:270:LEU:HD12	1.83	0.43
1:A:182:LYS:NZ	1:A:232:GLN:OE1	2.42	0.43
1:A:254:GLN:HE21	3:C:36:TRP:H	1.66	0.43
3:C:31:LEU:HD23	3:C:32:ASP:HB2	2.01	0.43
1:A:236:TYR:HA	1:A:237:ASN:HA	1.81	0.43
5:E:34:MET:HB3	5:E:79:LEU:HD11	2.00	0.42
1:A:261:ARG:NH1	2:B:132:ASP:O	2.44	0.42
2:B:69:LEU:H	2:B:72:HIS:CD2	2.36	0.42
1:A:145:ILE:HD11	1:A:198:ILE:HD13	2.01	0.42
5:E:170:GLY:HA3	5:E:190:THR:HB	2.01	0.42
4:D:60:ARG:NH1	4:D:76:SER:O	2.53	0.42
5:E:146:LEU:HD12	5:E:191:VAL:HG21	2.02	0.42
1:A:125:SER:HB2	3:C:238:TYR:CZ	2.55	0.41
2:B:192:ILE:HB	2:B:196:THR:HB	2.02	0.41
5:E:195:THR:O	5:E:199:GLU:N	2.54	0.41
4:D:181:THR:O	4:D:185:TYR:N	2.51	0.41
5:E:83:MET:HB3	5:E:86:LEU:HD21	2.02	0.41
1:A:72:LYS:HA	1:A:73:PRO:HD3	1.96	0.41
4:D:165:GLN:HB2	4:D:172:TYR:HE1	1.85	0.41
5:E:47:TRP:NE1	5:E:49:ALA:O	2.54	0.41
1:A:203:PRO:HD3	3:C:29:PHE:HB2	2.03	0.40
1:A:236:TYR:CD1	1:A:237:ASN:HB3	2.57	0.40
3:C:127:HIS:HB3	3:C:210:LEU:HD11	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/278 (92%)	245 (96%)	11 (4%)	0	100	100
2	B	214/222 (96%)	203 (95%)	11 (5%)	0	100	100
3	C	244/246 (99%)	236 (97%)	8 (3%)	0	100	100
4	D	211/213 (99%)	198 (94%)	13 (6%)	0	100	100
5	E	213/221 (96%)	201 (94%)	11 (5%)	1 (0%)	29	60
All	All	1138/1180 (96%)	1083 (95%)	54 (5%)	1 (0%)	54	81

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	E	122	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/242 (94%)	227 (100%)	0	100	100
2	B	188/194 (97%)	188 (100%)	0	100	100
3	C	217/217 (100%)	217 (100%)	0	100	100
4	D	186/187 (100%)	186 (100%)	0	100	100
5	E	185/189 (98%)	185 (100%)	0	100	100
All	All	1003/1029 (98%)	1003 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	254	GLN
1	A	265	ASN
2	B	72	HIS
2	B	76	HIS
2	B	111	GLN
3	C	87	GLN
3	C	95	GLN
4	D	89	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-9829. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)

6.1.1 Primary map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 240



Y Index: 240



Z Index: 240

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 276



Y Index: 173



Z Index: 249

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.035. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 5014 nm^3; this corresponds to an approximate mass of 4530 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.328 Å^{-1}

8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-9829 and PDB model 6JHS. Per-residue inclusion information can be found in section 3 on page 4.

9.1 Map-model overlays

9.1.1 Map-model overlay (i)



X



Y



Z

9.1.2 Map-model assembly overlay (i)



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.035 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.035).

9.4 Atom inclusion (i)



At the recommended contour level, 22% of all backbone atoms, 23% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.035) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.2309	-0.0150
A	0.2384	0.0080
B	0.2996	-0.0330
C	0.3115	-0.0200
D	0.1856	-0.0170
E	0.0993	-0.0140