Full wwPDB NMR Structure Validation Report (i)

Dec 13, 2023 – 07:52 PM EST

PDB ID	:	2JP7
Title	:	NMR structure of the Mex67 UBA domain
Authors	:	Hobeika, M.; Brockmann, C.; Iglesias, N.; Gwizdek, C.; Neuhaus, D.; Stutz, F.; Stewart, M.; Divita, G.; Dargemont, C.
Deposited on	:	2007-04-26

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	57	49% 39% 7% 5%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:544-A:597 (54)	0.47	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 6 single-model clusters were found.

Cluster number	Models
1	4, 5, 6, 8, 9, 11, 12, 14, 18, 19, 20
2	1, 3, 16
Single-model clusters	2; 7; 10; 13; 15; 17

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 938 atoms, of which 467 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called mRNA export factor MEX67.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	57	938	303	467	79	87	2	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: mRNA export factor MEX67

Chain A: 49% 39% 7% 5%

R543 V547 L555 E558 T559 K560 L561 N562 A563 T566 F567 M568 L569 Q572 N576 Y577 E578 V579 A580 I581 Q585 M588 I591 P592 R593 E594 A595 F596 V597 Q598 F599

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: mRNA export factor MEX67

Chain A: 47% 37% 9% • 5%

R543 L544 V547 Q548 L552 N553 K554 L555 E558 T559 K560 L561 N562 A563 E564 Y565 T566 F567 M568 L569 Q572 N576 Y577 E578 V579 A580 I581 Q585 P592 R593 E594 A595 F596 V597 Q598 F599

4.2.2 Score per residue for model 2

• Molecule 1: mRNA export factor MEX67

Chain A: 49% 40% • • 5%

R543 V547 L551 L555 H556 T559 K560 L561 N562 A563 E564 Y565 T566 F567 M588 L569 Q572 S573 N574 W575 N576 Y577 E578 V579 A580 I581 M588 I591 F596 V597 Q598 F599

4.2.3 Score per residue for model 3

- Molecule 1: mRNA export factor MEX67

Chain A: 49% 39% 5% • 5%

4.2.4 Score per residue for model 4

- Molecule 1: mRNA export factor MEX67

Chain A: 54% 37% • 5%



4.2.5 Score per residue for model 5 (medoid)

- Molecule 1: mRNA export factor MEX67

Chain A: 51% 35% 9% 5%

4.2.6 Score per residue for model 6

- Molecule 1: mRNA export factor MEX67

Chain A: 47% 37% 11% 5%



4.2.7 Score per residue for model 7

- Molecule 1: mRNA export factor MEX67

Chain A: 56% 32% 7% 5%

4.2.8 Score per residue for model 8

- Molecule 1: mRNA export factor MEX67

Chain A: 44% 47% • 5%



4.2.9 Score per residue for model 9

- Molecule 1: mRNA export factor MEX67

Chain A: 44% 46% • • 5%



4.2.10 Score per residue for model 10

- Molecule 1: mRNA export factor MEX67

Chain A: 58% 33% • • 5%

4.2.11 Score per residue for model 11

- Molecule 1: mRNA export factor MEX67

Chain A: 47% 42% • • 5%

4.2.12 Score per residue for model 12

- Molecule 1: mRNA export factor MEX67

Chain A: 46% 46% • 5%



4.2.13 Score per residue for model 13

- Molecule 1: mRNA export factor MEX67

Chain A: 49% 42% • 5%

4.2.14 Score per residue for model 14

- Molecule 1: mRNA export factor MEX67

Chain A: 40% 46% 7% • 5%



4.2.15 Score per residue for model 15

- Molecule 1: mRNA export factor MEX67

Chain A: 46% 42% 7% 5%



4.2.16 Score per residue for model 16

- Molecule 1: mRNA export factor MEX67

Chain A: 53% 35% 7% 5%



4.2.17 Score per residue for model 17

- Molecule 1: mRNA export factor MEX67

Chain A: 42% 44% 7% • 5%

4.2.18 Score per residue for model 18

- Molecule 1: mRNA export factor MEX67

Chain A: 40% 46% 9% 5%

R543 L544 V547 L552 L555 H556 T559 K560 L561 N562 A563 E564 Y565 T566 F567 A570 E571 Q572 N576 Y577 E578 V579 A580 I581 K582 G583 F584 Q585 M588 I591 P592 R593 E594 A595 F596 V597 Q598 F599

4.2.19 Score per residue for model 19

- Molecule 1: mRNA export factor MEX67

Chain A: 44% 46% 5% 5%

R543 L544 Q548 L551 K554 L555 E558 T559 K560 L561 N562 A563 T566 L569 S573 N574 W575 N576 Y577 E578 V579 A580 I581 F584 Q585 M588 I591 P592 R593 E594 A595 F596 V597 Q598 F599

4.2.20 Score per residue for model 20

- Molecule 1: mRNA export factor MEX67

Chain A: 51% 37% 5% • 5%

R543 V547 N553 K554 L555 E558 T559 K560 L561 N562 A563 T566 L569 S573 N576 Y577 E578 V579 A580 I581 K582 Q585 I591 P592 R593 E594 A595 F596 V597 Q598 F599

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	2.13
X-PLOR NIH	refinement	2.13

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.12±0.02	1±1/449 (0.2± 0.2%)	0.95±0.03	1±1/609 (0.1± 0.1%)
All	All	1.12	21/8980 (0.2%)	0.95	14/12180 (0.1%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	561	LEU	N-CA	-5.93	1.34	1.46	5	9
1	A	581	ILE	N-CA	-5.70	1.34	1.46	2	1
1	A	561	LEU	CA-C	-5.51	1.38	1.52	4	8
1	A	561	LEU	C-N	-5.30	1.21	1.34	10	1
1	A	566	THR	C-O	-5.10	1.13	1.23	6	2

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	585	GLN	CB-CA-C	5.92	122.23	110.40	11	7
1	A	597	VAL	CG1-CB-CG2	-5.34	102.35	110.90	5	2
1	A	585	GLN	CA-CB-CG	5.34	125.15	113.40	1	5

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	439	435	434	20±4
All	All	8780	8700	8680	394

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:552:LEU:HD13	1:A:567:PHE:HB3	0.79	1.54	15	4
1:A:563:ALA:HA	1:A:566:THR:HG22	0.78	1.54	18	6
1:A:562:ASN:HB2	1:A:595:ALA:HA	0.76	1.55	14	4
1:A:579:VAL:HA	1:A:582:LYS:HG2	0.75	1.57	10	1
1:A:573:SER:HB2	1:A:579:VAL:HG13	0.72	1.60	20	7
1:A:592:PRO:HB2	1:A:594:GLU:HG2	0.69	1.64	17	14
1:A:569:LEU:HD13	1:A:580:ALA:HB1	0.69	1.64	17	1
1:A:581:ILE:O	1:A:585:GLN:HB3	0.69	1.88	11	6
1:A:563:ALA:HA	1:A:566:THR:HB	0.67	1.67	17	14
1:A:561:LEU:HA	1:A:595:ALA:O	0.66	1.91	1	19
1:A:581:ILE:O	1:A:585:GLN:HG2	0.66	1.91	10	3
1:A:555:LEU:HD11	1:A:580:ALA:HB3	0.64	1.68	11	2
1:A:555:LEU:O	1:A:559:THR:HG22	0.64	1.92	10	19
1:A:578:GLU:HA	1:A:581:ILE:HB	0.63	1.69	2	18
1:A:581:ILE:O	1:A:585:GLN:HG3	0.63	1.94	6	3
1:A:582:LYS:HA	1:A:585:GLN:NE2	0.63	2.08	7	4
1:A:552:LEU:HD21	1:A:567:PHE:HA	0.62	1.70	6	1
1:A:558:GLU:HG3	1:A:577:TYR:HE1	0.62	1.53	1	1
1:A:562:ASN:CB	1:A:595:ALA:HA	0.62	2.25	9	6
1:A:558:GLU:HG3	1:A:577:TYR:CE1	0.61	2.30	1	2
1:A:581:ILE:O	1:A:585:GLN:HB2	0.61	1.96	16	2
1:A:559:THR:OG1	1:A:561:LEU:HG	0.61	1.96	9	19
1:A:588:MET:HA	1:A:591:ILE:HG12	0.61	1.72	12	5
1:A:564:GLU:HA	1:A:567:PHE:CE2	0.59	2.32	18	4
1:A:555:LEU:HD21	1:A:569:LEU:HD23	0.58	1.73	2	1
1:A:561:LEU:HD13	1:A:591:ILE:HG21	0.58	1.75	2	5
1:A:582:LYS:O	1:A:586:SER:HB3	0.58	1.98	16	2
1:A:576:ASN:HB3	1:A:579:VAL:CG1	0.57	2.29	1	8
1:A:563:ALA:CA	1:A:566:THR:HG22	0.57	2.29	18	5
1:A:558:GLU:HG3	1:A:577:TYR:OH	0.57	1.99	16	8
1:A:563:ALA:HA	1:A:566:THR:CB	0.57	2.29	2	6
1:A:584:PHE:HE1	1:A:596:PHE:HE1	0.57	1.42	17	1
1:A:588:MET:HB2	1:A:596:PHE:HZ	0.57	1.60	12	4
1:A:544:LEU:HD21	1:A:567:PHE:HE2	0.56	1.61	9	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:555:LEU:HD21	1:A:580:ALA:HB3	0.55	1.77	10	10
1:A:560:LYS:O	1:A:596:PHE:HA	0.55	2.00	8	2
1:A:555:LEU:HG	1:A:570:ALA:HB2	0.55	1.78	14	1
1:A:561:LEU:O	1:A:565:TYR:HB2	0.54	2.02	2	1
1:A:555:LEU:HD23	1:A:559:THR:HG22	0.54	1.79	11	2
1:A:578:GLU:C	1:A:581:ILE:HB	0.54	2.22	4	5
1:A:569:LEU:HA	1:A:572:GLN:HE21	0.54	1.62	5	2
1:A:565:TYR:CD2	1:A:595:ALA:HB2	0.54	2.38	11	3
1:A:567:PHE:O	1:A:571:GLU:HB3	0.53	2.04	3	1
1:A:593:ARG:HA	1:A:596:PHE:HB2	0.53	1.78	5	3
1:A:569:LEU:CD1	1:A:580:ALA:HB1	0.53	2.32	17	1
1:A:591:ILE:HB	1:A:596:PHE:CE2	0.52	2.39	11	2
1:A:554:LYS:O	1:A:558:GLU:HG2	0.52	2.04	10	3
1:A:595:ALA:O	1:A:597:VAL:HG13	0.52	2.05	1	11
1:A:556:HIS:HA	1:A:559:THR:OG1	0.52	2.04	2	1
1:A:544:LEU:HB3	1:A:548:GLN:HB2	0.52	1.82	7	4
1:A:551:LEU:HD11	1:A:577:TYR:HB2	0.51	1.83	12	1
1:A:588:MET:HB2	1:A:596:PHE:CZ	0.51	2.41	18	5
1:A:552:LEU:HD21	1:A:567:PHE:CA	0.51	2.34	6	1
1:A:555:LEU:HG	1:A:559:THR:HG23	0.50	1.83	2	1
1:A:573:SER:OG	1:A:580:ALA:HB2	0.50	2.05	2	9
1:A:562:ASN:ND2	1:A:595:ALA:HA	0.50	2.21	1	2
1:A:555:LEU:HD21	1:A:580:ALA:CB	0.50	2.36	10	9
1:A:578:GLU:CA	1:A:581:ILE:HB	0.50	2.35	4	5
1:A:576:ASN:HB2	1:A:579:VAL:HG12	0.50	1.83	12	1
1:A:551:LEU:CD1	1:A:577:TYR:HB2	0.50	2.37	12	1
1:A:569:LEU:HG	1:A:580:ALA:HB1	0.49	1.85	11	8
1:A:563:ALA:O	1:A:566:THR:HB	0.49	2.07	1	4
1:A:561:LEU:HD22	1:A:591:ILE:HG21	0.48	1.85	4	2
1:A:571:GLU:HA	1:A:575:TRP:NE1	0.48	2.22	9	1
1:A:562:ASN:HB2	1:A:597:VAL:CG1	0.48	2.38	19	4
1:A:551:LEU:HD23	1:A:575:TRP:HE3	0.48	1.68	14	9
1:A:565:TYR:O	1:A:568:MET:HG2	0.48	2.09	2	1
1:A:573:SER:HB2	1:A:579:VAL:HG22	0.48	1.85	19	1
1:A:556:HIS:O	1:A:560:LYS:HA	0.48	2.09	15	1
1:A:567:PHE:HB2	1:A:575:TRP:CH2	0.47	2.44	14	1
1:A:594:GLU:HA	1:A:597:VAL:HA	0.47	1.86	15	1
1:A:544:LEU:HD13	1:A:567:PHE:HE1	0.47	1.69	6	1
1:A:591:ILE:HB	1:A:596:PHE:CE1	0.47	2.45	9	2
1:A:556:HIS:HB2	1:A:566:THR:HG21	0.46	1.88	15	4
1:A:561:LEU:CD1	1:A:591:ILE:HG21	0.46	2.40	5	6

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:587:SER:O	1:A:591:ILE:HG12	0.45	2.11	7	1
1:A:593:ARG:HA	1:A:596:PHE:HB3	0.45	1.87	4	1
1:A:562:ASN:HB3	1:A:565:TYR:HD2	0.45	1.71	9	1
1:A:588:MET:HA	1:A:591:ILE:CG1	0.45	2.42	12	2
1:A:544:LEU:HD11	1:A:567:PHE:CE2	0.45	2.46	7	1
1:A:581:ILE:HA	1:A:584:PHE:CD2	0.45	2.47	14	1
1:A:552:LEU:HD12	1:A:566:THR:HG23	0.44	1.89	4	3
1:A:576:ASN:HB3	1:A:579:VAL:HG12	0.44	1.88	5	4
1:A:552:LEU:CD1	1:A:567:PHE:HB3	0.44	2.37	18	1
1:A:551:LEU:CD1	1:A:577:TYR:HA	0.44	2.43	2	1
1:A:554:LYS:HE3	1:A:577:TYR:CZ	0.44	2.47	15	1
1:A:562:ASN:HB2	1:A:597:VAL:HG11	0.43	1.89	19	2
1:A:555:LEU:HD23	1:A:570:ALA:HB2	0.43	1.90	18	1
1:A:592:PRO:HB2	1:A:594:GLU:CG	0.43	2.44	8	3
1:A:584:PHE:O	1:A:588:MET:HB3	0.43	2.14	18	3
1:A:588:MET:HA	1:A:596:PHE:CE2	0.43	2.48	15	1
1:A:591:ILE:HG22	1:A:595:ALA:HB3	0.43	1.90	10	1
1:A:552:LEU:HD12	1:A:566:THR:HG22	0.43	1.91	17	1
1:A:581:ILE:HD12	1:A:581:ILE:HA	0.42	1.56	6	1
1:A:576:ASN:HB3	1:A:579:VAL:HG11	0.42	1.90	1	1
1:A:567:PHE:O	1:A:571:GLU:HG2	0.42	2.14	8	2
1:A:556:HIS:ND1	1:A:561:LEU:O	0.42	2.51	11	1
1:A:545:ASN:HB2	1:A:546:PRO:HD2	0.42	1.92	5	1
1:A:567:PHE:O	1:A:571:GLU:HB2	0.42	2.14	9	1
1:A:563:ALA:CA	1:A:566:THR:HB	0.42	2.41	17	1
1:A:591:ILE:HB	1:A:596:PHE:HE1	0.42	1.73	9	1
1:A:582:LYS:HA	1:A:585:GLN:HE22	0.41	1.74	7	1
1:A:581:ILE:HA	1:A:581:ILE:HD13	0.41	1.39	12	2
1:A:568:MET:O	1:A:572:GLN:HG3	0.41	2.16	17	1
1:A:551:LEU:HD13	1:A:577:TYR:HA	0.41	1.92	2	1
1:A:582:LYS:NZ	1:A:582:LYS:HB2	0.41	2.29	3	1
1:A:581:ILE:HD13	1:A:581:ILE:HA	0.41	1.35	4	1
1:A:561:LEU:CD2	1:A:591:ILE:HG21	0.41	2.46	10	1
1:A:591:ILE:HB	1:A:596:PHE:HE2	0.41	1.75	7	1
1:A:563:ALA:HA	1:A:566:THR:OG1	0.41	2.15	2	1
1:A:574:ASN:OD1	1:A:576:ASN:HB2	0.41	2.16	5	1
1:A:569:LEU:CD2	1:A:580:ALA:HB1	0.41	2.46	8	1
1:A:563:ALA:O	1:A:567:PHE:HB2	0.40	2.15	2	1
1:A:566:THR:O	1:A:569:LEU:HB3	0.40	2.16	14	1
1:A:552:LEU:HD23	1:A:575:TRP:CZ3	0.40	2.50	6	1
1:A:561:LEU:HB2	1:A:566:THR:OG1	0.40	2.17	14	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	54/57 (95%)	48±2 (88±3%)	5±2 (9±3%)	1±0 (3±1%)	8 43
All	All	1080/1140 (95%)	953 (88%)	98 (9%)	29 (3%)	8 43

All 7 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	597	VAL	20
1	A	562	ASN	3
1	A	577	TYR	2
1	A	589	ASN	1
1	A	574	ASN	1
1	A	592	PRO	1
1	A	588	MET	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	48/51 (94%)	43±1 (90±3%)	5±1 (10±3%)	12 58
All	All	960/1020 (94%)	868 (90%)	92 (10%)	12 58

All 19 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	581	ILE	15
1	A	585	GLN	14
1	A	547	VAL	11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	572	GLN	11
1	A	560	LYS	8
1	A	576	ASN	7
1	A	553	ASN	6
1	A	588	MET	4
1	A	582	LYS	3
1	A	574	ASN	3
1	A	562	ASN	2
1	A	569	LEU	1
1	A	593	ARG	1
1	A	557	LEU	1
1	A	545	ASN	1
1	A	592	PRO	1
1	A	555	LEU	1
1	A	564	GLU	1
1	A	571	GLU	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided