wwPDB NMR Structure Validation Summary Report (i)

Oct 17, 2021 – 10:05 AM EDT

PDB ID	:	1K3M
Title	:	NMR STRUCTURE OF HUMAN INSULIN MUTANT ILE-A2-ALA, HIS-B10-ASP, PRO-B28-LYS, LYS-B29-PRO, 15 STRUCTURES
Authors	:	Xu, B.; Hua, Q.-X.; Nakagawa, S.H.; Jia, W.; Chu, Y.-C.; Katsoyannis, P.G.; Weiss, M.A.
Deposited on	:	2001-10-03

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.23.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	21	10% 38% 24% 5% 24%
2	B	30	7% 47% 17% 30%

2 Ensemble composition and analysis (i)

This entry contains 15 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:6-A:21, B:5-B:25 (37)	0.36	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

NmrClust was unable to cluster the ensemble.

Error message: Inconsistent models

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 772 atoms, of which 372 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called INSULIN.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	21	303	96	143	25	35	4	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	ILE	engineered mutation	UNP P01308

- Molecule 2 is a protein called INSULIN.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
2	B	30	469	156	229	38	44	2	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	10	ASP	HIS	engineered mutation	UNP P01308
B	28	LYS	PRO	engineered mutation	UNP P01308
B	29	PRO	LYS	engineered mutation	UNP P01308

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: INSULIN

Chain A: 10% 38% 24% 5% 24%

G1 A2 V3 E4 Q5 C6 C7 T8 S9 I10 C11 S12 L13 Y14 Q15 L16 E17 N18 Y19 C20 N21

- Molecule 2: INSULIN

Chain B: 7% 47% 17% 30%

F1 V2 N3 Q4 H5 L6 C7 G8 S9 D10 L11 V12 E13 A14 L15 Y16 L17 V18 C19 G20 E21 R22 G23 F24 F25 Y26 T27 K28 P29 T30

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 8. Colouring as in section 4.1 above.

- Molecule 1: INSULIN

Chain A: 10% 38% 24% 5% 24%

G1 A2 V3 E4 Q5 C6 C7 T8 S9 I10 C11 S12 L13 Y14 Q15 L16 E17 N18 Y19 C20 N21

- Molecule 2: INSULIN

Chain B: 7% 37% 27% 30%

F1 V2 N3 Q4 H5 L6 C7 G8 S9 D10 L11 V12 E13 A14 L15 Y16 L17 V18 C19 G20 E21 R22 G23 F24 F25 Y26 T27 K28 P29 T30

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *distance geometry and simulated annealing.*

Of the 60 calculated structures, 15 were deposited, based on the following criterion: *structures with the lowest energy.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.85
DGII	refinement	InsightII 2000

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
2	B	0.0±0.0	1.0±0.0
All	All	0	15

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
2	B	22	ARG	Sidechain	15

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	126	110	110	20±5
2	B	162	152	152	29±5
All	All	4320	3930	3930	570

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 69.

5 of 188 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:9:SER:O	1:A:10:ILE:HD13	0.95	1.61	1	8

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:13:LEU:HD11	2:B:18:VAL:HG22	0.90	1.44	14	4
2:B:6:LEU:HD21	2:B:14:ALA:CB	0.87	2.00	11	3
1:A:13:LEU:O	1:A:13:LEU:HD13	0.80	1.75	7	4
1:A:16:LEU:HD11	2:B:6:LEU:HD23	0.77	1.54	7	5

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	15/21 (71%)	9±1 (58±8%)	3±1 (22±8%)	3±1 (20±4%)	0 2
2	B	21/30 (70%)	16±1 (75±7%)	4±1 (17±7%)	2±1 (8±5%)	2 14
All	All	540/765 (71%)	368 (68%)	103 (19%)	69 (13%)	1 6

5 of 11 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	9	SER	15
1	A	11	CYS	15
2	B	8	GLY	14
1	A	13	LEU	10
2	B	20	GLY	4

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	16/19 (84%)	10±1 (65±6%)	6±1 (35±6%)	1 10
2	B	17/26 (65%)	11±1 (64±9%)	6±1 (36±9%)	1 8

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	495/675 (73%)	321 (65%)	174 (35%)	1 9

5 of 26 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	10	ILE	15
1	A	16	LEU	15
2	B	7	CYS	15
2	B	18	VAL	14
2	B	16	TYR	12

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided