Full wwPDB X-ray Structure Validation Report (i)

Sep 20, 2023 – 10:21 PM EDT

PDB ID	:	5KIK
Title	:	CmlA beta-hydroxylase in chemically reduced diferrous state
Authors	:	Knoot, C.J.; Lipscomb, J.D.
Deposited on	:	2016-06-16
Resolution	:	2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	551	

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 4402 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CmlA protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	521	4209	2669	752	775	13	0	2	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP F2RB80
A	-18	GLY	-	expression tag	UNP F2RB80
A	-17	SER	-	expression tag	UNP F2RB80
A	-16	SER	-	expression tag	UNP F2RB80
A	-15	HIS	-	expression tag	UNP F2RB80
A	-14	HIS	-	expression tag	UNP F2RB80
A	-13	HIS	-	expression tag	UNP F2RB80
A	-12	HIS	-	expression tag	UNP F2RB80
A	-11	HIS	-	expression tag	UNP F2RB80
A	-10	SER	-	expression tag	UNP F2RB80
A	-9	SER	-	expression tag	UNP F2RB80
A	-8	GLY	-	expression tag	UNP F2RB80
A	-7	LEU	-	expression tag	UNP F2RB80
A	-6	VAL	-	expression tag	UNP F2RB80
A	-5	PRO	-	expression tag	UNP F2RB80
A	-4	ARG	-	expression tag	UNP F2RB80
A	-3	GLY	-	expression tag	UNP F2RB80
A	-2	SER	-	expression tag	UNP F2RB80
A	-1	HIS	-	expression tag	UNP F2RB80
A	189	ASN	ASP	conflict	UNP F2RB80

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Fe		
2	A	2	2	2	0	0

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	K		
3	A	1	1	1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	6	3	3	0	0
			Total	C	O		
4	A	1	6	3	3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	178	178	178	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CmlA protein

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α, β, γ	153.53Å 153.53Å 93.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.88 – 2.20 44.84 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.8 (44.88-2.20) 98.8 (44.84-2.20)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	5.81 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.180 , 0.213 0.189 , 0.213	Depositor DCC
R_{free} test set	2850 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å^2)	49.4	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 46.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	4402	wwPDB-VP
Average B, all atoms (Å^2)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.70% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: K, FE, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.70	0/4320	0.93	14/5878 (0.2%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	456	ARG	NE-CZ-NH2	-15.93	112.34	120.30
1	A	456	ARG	NE-CZ-NH1	9.58	125.09	120.30
1	A	318	ARG	NE-CZ-NH1	7.95	124.28	120.30
1	A	288	ARG	NE-CZ-NH2	-7.76	116.42	120.30
1	A	223	ARG	NE-CZ-NH2	7.60	124.10	120.30
1	A	311	MET	CB-CG-SD	-7.45	90.05	112.40
1	A	288	ARG	NE-CZ-NH1	7.21	123.91	120.30
1	A	318	ARG	NE-CZ-NH2	-6.80	116.90	120.30
1	A	325	ARG	NE-CZ-NH2	-6.75	116.93	120.30
1	A	456	ARG	CG-CD-NE	-6.62	97.89	111.80
1	A	223	ARG	NE-CZ-NH1	-6.31	117.14	120.30
1	A	325	ARG	NE-CZ-NH1	6.01	123.31	120.30
1	A	312	LEU	CA-CB-CG	-5.25	103.24	115.30
1	A	36	ARG	NE-CZ-NH1	5.17	122.89	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4209	0	4071	26	0
2	A	2	0	0	0	0
3	A	1	0	0	0	0
4	A	12	0	16	0	0
5	A	178	0	0	3	3
All	All	4402	0	4087	26	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:PRO:HG3	1:A:301:LEU:HD11	1.70	0.73
1:A:36:ARG:HH11	1:A:36:ARG:HG3	1.55	0.71
1:A:495:VAL:HG12	1:A:495:VAL:O	1.94	0.67
1:A:274:PRO:CG	1:A:301:LEU:HD11	2.26	0.65
1:A:8:ILE:O	1:A:318:ARG:NH2	2.33	0.62
1:A:278:TYR:O	1:A:288:ARG:NH2	2.32	0.61
1:A:501:HIS:HD2	1:A:503:ALA:H	1.50	0.60
1:A:15:ALA:HA	1:A:157[B]:TRP:CE3	2.36	0.59
1:A:311:MET:HA	5:A:797:HOH:O	2.04	0.56
1:A:325:ARG:HD3	1:A:327:LEU:HD21	1.90	0.52
1:A:285:ALA:O	1:A:288:ARG:HD3	2.10	0.52
1:A:41:VAL:HG11	1:A:79:TYR:CD1	2.47	0.50
1:A:379:GLY:O	1:A:456:ARG:HD2	2.14	0.47
1:A:406:ASN:H	1:A:465:GLN:HE21	1.63	0.47
1:A:378:HIS:CE1	1:A:386:LYS:HD2	2.50	0.46
1:A:256:VAL:HB	1:A:289:PHE:HB3	1.99	0.45
1:A:157[B]:TRP:HZ3	5:A:855:HOH:O	2.00	0.44
1:A:17:TRP:CZ2	1:A:37:PHE:HB3	2.52	0.43
1:A:180:GLU:OE2	1:A:223:ARG:NH1	2.52	0.42
1:A:291:PHE:CD1	1:A:318:ARG:HD2	2.53	0.42
1:A:433:GLY:H	1:A:459:ASN:ND2	2.18	0.42
1:A:495:VAL:O	1:A:495:VAL:CG1	2.64	0.42
1:A:172:PHE:HA	1:A:490:GLY:O	2.21	0.41
1:A:275:VAL:HB	1:A:312:LEU:HD13	2.03	0.41
1:A:15:ALA:HA	1:A:157[B]:TRP:CD2	2.57	0.40
1:A:482:ALA:HB1	5:A:740:HOH:O	2.21	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:703:HOH:O	5:A:802:HOH:O[8_554]	1.63	0.57
5:A:747:HOH:O	5:A:757:HOH:O[8_554]	1.66	0.54
5:A:712:HOH:O	5:A:802:HOH:O[8_554]	2.15	0.05

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	519/551 (94%)	503 (97%)	15 (3%)	1 (0%)	47 55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	531	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	442/462 (96%)	423 (96%)	19 (4%)	29 36

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-1	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	36	ARG
1	A	59	ARG
1	A	87	GLU
1	A	250	GLU
1	A	252	VAL
1	A	254	GLU
1	A	312	LEU
1	A	347	ARG
1	A	374	PHE
1	A	384	ARG
1	A	395	GLU
1	A	396	ARG
1	A	422	VAL
1	A	481	TYR
1	A	497	TYR
1	A	526	LEU
1	A	527	ARG
1	A	531	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	459	ASN
1	A	465	GLN
1	A	501	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GOL	A	605	-	5,5,5	0.46	0	5,5,5	0.79	0
4	GOL	A	604	-	5,5,5	0.36	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	605	-	-	2/4/4/4	-
4	GOL	A	604	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	605	GOL	O1-C1-C2-C3
4	A	604	GOL	O1-C1-C2-C3
4	A	605	GOL	O1-C1-C2-O2
4	A	604	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	521/551 (94%)	0.07	26 (4%) 28 27	37, 53, 82, 137	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	497	TYR	8.8
1	A	249	GLU	4.9
1	A	512	ARG	4.5
1	A	531	ARG	3.8
1	A	499	GLU	3.8
1	A	235	PRO	3.4
1	A	56	ARG	3.2
1	A	516	ASP	3.1
1	A	341	LEU	3.1
1	A	500	SER	3.0
1	A	395	GLU	2.7
1	A	46	ASP	2.6
1	A	498	ASP	2.5
1	A	80	ARG	2.4
1	A	49	GLN	2.4
1	A	525	HIS	2.4
1	A	58	ARG	2.3
1	A	337	VAL	2.3
1	A	476	ASP	2.2
1	A	339	PRO	2.2
1	A	517	LYS	2.2
1	A	-1	HIS	2.1
1	A	496	ASP	2.1
1	A	383	ILE	2.1
1	A	47	ASP	2.1
1	A	439	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	GOL	A	604	6/6	0.87	0.26	68,74,79,86	0
4	GOL	A	605	6/6	0.94	0.17	62,70,74,83	0
3	K	A	603	1/1	0.98	0.06	56,56,56,56	0
2	FE	A	602	1/1	0.99	0.13	52,52,52,52	0
2	FE	A	601	1/1	0.99	0.16	45,45,45,45	0

6.5 Other polymers (i)

There are no such residues in this entry.