



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 23, 2025 – 12:13 AM JST

PDB ID : 9KQ7 / pdb_00009kq7
Title : The structure of the YcfA from Erwinia amylovora bound with ATP
Authors : Zhang, L.; Dou, C.; Zheng, Y.H.; Zhu, X.F.; Cheng, W.
Deposited on : 2024-11-25
Resolution : 2.84 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1367 (2.86-2.82)
Clashscore	180529	1455 (2.86-2.82)
Ramachandran outliers	177936	1422 (2.86-2.82)
Sidechain outliers	177891	1423 (2.86-2.82)
RSRZ outliers	164620	1368 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	234	<div> <div>75%</div> <div>15%</div> <div>• 6%</div> </div>
1	B	234	<div> <div>76%</div> <div>16%</div> <div>• 8%</div> </div>
1	C	234	<div> <div>78%</div> <div>13%</div> <div>• 6%</div> </div>
1	D	234	<div> <div>75%</div> <div>16%</div> <div>•• 6%</div> </div>
1	E	234	<div> <div>75%</div> <div>16%</div> <div>• 6%</div> </div>
1	F	234	<div> <div>79%</div> <div>13%</div> <div>• 6%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	234	 79%11% • 7%
1	H	234	 74%17% • 6%
1	I	234	 79%15% • 6%
1	J	234	 71%21% • 6%
1	K	234	 77%14% • 7%
1	L	234	 69%22% • 7%
1	M	234	 75%16% • 7%
1	N	234	 76%15% • 7%

2 Entry composition


There are 3 unique types of molecules in this entry. The entry contains 24784 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Asparagine synthetase domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	216	Total	C	N	O	S	0	0	0
			1703	1084	293	316	10			
1	E	220	Total	C	N	O	S	0	0	0
			1731	1101	298	322	10			
1	G	218	Total	C	N	O	S	0	0	0
			1718	1093	296	319	10			
1	I	221	Total	C	N	O	S	0	0	0
			1738	1106	299	323	10			
1	J	219	Total	C	N	O	S	0	0	0
			1725	1098	297	320	10			
1	L	218	Total	C	N	O	S	0	0	0
			1718	1093	296	319	10			
1	M	218	Total	C	N	O	S	0	0	0
			1716	1092	295	319	10			
1	A	219	Total	C	N	O	S	0	0	0
			1725	1098	297	320	10			
1	C	220	Total	C	N	O	S	0	0	0
			1731	1101	298	322	10			
1	D	219	Total	C	N	O	S	0	0	0
			1725	1098	297	320	10			
1	F	221	Total	C	N	O	S	0	0	0
			1738	1106	299	323	10			
1	H	219	Total	C	N	O	S	0	0	0
			1725	1098	297	320	10			
1	K	218	Total	C	N	O	S	0	0	0
			1718	1093	296	319	10			
1	N	218	Total	C	N	O	S	0	0	0
			1716	1092	295	319	10			

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	E	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	G	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	I	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	J	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	L	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	M	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	F	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	H	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	K	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	N	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	19	Total O 19 19	0	0
3	E	11	Total O 11 11	0	0
3	G	9	Total O 9 9	0	0
3	I	15	Total O 15 15	0	0
3	J	8	Total O 8 8	0	0
3	L	10	Total O 10 10	0	0
3	M	8	Total O 8 8	0	0
3	A	20	Total O 20 20	0	0
3	C	29	Total O 29 29	0	0
3	D	30	Total O 30 30	0	0
3	F	20	Total O 20 20	0	0
3	H	9	Total O 9 9	0	0
3	K	12	Total O 12 12	0	0
3	N	23	Total O 23 23	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Asparagine synthetase domain-containing protein




- Molecule 1: Asparagine synthetase domain-containing protein



- Molecule 1: Asparagine synthetase domain-containing protein



- Molecule 1: Asparagine synthetase domain-containing protein






- Molecule 1: Asparagine synthetase domain-containing protein

Chain J: 71% 21% 6%




- Molecule 1: Asparagine synthetase domain-containing protein

Chain L: 69% 22% 7%



- Molecule 1: Asparagine synthetase domain-containing protein

Chain M: 75% 16% 7%



- Molecule 1: Asparagine synthetase domain-containing protein


Chain A: 75% 15% 6%



- Molecule 1: Asparagine synthetase domain-containing protein


Chain C: 78% 13% 6%





- Molecule 1: Asparagine synthetase domain-containing protein

Chain D: 75% 16% 6%




- Molecule 1: Asparagine synthetase domain-containing protein

Chain F: 79% 13% 6%



- Molecule 1: Asparagine synthetase domain-containing protein

Chain H: 74% 17% 6%



- Molecule 1: Asparagine synthetase domain-containing protein

Chain K: 77% 14% 7%



- Molecule 1: Asparagine synthetase domain-containing protein

Chain N: 76% 15% 7%





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	69.71Å 134.33Å 235.96Å 90.00° 98.43° 90.00°	Depositor
Resolution (Å)	33.34 – 2.84 33.34 – 2.84	Depositor EDS
% Data completeness (in resolution range)	99.8 (33.34-2.84) 99.7 (33.34-2.84)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 2.85Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, R_{free}	0.210 , 0.252 0.218 , 0.254	Depositor DCC
R_{free} test set	5003 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	65.2	Xtriage
Anisotropy	0.326	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 28.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.217 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	24784	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	0/1760	1.30	10/2382 (0.4%)
1	B	0.88	1/1736 (0.1%)	1.32	10/2347 (0.4%)
1	C	0.97	1/1765 (0.1%)	1.36	9/2388 (0.4%)
1	D	0.99	0/1759	1.38	13/2380 (0.5%)
1	E	0.86	1/1765 (0.1%)	1.35	11/2388 (0.5%)
1	F	0.83	0/1773	1.29	11/2400 (0.5%)
1	G	0.77	0/1752	1.28	11/2370 (0.5%)
1	H	0.87	0/1759	1.38	18/2380 (0.8%)
1	I	0.87	0/1773	1.34	9/2400 (0.4%)
1	J	0.84	0/1759	1.33	10/2380 (0.4%)
1	K	0.86	0/1752	1.38	13/2370 (0.5%)
1	L	0.76	0/1752	1.29	14/2370 (0.6%)
1	M	0.79	0/1749	1.25	7/2365 (0.3%)
1	N	0.93	0/1749	1.38	16/2365 (0.7%)
All	All	0.87	3/24603 (0.0%)	1.33	162/33285 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	4
1	D	0	2
1	E	0	2
1	F	0	3
1	G	0	2
1	H	0	3
1	I	0	3

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	J	0	2
1	K	0	2
1	L	0	3
1	M	0	2
1	N	0	2
All	All	0	34

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	127	HIS	CE1-NE2	-5.62	1.26	1.32
1	C	106	HIS	CD2-NE2	5.22	1.43	1.37
1	B	223	LEU	C-O	5.10	1.33	1.23

The worst 5 of 162 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	K	198	ASP	CA-CB-CG	11.21	123.81	112.60
1	D	198	ASP	CA-CB-CG	10.34	122.94	112.60
1	H	59	ASN	CA-CB-CG	-9.98	102.62	112.60
1	H	83	GLU	CB-CG-CD	8.78	127.52	112.60
1	N	145	GLU	CB-CG-CD	8.60	127.23	112.60

There are no chirality outliers.

5 of 34 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	114	ARG	Sidechain
1	B	171	ARG	Sidechain
1	E	136	ARG	Sidechain
1	E	47	ARG	Sidechain
1	G	47	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1725	0	1719	27	0
1	B	1703	0	1699	21	0
1	C	1731	0	1726	18	0
1	D	1725	0	1721	31	0
1	E	1731	0	1726	28	3
1	F	1738	0	1733	14	0
1	G	1718	0	1712	20	0
1	H	1725	0	1721	16	0
1	I	1738	0	1733	21	0
1	J	1725	0	1721	32	0
1	K	1718	0	1712	12	0
1	L	1718	0	1712	39	3
1	M	1716	0	1713	24	0
1	N	1716	0	1713	15	0
2	A	31	0	12	4	0
2	B	31	0	12	3	0
2	C	31	0	12	1	0
2	D	31	0	12	1	0
2	E	31	0	12	4	0
2	F	31	0	12	1	0
2	G	31	0	12	1	0
2	H	31	0	12	2	0
2	I	31	0	12	4	0
2	J	31	0	12	2	0
2	K	31	0	12	1	0
2	L	31	0	12	0	0
2	M	31	0	12	0	0
2	N	31	0	12	3	0
3	A	20	0	0	0	0
3	B	19	0	0	2	0
3	C	29	0	0	0	0
3	D	30	0	0	1	0
3	E	11	0	0	1	0
3	F	20	0	0	1	0
3	G	9	0	0	0	0
3	H	9	0	0	1	0
3	I	15	0	0	1	0
3	J	8	0	0	1	0
3	K	12	0	0	0	0
3	L	10	0	0	1	0
3	M	8	0	0	1	0
3	N	23	0	0	1	0
All	All	24784	0	24229	316	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 316 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:130:MET:HE1	1:E:159:TYR:OH	1.43	1.17
1:L:103:LEU:HD21	1:L:211:ALA:HB2	1.33	1.07
1:A:19:ASP:OD2	1:A:126:THR:HG21	1.54	1.07
1:L:103:LEU:CD2	1:L:211:ALA:HB2	1.86	1.05
1:E:130:MET:CE	1:E:159:TYR:OH	2.12	0.97

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:193:SER:OG	1:L:86:GLU:OE1[2_455]	2.09	0.11
1:E:86:GLU:OE2	1:L:95:ASN:N[2_455]	2.16	0.04
1:E:187:ILE:CG1	1:L:78:ARG:CG[2_455]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/234 (93%)	214 (99%)	3 (1%)	0	100	100
1	B	214/234 (92%)	213 (100%)	1 (0%)	0	100	100
1	C	218/234 (93%)	216 (99%)	1 (0%)	1 (0%)	25	44
1	D	217/234 (93%)	216 (100%)	1 (0%)	0	100	100
1	E	218/234 (93%)	211 (97%)	6 (3%)	1 (0%)	25	44
1	F	219/234 (94%)	218 (100%)	1 (0%)	0	100	100
1	G	216/234 (92%)	212 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	217/234 (93%)	216 (100%)	1 (0%)	0	100	100
1	I	219/234 (94%)	218 (100%)	1 (0%)	0	100	100
1	J	217/234 (93%)	216 (100%)	1 (0%)	0	100	100
1	K	216/234 (92%)	212 (98%)	3 (1%)	1 (0%)	25	44
1	L	216/234 (92%)	212 (98%)	3 (1%)	1 (0%)	25	44
1	M	216/234 (92%)	214 (99%)	2 (1%)	0	100	100
1	N	216/234 (92%)	214 (99%)	2 (1%)	0	100	100
All	All	3036/3276 (93%)	3002 (99%)	30 (1%)	4 (0%)	48	69

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	7	VAL
1	E	162	ALA
1	L	83	GLU
1	K	131	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/200 (94%)	179 (95%)	9 (5%)	21	43
1	B	185/200 (92%)	178 (96%)	7 (4%)	28	53
1	C	189/200 (94%)	183 (97%)	6 (3%)	34	59
1	D	188/200 (94%)	177 (94%)	11 (6%)	16	33
1	E	189/200 (94%)	183 (97%)	6 (3%)	34	59
1	F	190/200 (95%)	182 (96%)	8 (4%)	25	49
1	G	187/200 (94%)	181 (97%)	6 (3%)	34	59
1	H	188/200 (94%)	181 (96%)	7 (4%)	29	54
1	I	190/200 (95%)	183 (96%)	7 (4%)	29	54

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	J	188/200 (94%)	178 (95%)	10 (5%)	19	38
1	K	187/200 (94%)	176 (94%)	11 (6%)	16	33
1	L	187/200 (94%)	181 (97%)	6 (3%)	34	59
1	M	187/200 (94%)	176 (94%)	11 (6%)	16	33
1	N	187/200 (94%)	180 (96%)	7 (4%)	29	54
All	All	2630/2800 (94%)	2518 (96%)	112 (4%)	25	48

5 of 112 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	178	SER
1	N	199	VAL
1	D	105	ILE
1	N	181	SER
1	K	178	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	87	GLN
1	K	108	HIS
1	H	106	HIS
1	H	153	ASN
1	K	153	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ATP	K	301	-	26,33,33	0.73	0	31,52,52	0.84	1 (3%)
2	ATP	L	301	-	26,33,33	0.76	0	31,52,52	0.75	1 (3%)
2	ATP	M	301	-	26,33,33	0.75	0	31,52,52	0.79	1 (3%)
2	ATP	H	301	-	26,33,33	0.75	0	31,52,52	0.76	1 (3%)
2	ATP	B	301	-	26,33,33	0.76	0	31,52,52	0.76	1 (3%)
2	ATP	C	301	-	26,33,33	0.78	0	31,52,52	0.79	1 (3%)
2	ATP	J	301	-	26,33,33	0.75	0	31,52,52	0.86	2 (6%)
2	ATP	D	301	-	26,33,33	0.75	0	31,52,52	0.85	1 (3%)
2	ATP	N	301	-	26,33,33	0.78	0	31,52,52	0.76	1 (3%)
2	ATP	A	301	-	26,33,33	0.77	0	31,52,52	0.75	1 (3%)
2	ATP	I	301	-	26,33,33	0.75	0	31,52,52	0.76	1 (3%)
2	ATP	F	301	-	26,33,33	0.76	0	31,52,52	0.77	1 (3%)
2	ATP	G	301	-	26,33,33	0.73	0	31,52,52	0.78	1 (3%)
2	ATP	E	301	-	26,33,33	0.73	0	31,52,52	0.76	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	K	301	-	-	1/18/38/38	0/3/3/3
2	ATP	L	301	-	-	0/18/38/38	0/3/3/3
2	ATP	M	301	-	-	3/18/38/38	0/3/3/3
2	ATP	H	301	-	-	5/18/38/38	0/3/3/3
2	ATP	B	301	-	-	7/18/38/38	0/3/3/3
2	ATP	C	301	-	-	0/18/38/38	0/3/3/3
2	ATP	J	301	-	-	0/18/38/38	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	D	301	-	-	5/18/38/38	0/3/3/3
2	ATP	N	301	-	-	0/18/38/38	0/3/3/3
2	ATP	A	301	-	-	1/18/38/38	0/3/3/3
2	ATP	I	301	-	-	5/18/38/38	0/3/3/3
2	ATP	F	301	-	-	0/18/38/38	0/3/3/3
2	ATP	G	301	-	-	4/18/38/38	0/3/3/3
2	ATP	E	301	-	-	3/18/38/38	0/3/3/3

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	301	ATP	C5-C6-N6	2.41	124.01	120.35
2	C	301	ATP	C5-C6-N6	2.36	123.94	120.35
2	K	301	ATP	C5-C6-N6	2.35	123.92	120.35
2	D	301	ATP	C5-C6-N6	2.35	123.92	120.35
2	F	301	ATP	C5-C6-N6	2.34	123.90	120.35

There are no chirality outliers.

5 of 34 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	301	ATP	O4'-C4'-C5'-O5'
2	H	301	ATP	PB-O3B-PG-O2G
2	H	301	ATP	PB-O3B-PG-O3G
2	G	301	ATP	O4'-C4'-C5'-O5'
2	I	301	ATP	O4'-C4'-C5'-O5'

There are no ring outliers.

12 monomers are involved in 27 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	K	301	ATP	1	0
2	H	301	ATP	2	0
2	B	301	ATP	3	0
2	C	301	ATP	1	0
2	J	301	ATP	2	0
2	D	301	ATP	1	0
2	N	301	ATP	3	0
2	A	301	ATP	4	0

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	I	301	ATP	4	0
2	F	301	ATP	1	0
2	G	301	ATP	1	0
2	E	301	ATP	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Ligand ATP L 301















Ligand ATP J 301













Ligand ATP I 301



Ligand ATP F 301







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	219/234 (93%)	-1.70	0 100 100	42, 63, 96, 122	0
1	B	216/234 (92%)	-1.35	0 100 100	43, 61, 86, 105	0
1	C	220/234 (94%)	-1.64	0 100 100	32, 49, 72, 83	0
1	D	219/234 (93%)	-1.64	0 100 100	28, 50, 95, 134	0
1	E	220/234 (94%)	-1.44	0 100 100	54, 81, 130, 177	0
1	F	221/234 (94%)	-1.71	0 100 100	40, 61, 88, 114	0
1	G	218/234 (93%)	-1.58	0 100 100	60, 84, 107, 124	0
1	H	219/234 (93%)	-1.74	0 100 100	42, 58, 87, 119	0
1	I	221/234 (94%)	-1.48	0 100 100	38, 61, 87, 113	0
1	J	219/234 (93%)	-1.41	0 100 100	42, 67, 116, 157	0
1	K	218/234 (93%)	-1.62	0 100 100	43, 63, 104, 134	0
1	L	218/234 (93%)	-1.55	0 100 100	60, 87, 119, 150	0
1	M	218/234 (93%)	-1.63	0 100 100	56, 79, 105, 145	0
1	N	218/234 (93%)	-1.63	0 100 100	33, 49, 72, 96	0
All	All	3064/3276 (93%)	-1.58	0 100 100	28, 66, 106, 177	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ATP	B	301	31/31	0.99	0.04	49,65,91,109	0
2	ATP	E	301	31/31	0.99	0.03	63,79,120,135	0
2	ATP	G	301	31/31	0.99	0.03	72,96,110,123	0
2	ATP	I	301	31/31	0.99	0.04	59,79,104,109	0
2	ATP	J	301	31/31	0.99	0.04	52,75,102,137	0
2	ATP	L	301	31/31	0.99	0.03	75,101,124,141	0
2	ATP	M	301	31/31	0.99	0.04	75,95,115,123	0
2	ATP	A	301	31/31	0.99	0.03	49,74,93,97	0
2	ATP	C	301	31/31	0.99	0.04	36,58,73,80	0
2	ATP	D	301	31/31	0.99	0.03	48,58,91,94	0
2	ATP	F	301	31/31	0.99	0.03	52,71,90,109	0
2	ATP	H	301	31/31	0.99	0.03	48,67,101,106	0
2	ATP	K	301	31/31	0.99	0.03	55,69,119,132	0
2	ATP	N	301	31/31	0.99	0.04	44,59,73,80	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ATP B 301:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP E 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP G 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP I 301:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP J 301:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP L 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ATP M 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ATP C 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP F 301:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP H 301:


$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ATP K 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.