wwPDB NMR Structure Validation Summary Report (i)

Apr 21, 2024 – 04:43 PM EDT

PDB ID	:	2KXJ
Title	:	Solution structure of UBX domain of human UBXD2 protein
Authors	:	Wu, Q.; Huang, H.; Zhang, J.; Hu, Q.; Wu, J.; Shi, Y.
Deposited on	:	2010-05-06

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	90	57% 28% • 11%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 9 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:318-A:397 (80)	0.55	9

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 3 single-model clusters were found.

Cluster number	Models
1	2, 3, 5, 9, 10, 15, 19
2	1, 7, 8, 16, 20
3	4, 14, 17
4	6, 11
Single-model clusters	12; 13; 18

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1409 atoms, of which 696 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called UBX domain-containing protein 4.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	90	1409	451	696	128	131	3	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	308	MET	-	expression tag	UNP Q92575
A	309	GLY	-	expression tag	UNP Q92575
A	310	HIS	-	expression tag	UNP Q92575
A	311	HIS	-	expression tag	UNP Q92575
A	312	HIS	-	expression tag	UNP Q92575
A	313	HIS	-	expression tag	UNP Q92575
A	314	HIS	-	expression tag	UNP Q92575
A	315	HIS	-	expression tag	UNP Q92575
A	316	MET	-	expression tag	UNP Q92575

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: UBX domain-containing protein 4

Chain A: 57% 28% • 11%

M308 G309 H310 H311 H312 H313 H314 H315 M316 S317 R321 I322 Q323 F324 R325 L326 P327 D328 S331 F332 T333 N334 Q335 F336 P337 S338 A363 R368 R369 E370 K373 K377 K378 K379 L380 L381 D382 L385 V392 L393 L394 P395 A396 G397

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 9. Colouring as in section 4.1 above.

- Molecule 1: UBX domain-containing protein 4

Chain A: 54% 28% 7% 11%

M308 G309 H310 H311 H312 H313 H314 H315 M316 S317 T318 V319 I322 Q323 F324 R325 L326 P327 D328 G329 S330 S331 F332 F336 P337 S338 Q351 Y357 G358 N359 A363 R368 R369 E370 F371 T372 K373 K377 L380 L381 D382 L383 E384 L385 S388 V392 L393 L394 P395 A396

G397

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	1.1
CNS	refinement	1.1

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	627	620	620	18±4
All	All	12540	12400	12400	352

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

5 of 96 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:322:ILE:HD12	1:A:385:LEU:HD12	1.02	1.32	16	6
1:A:322:ILE:HD11	1:A:380:LEU:HD23	0.83	1.50	8	14
1:A:380:LEU:HD23	1:A:385:LEU:HD13	0.81	1.51	15	5
1:A:380:LEU:O	1:A:385:LEU:HD23	0.77	1.78	3	6
1:A:381:LEU:HD13	1:A:382:ASP:N	0.76	1.96	20	4

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	79/90 (88%)	70±2 (89±3%)	8±2 (11±3%)	1±1 (1±1%)	24 71
All	All	1580/1800 (88%)	1399 (89%)	169 (11%)	12 (1%)	24 71

5 of 7 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	355	ASN	3
1	A	370	GLU	2
1	A	384	GLU	2
1	A	358	GLY	2
1	A	318	THR	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	67/76 (88%)	48±2 (72±3%)	18±2 (28±3%)	2 20
All	All	1340/1520 (88%)	970 (72%)	370 (28%)	2 20

5 of 46 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	322	ILE	20
1	A	326	LEU	20
1	A	331	SER	20
1	A	377	LYS	20
1	A	324	PHE	18

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided