wwPDB X-ray Structure Validation Summary Report (i)

Sep 23, 2023 – 04:49 PM EDT

PDB ID	:	5L0Y
Title	:	Crystal Structure of a Sec72-ssa1 c-terminal peptide fusion protein
Authors	:	Tripathi, A.; Rapoport, T.A.
Deposited on	:	2016-07-28
Resolution	:	2.87 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2691 (2.90-2.86)
Clashscore	141614	2947 (2.90-2.86)
Ramachandran outliers	138981	2868 (2.90-2.86)
Sidechain outliers	138945	2871 (2.90-2.86)
RSRZ outliers	127900	2629 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	2%, 63%, 29%, 6%, •
1	B	159	3%, 60%, 30%, •, 6%
1	C	159	8%, 58%, 32%, •, 6%
1	D	159	3%, 57%, 31%, 6%, 6%
1	E	159	2%, 57%, 33%, 5%, 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	159	%, 63%, 21%, 12%
1	G	159	8%, 52%, 33%, 9%
1	H	159	10%, 50%, 27%, 6%, 16%
2	I	7	14%, 57%, 14%, 14%
2	J	7	43%, 57%
2	K	7	14%, 71%, 14%, 14%
2	L	7	29%, 29%, 29%, 14%
2	M	7	29%, 29%, 14%, 29%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 9720 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sec72-ssa1 c-terminal peptide fusion protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	155	1248	772	233	237	6	0	0	0
1	B	149	1201	743	227	225	6	0	0	0
1	C	150	1209	747	229	227	6	0	0	0
1	D	150	1212	749	224	233	6	0	0	0
1	E	151	1220	756	228	230	6	0	0	0
1	F	140	1132	702	211	213	6	0	0	0
1	G	144	1162	722	222	212	6	0	0	0
1	H	133	1089	677	205	201	6	0	0	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	215	HIS	-	linker	UNP G0SH41
A	216	ASP	-	linker	UNP G0SH41
A	217	ASN	-	linker	UNP G0SH41
A	218	ASP	-	linker	UNP G0SH41
B	215	HIS	-	linker	UNP G0SH41
B	216	ASP	-	linker	UNP G0SH41
B	217	ASN	-	linker	UNP G0SH41
B	218	ASP	-	linker	UNP G0SH41
C	215	HIS	-	linker	UNP G0SH41
C	216	ASP	-	linker	UNP G0SH41
C	217	ASN	-	linker	UNP G0SH41
C	218	ASP	-	linker	UNP G0SH41
D	215	HIS	-	linker	UNP G0SH41

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	216	ASP	-	linker	UNP G0SH41
D	217	ASN	-	linker	UNP G0SH41
D	218	ASP	-	linker	UNP G0SH41
E	215	HIS	-	linker	UNP G0SH41
E	216	ASP	-	linker	UNP G0SH41
E	217	ASN	-	linker	UNP G0SH41
E	218	ASP	-	linker	UNP G0SH41
F	215	HIS	-	linker	UNP G0SH41
F	216	ASP	-	linker	UNP G0SH41
F	217	ASN	-	linker	UNP G0SH41
F	218	ASP	-	linker	UNP G0SH41
G	215	HIS	-	linker	UNP G0SH41
G	216	ASP	-	linker	UNP G0SH41
G	217	ASN	-	linker	UNP G0SH41
G	218	ASP	-	linker	UNP G0SH41
H	215	HIS	-	linker	UNP G0SH41
H	216	ASP	-	linker	UNP G0SH41
H	217	ASN	-	linker	UNP G0SH41
H	218	ASP	-	linker	UNP G0SH41

- Molecule 2 is a protein called PRO-THR-VAL-GLU-GLU-VAL-ASP.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	I	6	48	28	6	14	0	0	0
2	J	7	55	33	7	15	0	0	0
2	K	7	55	33	7	15	0	0	0
2	L	6	48	28	6	14	0	0	0
2	M	5	41	24	5	12	0	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Sec72-ssa1 c-terminal peptide fusion protein

Chain A:



• Molecule 1: Sec72-ssa1 c-terminal peptide fusion protein

Chain B:



• Molecule 1: Sec72-ssa1 c-terminal peptide fusion protein

Chain C:



• Molecule 1: Sec72-ssa1 c-terminal peptide fusion protein

Chain D:



• Molecule 1: Sec72-ssa1 c-terminal peptide fusion protein

Chain E:



• Molecule 1: Sec72-ssa1 c-terminal peptide fusion protein

Chain F:



• Molecule 1: Sec72-ssa1 c-terminal peptide fusion protein

Chain G:



• Molecule 1: Sec72-ssa1 c-terminal peptide fusion protein

Chain H:



- Molecule 2: PRO-THR-VAL-GLU-GLU-VAL-ASP

Chain I:



- Molecule 2: PRO-THR-VAL-GLU-GLU-VAL-ASP

Chain J:



- Molecule 2: PRO-THR-VAL-GLU-GLU-VAL-ASP

Chain K:



- Molecule 2: PRO-THR-VAL-GLU-GLU-VAL-ASP

Chain L:



- Molecule 2: PRO-THR-VAL-GLU-GLU-VAL-ASP

Chain M:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α, β, γ	76.50Å 118.53Å 164.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	96.08 – 2.87 164.05 – 2.87	Depositor EDS
% Data completeness (in resolution range)	99.3 (96.08-2.87) 99.3 (164.05-2.87)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.12 (at 2.86Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.236 , 0.283 0.237 , 0.281	Depositor DCC
R_{free} test set	1747 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å^2)	69.5	Xtriage
Anisotropy	0.406	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 57.4	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	9720	wwPDB-VP
Average B, all atoms (Å^2)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.10% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.49	0/1268	0.53	0/1700
1	B	0.33	0/1221	0.48	0/1638
1	C	0.37	0/1229	0.54	1/1649 (0.1%)
1	D	0.40	0/1230	0.53	0/1648
1	E	0.36	0/1239	0.51	0/1658
1	F	0.43	1/1149 (0.1%)	0.53	0/1539
1	G	0.39	0/1181	0.59	1/1583 (0.1%)
1	H	0.40	0/1105	0.59	2/1476 (0.1%)
2	I	0.69	0/47	0.53	0/62
2	J	0.45	0/55	0.69	0/73
2	K	0.34	0/55	0.57	0/73
2	L	0.29	0/47	0.50	0/62
2	M	0.20	0/40	0.42	0/52
All	All	0.40	1/9866 (0.0%)	0.54	4/13213 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	190	GLU	CG-CD	5.33	1.59	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	H	205	LEU	CA-CB-CG	7.84	133.34	115.30
1	G	117	LEU	CA-CB-CG	6.39	129.99	115.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	116	GLY	N-CA-C	-5.34	99.75	113.10
1	H	202	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	114	PRO	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1248	0	1230	43	0
1	B	1201	0	1186	39	1
1	C	1209	0	1192	46	0
1	D	1212	0	1183	43	0
1	E	1220	0	1206	34	0
1	F	1132	0	1118	26	1
1	G	1162	0	1160	39	0
1	H	1089	0	1082	43	0
2	I	48	0	40	7	0
2	J	55	0	48	4	0
2	K	55	0	48	3	0
2	L	48	0	40	4	0
2	M	41	0	33	3	0
All	All	9720	0	9566	301	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 301 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:155:GLN:HA	1:A:188:GLU:HB2	1.34	1.09

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:109:ARG:HD2	1:H:119:ARG:HE	1.30	0.96
1:A:156:GLY:N	1:A:188:GLU:OE1	1.99	0.94
1:A:158:ALA:N	1:A:188:GLU:OE2	2.03	0.92
1:A:157:ASN:N	1:A:188:GLU:OE1	2.04	0.91

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:213:ASP:OD1	1:F:190:GLU:OE1[2_655]	1.42	0.78

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/159 (95%)	147 (97%)	4 (3%)	0	100	100
1	B	147/159 (92%)	142 (97%)	5 (3%)	0	100	100
1	C	148/159 (93%)	141 (95%)	7 (5%)	0	100	100
1	D	144/159 (91%)	141 (98%)	3 (2%)	0	100	100
1	E	145/159 (91%)	137 (94%)	8 (6%)	0	100	100
1	F	136/159 (86%)	133 (98%)	3 (2%)	0	100	100
1	G	140/159 (88%)	131 (94%)	9 (6%)	0	100	100
1	H	127/159 (80%)	124 (98%)	3 (2%)	0	100	100
2	I	4/7 (57%)	4 (100%)	0	0	100	100
2	J	5/7 (71%)	5 (100%)	0	0	100	100
2	K	5/7 (71%)	5 (100%)	0	0	100	100
2	L	4/7 (57%)	4 (100%)	0	0	100	100
2	M	3/7 (43%)	3 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1159/1307 (89%)	1117 (96%)	42 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/127 (98%)	103 (83%)	21 (17%)	2	5
1	B	118/127 (93%)	102 (86%)	16 (14%)	3	10
1	C	119/127 (94%)	107 (90%)	12 (10%)	7	21
1	D	120/127 (94%)	100 (83%)	20 (17%)	2	6
1	E	121/127 (95%)	99 (82%)	22 (18%)	1	4
1	F	111/127 (87%)	97 (87%)	14 (13%)	4	12
1	G	114/127 (90%)	95 (83%)	19 (17%)	2	6
1	H	107/127 (84%)	88 (82%)	19 (18%)	2	4
2	I	6/7 (86%)	4 (67%)	2 (33%)	0	0
2	J	7/7 (100%)	7 (100%)	0	100	100
2	K	7/7 (100%)	6 (86%)	1 (14%)	3	9
2	L	6/7 (86%)	3 (50%)	3 (50%)	0	0
2	M	5/7 (71%)	4 (80%)	1 (20%)	1	3
All	All	965/1051 (92%)	815 (84%)	150 (16%)	2	7

5 of 150 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	135	GLN
2	I	221	THR
1	G	173	ARG
1	H	166	LYS
1	C	217	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	128	ASN
1	B	215	HIS
1	D	90	GLN
1	E	104	GLN
1	G	138	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	155/159 (97%)	0.23	3 (1%)	66	65	24, 58, 105, 152	0
1	B	149/159 (93%)	0.21	4 (2%)	54	51	34, 67, 110, 158	0
1	C	150/159 (94%)	0.55	12 (8%)	12	9	38, 77, 148, 256	0
1	D	150/159 (94%)	0.25	5 (3%)	46	41	35, 73, 132, 294	0
1	E	151/159 (94%)	0.34	3 (1%)	65	63	46, 84, 123, 142	0
1	F	140/159 (88%)	0.14	1 (0%)	87	87	32, 54, 102, 133	0
1	G	144/159 (90%)	0.59	13 (9%)	9	6	46, 87, 131, 170	0
1	H	133/159 (83%)	0.65	16 (12%)	4	3	36, 86, 168, 233	0
2	I	6/7 (85%)	0.73	0	100	100	73, 86, 99, 99	0
2	J	7/7 (100%)	-0.08	0	100	100	40, 59, 71, 72	0
2	K	7/7 (100%)	0.48	1 (14%)	2	2	95, 110, 114, 116	0
2	L	6/7 (85%)	0.38	0	100	100	82, 101, 110, 116	0
2	M	5/7 (71%)	0.33	0	100	100	72, 82, 94, 98	0
All	All	1203/1307 (92%)	0.37	58 (4%)	30	26	24, 73, 133, 294	0

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	68	VAL	11.1
1	C	190	GLU	6.5
1	H	187	PHE	5.7
1	G	118	VAL	5.1
1	G	201	ILE	4.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.