



wwPDB NMR Structure Validation Summary Report ⓘ

Oct 17, 2021 – 09:28 AM EDT

PDB ID : 1L3N
Title : The Solution Structure of Reduced Dimeric Copper Zinc SOD: the Structural Effects of Dimerization
Authors : Banci, L.; Bertini, I.; Cramaro, F.; Del Conte, R.; Viezzoli, M.S.
Deposited on : 2002-02-28

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.23.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	153	 88% 12% .
1	B	153	 80% 16% .

2 Ensemble composition and analysis

This entry contains 30 models. Model 15 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:153, B:1-B:23, B:27-B:107, B:111-B:153 (300)	0.56	15

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 5 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 5, 6, 7, 9, 12, 14, 15, 16, 18, 21, 22, 24, 26, 29
2	4, 8, 30
3	10, 20, 23
4	11, 13
Single-model clusters	17; 19; 25; 27; 28

3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4378 atoms, of which 2156 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called superoxide dismutase [Cu-Zn].

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	153	2187	679	1078	203	225	2	0
1	B	153	2187	679	1078	203	225	2	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	6	ALA	CYS	engineered mutation	UNP P00441
A	111	SER	CYS	engineered mutation	UNP P00441
B	6	ALA	CYS	engineered mutation	UNP P00441
B	111	SER	CYS	engineered mutation	UNP P00441

- Molecule 2 is COPPER (I) ION (three-letter code: CU1) (formula: Cu).

Mol	Chain	Residues	Atoms	
			Total	Cu
2	A	1	1	1
2	B	1	1	1

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
			Total	Zn
3	A	1	1	1
3	B	1	1	1

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: superoxide dismutase [Cu-Zn]




- Molecule 1: superoxide dismutase [Cu-Zn]




4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 15. Colouring as in section 4.1 above.

- Molecule 1: superoxide dismutase [Cu-Zn]



- Molecule 1: superoxide dismutase [Cu-Zn]



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing and restrained energy minimization in vacuum*.

Of the 1400 calculated structures, 30 were deposited, based on the following criterion: *structure with the lowest energy target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CALIBA	structure solution	
GLOMSA	structure solution	
DYANA	structure solution	
Amber	refinement	

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CU1, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.50±0.01	0±0/1127 (0.0± 0.0%)	1.02±0.02	0±0/1519 (0.0± 0.0%)
1	B	0.51±0.01	0±0/1081 (0.0± 0.0%)	1.03±0.02	1±1/1457 (0.0± 0.0%)
All	All	0.51	0/66240 (0.0%)	1.03	23/89280 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.7±1.0
1	B	0.0±0.0	1.7±1.1
All	All	0	102

There are no bond-length outliers.

5 of 10 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	143	ARG	NE-CZ-NH1	7.99	124.30	120.30	27	6
1	A	115	ARG	NE-CZ-NH2	-7.55	116.53	120.30	19	1
1	B	69	ARG	NE-CZ-NH1	7.12	123.86	120.30	17	2
1	A	143	ARG	NE-CZ-NH1	7.00	123.80	120.30	11	3
1	B	69	ARG	NE-CZ-NH2	-6.58	117.01	120.30	29	4

There are no chirality outliers.

5 of 21 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	B	64	PHE	Sidechain	11
1	A	64	PHE	Sidechain	11
1	A	115	ARG	Sidechain	10
1	A	69	ARG	Sidechain	9
1	B	115	ARG	Sidechain	9

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1109	1078	1077	8±2
1	B	1064	1047	1046	7±2
All	All	65310	63750	63690	396

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

5 of 124 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:17:ILE:CG2	1:B:54:THR:HG22	0.85	2.01	24	5
1:B:16:GLY:HA2	1:B:35:ILE:HG22	0.77	1.55	9	18
1:A:16:GLY:HA2	1:A:35:ILE:HG22	0.77	1.56	27	20
1:A:54:THR:HG22	1:B:17:ILE:HG22	0.77	1.55	10	7
1:A:54:THR:HG22	1:B:17:ILE:CG2	0.76	2.10	5	8

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	151/153 (99%)	117±3 (78±2%)	26±3 (17±2%)	8±2 (5±2%)	4 24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	145/153 (95%)	113±5 (78±3%)	25±4 (17±3%)	7±2 (5±1%)	4	24
All	All	8880/9180 (97%)	6910 (78%)	1514 (17%)	456 (5%)	4	24

5 of 71 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	81	VAL	30
1	A	10	GLY	28
1	B	10	GLY	24
1	A	81	VAL	23
1	A	60	ALA	18

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/117 (100%)	102±3 (87±2%)	15±3 (13±2%)	7	48
1	B	112/117 (96%)	97±3 (87±3%)	15±3 (13±3%)	7	48
All	All	6870/7020 (98%)	5977 (87%)	893 (13%)	7	48

5 of 147 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	46	HIS	30
1	B	46	HIS	28
1	B	53	ASN	26
1	A	54	THR	24
1	A	15	GLN	21

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided