



wwPDB EM Validation Summary Report ⓘ

Oct 19, 2025 – 12:49 AM JST

PDB ID : 9LEM / pdb_00009lem
EMDB ID : EMD-63029
Title : Composite map of Sag-18RS21-tRNASer Golld RNA
Authors : Zhang, K.; Li, S.
Deposited on : 2025-01-07
Resolution : 2.98 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : **FAILED**
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
EM percentile statistics : **NOT EXECUTED**
MapQ : **FAILED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY











The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
RNA backbone	6643	2191

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	476	 57% 29% 5% 9%
1	B	476	 57% 29% 5% 9%
1	C	476	 57% 28% 6% 9%
1	D	476	 57% 29% 5% 9%
1	E	476	 57% 29% 6% 9%
1	F	476	 57% 29% 5% 9%
1	G	476	 57% 29% 5% 9%
1	H	476	 57% 29% 5% 9%
1	I	476	 57% 29% 6% 9%
1	J	476	 57% 29% 5% 9%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 92800 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a RNA chain called Sag-18RS21-tRNA^{Ser} Golld RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	J	434	Total	C	N	O	P	0	0
			9280	4149	1673	3024	434		
1	A	434	Total	C	N	O	P	0	0
			9280	4149	1673	3024	434		
1	B	434	Total	C	N	O	P	0	0
			9280	4149	1673	3024	434		
1	C	434	Total	C	N	O	P	0	0
			9280	4149	1673	3024	434		
1	D	434	Total	C	N	O	P	0	0
			9280	4149	1673	3024	434		
1	E	434	Total	C	N	O	P	0	0
			9280	4149	1673	3024	434		
1	F	434	Total	C	N	O	P	0	0
			9280	4149	1673	3024	434		
1	G	434	Total	C	N	O	P	0	0
			9280	4149	1673	3024	434		
1	H	434	Total	C	N	O	P	0	0
			9280	4149	1673	3024	434		
1	I	434	Total	C	N	O	P	0	0
			9280	4149	1673	3024	434		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Sag-18RS21-tRNA^{Ser} Goldd RNA




• Molecule 1: Sag-18RS21-tRNA^{Ser} Goldd RNA




• Molecule 1: Sag-18RS21-tRNA^{Ser} Goldd RNA




• Molecule 1: Sag-18RS21-tRNA^{Ser} Goldd RNA




• Molecule 1: Sag-18RS21-tRNA^{Ser} Goldd RNA






• Molecule 1: Sag-18RS21-tRNA^{Ser} Goldd RNA

Chain E: 57% 29% 6% 9%




• Molecule 1: Sag-18RS21-tRNA^{Ser} Goldd RNA

Chain F: 57% 29% 5% 9%




• Molecule 1: Sag-18RS21-tRNA^{Ser} Goldd RNA

Chain G: 




• Molecule 1: Sag-18RS21-tRNA^{Ser} Gold RNA


Chain H: 



• Molecule 1: Sag-18RS21-tRNA^{Ser} Gold RNA

Chain I: 





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, D5	Depositor
Number of particles used	1296920	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	46	Depositor
Minimum defocus (nm)	700	Depositor
Maximum defocus (nm)	1500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/10389	0.49	1/16191 (0.0%)
1	B	0.43	0/10389	0.49	1/16191 (0.0%)
1	C	0.43	0/10389	0.49	2/16191 (0.0%)
1	D	0.43	0/10389	0.49	2/16191 (0.0%)
1	E	0.43	0/10389	0.49	1/16191 (0.0%)
1	F	0.43	0/10389	0.49	1/16191 (0.0%)
1	G	0.43	0/10389	0.49	2/16191 (0.0%)
1	H	0.43	0/10389	0.49	1/16191 (0.0%)
1	I	0.43	0/10389	0.49	2/16191 (0.0%)
1	J	0.43	0/10389	0.49	2/16191 (0.0%)
All	All	0.43	0/103890	0.49	15/161910 (0.0%)

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	1	A	C2'-C3'-O3'	5.47	117.70	109.50
1	A	1	A	C2'-C3'-O3'	5.47	117.70	109.50
1	G	1	A	C2'-C3'-O3'	5.47	117.70	109.50
1	B	1	A	C2'-C3'-O3'	5.46	117.69	109.50
1	C	1	A	C2'-C3'-O3'	5.46	117.68	109.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9280	0	4661	58	0
1	B	9280	0	4661	66	0
1	C	9280	0	4661	71	0
1	D	9280	0	4661	55	0
1	E	9280	0	4661	67	0
1	F	9280	0	4661	62	0
1	G	9280	0	4661	52	0
1	H	9280	0	4661	61	0
1	I	9280	0	4661	78	0
1	J	9280	0	4661	57	0
All	All	92800	0	46610	538	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 538 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:171:C:O2	1:I:105:G:N2	2.02	0.93
1:C:401:G:N2	1:C:449:C:H42	1.67	0.93
1:A:401:G:N2	1:A:449:C:H42	1.67	0.93
1:I:401:G:N2	1:I:449:C:H42	1.67	0.93
1:F:401:G:N2	1:F:449:C:H42	1.67	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	433/476 (90%)	122 (28%)	5 (1%)
1	B	433/476 (90%)	122 (28%)	5 (1%)
1	C	433/476 (90%)	122 (28%)	5 (1%)
1	D	433/476 (90%)	122 (28%)	5 (1%)
1	E	433/476 (90%)	122 (28%)	5 (1%)
1	F	433/476 (90%)	122 (28%)	5 (1%)
1	G	433/476 (90%)	122 (28%)	5 (1%)
1	H	433/476 (90%)	122 (28%)	5 (1%)
1	I	433/476 (90%)	122 (28%)	5 (1%)
1	J	433/476 (90%)	122 (28%)	5 (1%)
All	All	4330/4760 (90%)	1220 (28%)	50 (1%)

5 of 1220 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	J	2	G
1	J	3	U
1	J	4	A
1	J	5	G
1	J	8	G

5 of 50 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	E	359	A
1	G	1	A
1	I	360	A
1	E	360	A
1	F	43	C

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.