Full wwPDB NMR Structure Validation Report (i)

Jun 16, 2024 – 07:34 PM EDT

PDB ID	:	5LGM
BMRB ID	:	34024
Title	:	Gp5.7 mutant L42A
Authors	:	Liu, B.; Matthews, S.
Deposited on	:	2016-07-07

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 88%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	69	55% 20% • 22%

2 Ensemble composition and analysis (i)

This entry contains 10 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:16-A:28, A:34-A:56, A:67-A:84 (54)	0.44	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 5, 6, 9
2	3, 7
Single-model clusters	8; 10

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1032 atoms, of which 518 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Fusion protein 5.5/5.7.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	69	1032	315	518	97	97	5	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	ALA	LEU	conflict	UNP P03787

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Fusion protein 5.5/5.7

Chain A: 55% 20% • 22%

M16 Y19 L20 L23 Q24 A25 I26 K27 S28 C29 P30 K31 T32 F33 Q34 S35 V38 A48 R51 I54 A57 T58 T59 S60 G61 R62 N63 G64 G65 A66 F76 L77 K78 R79 V84

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

• Molecule 1: Fusion protein 5.5/5.7

Chain A: 52% 20% 6% 22%

M16 Y19 L20 K21 V22 L23 Q24 A25 I26 K27 S28 C29 P30 K31 T32 F33 Q34 S35 N36 Y37 A48 R51 I54 A57 T58 T59 S60 G61 R62 N63 G64 G65 A66 W67 E68 F76 L77 K78 R79 V84

4.2.2 Score per residue for model 2 (medoid)

• Molecule 1: Fusion protein 5.5/5.7

Chain A: 55% 17% 6% 22%

M16 Y19 L20 L23 Q24 A25 C29 P30 K31 T32 F33 Q34 S35 V38 R39 A48 R51 I54 S55 C56 A57 T58 T59 S60 G61 R62 N63 G64 G65 A66 G73 L77 V84

4.2.3 Score per residue for model 3

- Molecule 1: Fusion protein 5.5/5.7

Chain A: 51% 22% • • 22%



4.2.4 Score per residue for model 4

- Molecule 1: Fusion protein 5.5/5.7

Chain A: 57% 16% 6% 22%



4.2.5 Score per residue for model 5

- Molecule 1: Fusion protein 5.5/5.7

Chain A: 55% 17% 6% 22%

4.2.6 Score per residue for model 6

- Molecule 1: Fusion protein 5.5/5.7

Chain A: 54% 22% •• 22%

4.2.7 Score per residue for model 7

- Molecule 1: Fusion protein 5.5/5.7

Chain A: 51% 22% 6% 22%



4.2.8 Score per residue for model 8

- Molecule 1: Fusion protein 5.5/5.7

Chain A:



4.2.9 Score per residue for model 9

- Molecule 1: Fusion protein 5.5/5.7

Chain A:



4.2.10 Score per residue for model 10

- Molecule 1: Fusion protein 5.5/5.7

Chain A:



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *distance geometry, simulated annealing.*

Of the 10 calculated structures, 10 were deposited, based on the following criterion: *all calculated structures submitted.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	759
Number of shifts mapped to atoms	759
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	88%

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.3±0.5
All	All	0	3

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	19	TYR	Sidechain	3

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	413	420	416	8±4
All	All	4130	4200	4160	83

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:19:TYR:OH	1:A:48:ALA:HA	0.73	1.83	6	10
1:A:76:PHE:O	1:A:79:ARG:HG2	0.65	1.91	8	8

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:16:MET:O	1:A:20:LEU:HG	0.64	1.92	9	10
1:A:77:LEU:HG	1:A:78:LYS:N	0.62	2.07	10	1
1:A:24:GLN:NE2	1:A:74:THR:HA	0.62	2.09	9	1
1:A:24:GLN:O	1:A:27:LYS:HG3	0.62	1.94	9	3
1:A:24:GLN:NE2	1:A:78:LYS:HB2	0.60	2.11	9	1
1:A:73:GLY:O	1:A:77:LEU:HB3	0.58	1.98	9	2
1:A:24:GLN:OE1	1:A:78:LYS:HB2	0.57	2.00	8	2
1:A:24:GLN:OE1	1:A:78:LYS:HD3	0.55	2.01	3	1
1:A:24:GLN:OE1	1:A:74:THR:HA	0.54	2.02	8	1
1:A:26:ILE:CG2	1:A:34:GLN:HB3	0.53	2.34	10	1
1:A:23:LEU:O	1:A:26:ILE:HG22	0.52	2.04	8	3
1:A:38:VAL:HG21	1:A:67:TRP:H	0.51	1.65	9	2
1:A:34:GLN:O	1:A:38:VAL:HG23	0.51	2.05	2	5
1:A:24:GLN:O	1:A:27:LYS:HG2	0.51	2.05	7	2
1:A:24:GLN:HE22	1:A:78:LYS:HB2	0.50	1.66	9	1
1:A:27:LYS:HE3	1:A:74:THR:OG1	0.49	2.06	6	1
1:A:54:ILE:HB	1:A:67:TRP:CE3	0.49	2.43	8	1
1:A:23:LEU:HB3	1:A:34:GLN:NE2	0.49	2.21	8	3
1:A:23:LEU:HD21	1:A:48:ALA:HB2	0.47	1.86	9	3
1:A:73:GLY:HA2	1:A:76:PHE:CE1	0.45	2.46	4	3
1:A:34:GLN:HG2	1:A:38:VAL:CG2	0.45	2.41	9	1
1:A:24:GLN:CD	1:A:78:LYS:HB2	0.45	2.32	9	1
1:A:34:GLN:CD	1:A:68:GLU:HA	0.45	2.33	1	1
1:A:41:ASN:HD22	1:A:44:LEU:HD12	0.44	1.73	8	1
1:A:24:GLN:HE21	1:A:74:THR:HG23	0.44	1.73	9	1
1:A:73:GLY:HA2	1:A:76:PHE:CZ	0.43	2.47	3	1
1:A:20:LEU:O	1:A:77:LEU:HD21	0.43	2.14	9	1
1:A:41:ASN:OD1	1:A:44:LEU:HB2	0.43	2.14	10	1
1:A:23:LEU:HD13	1:A:77:LEU:HD21	0.42	1.92	10	1
1:A:20:LEU:HA	1:A:77:LEU:HD11	0.42	1.91	8	2
1:A:16:MET:SD	1:A:16:MET:N	0.42	2.93	8	1
1:A:19:TYR:OH	1:A:51:ARG:HD3	0.41	2.15	8	2
1:A:38:VAL:HG23	1:A:45:VAL:HG21	0.41	1.92	6	1
1:A:39:ARG:NE	1:A:39:ARG:HA	0.41	2.31	2	1
1:A:26:ILE:HG21	1:A:34:GLN:CD	0.40	2.35	1	1
1:A:69:ILE:HG12	1:A:70:THR:O	0.40	2.17	3	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	52/69 (75%)	46±1 (88±2%)	5±1 (10±2%)	1±1 (2±1%)	10 50
All	All	520/690 (75%)	457 (88%)	52 (10%)	11 (2%)	10 50

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	35	SER	7
1	A	82	GLY	4

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	44/54 (81%)	37±2 (84±4%)	7±2 (16±4%)	5 42
All	All	440/540 (81%)	371 (84%)	69 (16%)	5 42

All 14 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	19	TYR	10
1	A	23	LEU	10
1	A	54	ILE	9
1	A	77	LEU	9
1	A	51	ARG	7
1	A	24	GLN	6
1	A	27	LYS	6
1	A	39	ARG	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	16	MET	2
1	A	55	SER	2
1	A	34	GLN	2
1	A	26	ILE	1
1	A	37	TYR	1
1	A	78	LYS	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 88% for the well-defined parts and 85% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *data1.str*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	759
Number of shifts mapped to atoms	759
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	67	-0.60 ± 0.09	Should be checked
$^{13}C_{\beta}$	60	0.13 ± 0.17	None needed (< 0.5 ppm)
$^{13}C'$	64	-0.55 ± 0.13	Should be applied
^{15}N	66	0.47 ± 0.27	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 88%, i.e. 638 atoms were assigned a chemical shift out of a possible 721. 0 out of 8 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	272/274 (99%)	111/112 (99%)	107/108 (99%)	54/54 (100%)
Sidechain	329/399 (82%)	225/261 (86%)	104/118 (88%)	0/20 (0%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	37/48 (77%)	19/23 (83%)	17/22 (77%)	1/3 (33%)
Overall	638/721 (88%)	355/396 (90%)	228/248 (92%)	55/77 (71%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 85%, i.e. 757 atoms were assigned a chemical shift out of a possible 889. 0 out of 8 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	336/350 (96%)	139/144 (97%)	131/138 (95%)	66/68 (97%)
Sidechain	376/481 (78%)	257/314 (82%)	119/142 (84%)	0/25 (0%)
Aromatic	45/58 (78%)	23/28 (82%)	21/27 (78%)	1/3 (33%)
Overall	757/889 (85%)	419/486 (86%)	271/307 (88%)	67/96 (70%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A: