



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 30, 2021 – 07:16 PM EST

PDB ID : 3LQ7
Title : Crystal structure of glutathione s-transferase from agrobacterium tumefaciens str. c58
Authors : Patskovsky, Y.; Toro, R.; Gilmore, M.; Chang, S.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYS-GXRC)
Deposited on : 2010-02-08
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.16
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	240	 7% (poor fit), 72% (0-1 outliers), 12% (2 outliers), 16% (3+ outliers or not modelled)
1	B	240	 15% (poor fit), 70% (0-1 outliers), 10% (2 outliers), 18% (3+ outliers or not modelled)
1	C	240	 6% (poor fit), 77% (0-1 outliers), 6% (2 outliers), 15% (3+ outliers or not modelled)

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 4945 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutathione S-transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	202	Total 1650	C 1056	N 300	O 289	S 5	0	3	0
1	B	197	Total 1608	C 1033	N 289	O 281	S 5	0	1	0
1	C	203	Total 1643	C 1049	N 297	O 291	S 6	0	1	0

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP A9CFJ9
A	0	SER	-	expression tag	UNP A9CFJ9
A	1	LEU	-	expression tag	UNP A9CFJ9
A	231	GLU	-	expression tag	UNP A9CFJ9
A	232	GLY	-	expression tag	UNP A9CFJ9
A	233	HIS	-	expression tag	UNP A9CFJ9
A	234	HIS	-	expression tag	UNP A9CFJ9
A	235	HIS	-	expression tag	UNP A9CFJ9
A	236	HIS	-	expression tag	UNP A9CFJ9
A	237	HIS	-	expression tag	UNP A9CFJ9
A	238	HIS	-	expression tag	UNP A9CFJ9
B	-1	MET	-	expression tag	UNP A9CFJ9
B	0	SER	-	expression tag	UNP A9CFJ9
B	1	LEU	-	expression tag	UNP A9CFJ9
B	231	GLU	-	expression tag	UNP A9CFJ9
B	232	GLY	-	expression tag	UNP A9CFJ9
B	233	HIS	-	expression tag	UNP A9CFJ9
B	234	HIS	-	expression tag	UNP A9CFJ9
B	235	HIS	-	expression tag	UNP A9CFJ9
B	236	HIS	-	expression tag	UNP A9CFJ9
B	237	HIS	-	expression tag	UNP A9CFJ9
B	238	HIS	-	expression tag	UNP A9CFJ9
C	-1	MET	-	expression tag	UNP A9CFJ9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	SER	-	expression tag	UNP A9CFJ9
C	1	LEU	-	expression tag	UNP A9CFJ9
C	231	GLU	-	expression tag	UNP A9CFJ9
C	232	GLY	-	expression tag	UNP A9CFJ9
C	233	HIS	-	expression tag	UNP A9CFJ9
C	234	HIS	-	expression tag	UNP A9CFJ9
C	235	HIS	-	expression tag	UNP A9CFJ9
C	236	HIS	-	expression tag	UNP A9CFJ9
C	237	HIS	-	expression tag	UNP A9CFJ9
C	238	HIS	-	expression tag	UNP A9CFJ9

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	15	Total O 15 15	0	0
2	B	21	Total O 21 21	0	0
2	C	8	Total O 8 8	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	54.92Å 111.25Å 246.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 38.44 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.7 (20.00-2.30) 98.6 (38.44-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.47 (at 2.31Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.243 , 0.288 0.245 , 0.291	Depositor DCC
R_{free} test set	1056 reflections (3.16%)	wwPDB-VP
Wilson B-factor (Å ²)	47.8	Xtrriage
Anisotropy	0.630	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 48.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4945	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/1700	0.58	0/2301
1	B	0.40	0/1650	0.57	0/2233
1	C	0.39	0/1685	0.56	0/2279
All	All	0.39	0/5035	0.57	0/6813

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1650	0	1647	23	0
1	B	1608	0	1606	16	0
1	C	1643	0	1635	5	0
2	A	15	0	0	0	0
2	B	21	0	0	0	0
2	C	8	0	0	0	0
All	All	4945	0	4888	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:34[B]:ARG:HH21	1:A:34[B]:ARG:CB	1.60	1.13
1:A:34[B]:ARG:NH2	1:A:34[B]:ARG:HB2	1.67	1.08
1:A:34[B]:ARG:HH21	1:A:34[B]:ARG:CG	1.70	1.03
1:A:34[B]:ARG:HH21	1:A:34[B]:ARG:HB2	1.16	0.99
1:A:34[B]:ARG:CB	1:A:34[B]:ARG:NH2	2.35	0.81
1:A:34[B]:ARG:HH21	1:A:34[B]:ARG:HG3	1.46	0.80
1:B:91:MET:HE1	1:B:101:LEU:HD11	1.69	0.75
1:A:34[B]:ARG:CG	1:A:34[B]:ARG:NH2	2.42	0.72
1:B:45:VAL:HG21	1:B:97:HIS:HB2	1.79	0.65
1:A:34[B]:ARG:NH2	1:A:34[B]:ARG:HG3	2.07	0.65
1:A:23:GLU:HB3	1:A:53:ARG:HB3	1.84	0.59
1:A:106:LEU:HD23	1:B:81:LEU:HB2	1.87	0.56
1:C:120:ASN:C	1:C:120:ASN:HD22	2.11	0.54
1:A:104:ASP:HB3	1:A:107:ARG:HH11	1.75	0.51
1:B:102:PRO:HB3	1:B:107:ARG:HH11	1.75	0.51
1:B:44:GLU:HG2	1:B:100:LEU:HD21	1.93	0.50
1:B:43:GLU:OE1	1:B:215[B]:ARG:NH1	2.46	0.49
1:A:120:ASN:HD22	1:A:120:ASN:C	2.17	0.48
1:A:104:ASP:HB3	1:A:107:ARG:HD3	1.95	0.48
1:A:132:VAL:HA	1:A:136:GLU:HB2	1.96	0.48
1:B:45:VAL:CG2	1:B:97:HIS:HB2	2.44	0.47
1:C:126:ILE:HG23	1:C:188:LEU:HD13	1.96	0.46
1:C:182:ILE:HD13	1:C:207:GLY:HA3	1.98	0.46
1:B:68:GLN:HE21	1:B:84:PHE:HB3	1.81	0.46
1:B:182:ILE:HD13	1:B:207:GLY:HA3	1.98	0.45
1:A:126:ILE:HG23	1:A:188:LEU:HD13	1.99	0.44
1:A:182:ILE:O	1:A:186:ARG:HG3	2.18	0.43
1:B:220:GLN:HE21	1:B:220:GLN:HB3	1.54	0.43
1:C:27:ASP:HB2	1:C:32:LEU:HB2	1.98	0.43
1:A:16:ASN:HA	1:A:17:PRO:HD3	1.90	0.43
1:A:94:ALA:HB1	1:A:101:LEU:HG	2.01	0.42
1:A:182:ILE:HD13	1:A:207:GLY:HA3	2.00	0.42
1:A:122:ILE:HD11	1:B:70:PHE:CZ	2.54	0.42
1:A:15:PRO:HB2	1:A:16:ASN:H	1.70	0.42
1:B:133:TRP:HD1	1:B:134:LEU:HG	1.84	0.42
1:A:81:LEU:HB2	1:B:106:LEU:HD22	2.01	0.41
1:B:76:TYR:HB3	1:B:83:LEU:HB2	2.03	0.41
1:A:70:PHE:CZ	1:B:122:ILE:HD11	2.56	0.41
1:B:120:ASN:C	1:B:120:ASN:HD22	2.24	0.41
1:C:20:THR:HG23	1:C:52:ARG:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	201/240 (84%)	197 (98%)	3 (2%)	1 (0%)	29	35
1	B	192/240 (80%)	188 (98%)	4 (2%)	0	100	100
1	C	200/240 (83%)	190 (95%)	10 (5%)	0	100	100
All	All	593/720 (82%)	575 (97%)	17 (3%)	1 (0%)	47	58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	85	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/201 (85%)	164 (96%)	7 (4%)	30	43
1	B	167/201 (83%)	158 (95%)	9 (5%)	22	30
1	C	170/201 (85%)	158 (93%)	12 (7%)	14	19
All	All	508/603 (84%)	480 (94%)	28 (6%)	21	30

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	SER
1	A	82	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	91	MET
1	A	110	ARG
1	A	120	ASN
1	A	148	ARG
1	A	200	LEU
1	B	45	VAL
1	B	53	ARG
1	B	91	MET
1	B	120	ASN
1	B	143	GLU
1	B	148	ARG
1	B	188	LEU
1	B	200	LEU
1	B	220	GLN
1	C	27	ASP
1	C	53	ARG
1	C	91	MET
1	C	103	GLU
1	C	110	ARG
1	C	120	ASN
1	C	138	ASN
1	C	143	GLU
1	C	154	LEU
1	C	188	LEU
1	C	193	ILE
1	C	220	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	120	ASN
1	A	152	GLN
1	B	68	GLN
1	B	72	GLN
1	B	120	ASN
1	B	128	ASN
1	B	220	GLN
1	C	68	GLN
1	C	120	ASN
1	C	128	ASN
1	C	138	ASN
1	C	152	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	220	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	202/240 (84%)	0.48	17 (8%) 11 15	37, 61, 98, 125	0
1	B	197/240 (82%)	0.91	35 (17%) 1 1	38, 65, 121, 138	0
1	C	203/240 (84%)	0.58	15 (7%) 14 19	41, 75, 109, 129	0
All	All	602/720 (83%)	0.65	67 (11%) 5 7	37, 67, 111, 138	0

All (67) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	224	PHE	11.6
1	B	54	LEU	7.3
1	B	22	PHE	6.5
1	B	223	VAL	6.2
1	A	106	LEU	6.0
1	C	54	LEU	5.1
1	B	52	ARG	4.8
1	A	24	ARG	4.8
1	C	67	TYR	4.8
1	A	55	SER	4.2
1	B	144	ALA	4.2
1	B	68	GLN	4.0
1	B	24	ARG	4.0
1	B	51	VAL	3.9
1	B	133	TRP	3.6
1	B	143	GLU	3.5
1	B	225	THR	3.5
1	B	142	HIS	3.5
1	C	24	ARG	3.4
1	A	56	PHE	3.4
1	B	53	ARG	3.3
1	B	146	LEU	3.2
1	A	54	LEU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	22	PHE	3.1
1	B	32	LEU	3.1
1	B	193	ILE	3.1
1	A	115	MET	3.0
1	B	141	TRP	3.0
1	B	138	ASN	3.0
1	C	66	ALA	2.9
1	C	22	PHE	2.9
1	A	79	GLY	2.8
1	C	106	LEU	2.8
1	A	226	ALA	2.8
1	C	73	ILE	2.7
1	A	227	ALA	2.7
1	B	135	PHE	2.7
1	B	106	LEU	2.7
1	C	96	HIS	2.7
1	B	23	GLU	2.5
1	B	137	ARG	2.5
1	B	87	GLY	2.4
1	C	166	ASP	2.4
1	B	113	ALA	2.3
1	B	119	LEU	2.3
1	B	115	MET	2.3
1	A	88	ALA	2.3
1	A	91	MET	2.3
1	C	144	ALA	2.3
1	C	100	LEU	2.2
1	B	21	VAL	2.2
1	B	227	ALA	2.2
1	A	26	PRO	2.2
1	C	221	LEU	2.1
1	A	166	ASP	2.1
1	B	80	ASP	2.1
1	A	23	GLU	2.1
1	B	129	PHE	2.1
1	B	86	SER	2.1
1	C	30	ARG	2.1
1	C	53	ARG	2.1
1	C	165	GLY	2.1
1	B	112	VAL	2.1
1	B	116	PHE	2.1
1	B	140	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	90	VAL	2.0
1	A	223	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.